arXiv:math/0112251v3 [math.RT] 9 Jan 2005

$\mathcal{L}$-MODULES AND MICRO-SUPPORT

LESLIE SAPER

Abstract. $\mathcal{L}$-modules are a combinatorial analogue of constructible sheaves on the reductive Borel-Serre compactification of a locally symmetric space. We define the *micro-support* of an $\mathcal{L}$-module; it is a set of irreducible modules for the Levi quotients of the parabolic $\mathbb{Q}$-subgroups associated to the strata. We prove a vanishing theorem for the global cohomology of an $\mathcal{L}$-module in term of the micro-support. We calculate the micro-support of the middle weight profile weighted cohomology and the middle perversity intersection cohomology $\mathcal{L}$-modules. (For intersection cohomology we must assume the $\mathbb{Q}$-root system has no component of type D_n, E_n, or F_4.) Finally we prove a functoriality theorem concerning the behavior of micro-support upon restriction of an $\mathcal{L}$-module to the pre-image of a Satake stratum. As an application we settle a conjecture made independently by Rapoport and by Goresky and MacPherson, namely, that the intersection cohomology (for either middle perversity) of the reductive Borel-Serre compactification of a Hermitian locally symmetric space is isomorphic to the intersection cohomology of the Baily-Borel-Satake compactification. We also obtain a new proof of the main result of Goresky, Harder, and MacPherson on weighted cohomology as well as generalizations of both of these results to general Satake compactifications with equal-rank real boundary components.

Contents

0. Introduction 2
Part I. $\mathcal{L}$-modules and Micro-support 13
1. Geodesic Action and Related Bundles 13
2. The Reductive Borel-Serre Compactification 16
3. $\mathcal{L}$-modules 18
4. Realization of $\mathcal{L}$-modules 23
5. Example: the Intersection Cohomology $\mathcal{L}$-module 27
6. Example: the Weighted Cohomology $\mathcal{L}$-module 30
7. Micro-support of an $\mathcal{L}$-module 31
8. Conjugate Self-contragredient Modules and Fundamental Weyl Elements 32

1991 *Mathematics Subject Classification.* Primary 11F75, 22E40, 32S60, 55N33; Secondary 14G35, 22E45.

Key words and phrases. Intersection cohomology, Shimura varieties, locally symmetric varieties, compactifications.

This research was supported in part by the National Science Foundation under grants DMS-8957216, DMS-9100383, and DMS-9870162, a grant from The Duke Endowment, an Alfred P. Sloan Research Fellowship, a grant from the Maximilian-Bickhoff-Stiftung to visit the Katholischen Universität Eichstätt as Hermann-Minkowski Gastprofessur, and a grant from the Institut des Hautes Études Scientifiques. The author wishes to thank these organizations for their hospitality and support.

The original manuscript was prepared with the $\mathcal{A}\mathcal{M}\mathcal{S}$-LaTeX macro system and the XY-pic package.

9. Alternate Forms of Definition 7.2(i) 35
Part II. A Global Vanishing Theorem for $\mathcal{L}$-modules 37
10. The Vanishing Theorem 37
11. L^2-cohomology with Boundary Values 39
12. L^2-cohomology of $\mathcal{L}$-modules 44
13. Analytic Lemmas 48
14. Vanishing Theorem for L^2-cohomology 51
15. Proof of Theorem 10.4 53
Part III. Micro-support Calculations 55
16. Micro-support of Weighted Cohomology 55
17. Micro-support of Intersection Cohomology 57
18. Proof of Proposition 17.7 60
19. Proof of Lemma 17.8 61
20. Proof of Theorem 17.9 64
Part IV. Satake Compactifications and Functoriality of Micro-support 66
21. Satake Compactifications 67
22. Functoriality under Inverse Images 71
23. Functoriality under Inverse Images (continued) 76
24. The Basic Lemma 78
25. The Basic Lemma in the Presence of Equal-Rank Real Boundary Components 80
26. Restriction to Fibers of $\pi\colon \widehat{X} \to X^*_\sigma$ 84
27. Application: the Conjecture of Rapoport and Goresky-MacPherson 85
References 86

0. INTRODUCTION

Let $X = \Gamma\backslash D$ be an arithmetic quotient of the symmetric space D associated to a connected reductive algebraic group G defined over $\mathbb{Q}$. (Here $D = G(\mathbb{R})/KA_G$, where $K \subseteq G(\mathbb{R})$ is a maximal compact subgroup, $A_G = S_G(\mathbb{R})^0$, and S_G is the maximal $\mathbb{Q}$-split torus in the center of G.) If X is not compact, Borel and Serre [11] construct a compact real analytic manifold with corners $\overline{X}$ whose interior is X. The faces of $\overline{X}$ are indexed by $\mathcal{P}$, the Γ-conjugacy classes of parabolic $\mathbb{Q}$-subgroups P of G. Each such face Y_P is fibered over an arithmetic quotient $X_P = \Gamma_{L_P}\backslash D_P$ of the symmetric space D_P associated to the Levi quotient L_P of P. The fibers are compact nilmanifolds (an arithmetic quotient of the real points of the unipotent radical of P) and Zucker [56] collapses these fibers (for all Y_P) to obtain a new compactification $\widehat{X}$ of X, the *reductive Borel-Serre compactification*.

The reductive Borel-Serre compactification is important because it is canonical, its singularities are reasonably easy to understand, and the more singular Satake compactifications X^* (such as the Baily-Borel-Satake compactification in the case that D is Hermitian symmetric) may be realized as quotients of it [57]. It would thus be desirable to transfer cohomological calculations from X^* to $\widehat{X}$. In this paper we develop a combinatorial tool, the theory of $\mathcal{L}$-modules and their micro-support, to study the cohomology of constructible complexes of sheaves on $\widehat{X}$. As one application, we settle a 15 year old conjecture on the intersection cohomology made independently by Rapoport [39] and by Goresky and MacPherson [24]. In fact we prove a generalization:

Theorem 27.1 (Rapoport/Goresky-MacPherson Conjecture)**.** *Let X^* be a Satake compactification of X for which all real boundary components $D_{R,h}$ are equal-rank, and let $\pi\colon \widehat{X} \to X^*$ be the projection from the reductive Borel-Serre compactification. Let E be a regular representation of G and let $\mathbb{E}$ be the corresponding locally constant sheaf on X. Let p be a middle perversity. Then there is a natural quasi-isomorphism of intersection cohomology sheaves $\pi_*\mathcal{I}_p\mathcal{C}(\widehat{X};\mathbb{E}) \cong \mathcal{I}_p\mathcal{C}(X^*;\mathbb{E})$.*

Recall that D is said to be *equal-rank* if $\mathbb{C}\text{-rank}\,{}^0G = \operatorname{rank} K$; here ${}^0G$ is the intersection of the kernels of the squares of all characters of G defined over $\mathbb{Q}$. The original conjecture was for D Hermitian symmetric and X^* the Baily-Borel-Satake compactification. Previously Zucker had observed that the conjecture held for $G = \mathrm{Sp}(4)$ with $\mathbb{E} = \mathbb{C}$. Furthermore, Goresky and MacPherson [24] announced a proof for $G = \mathrm{Sp}(4)$, $G = \mathrm{Sp}(6)$, and (for $\mathbb{E} = \mathbb{C}$) $G = \mathrm{Sp}(8)$. A proof in the case where $\mathbb{Q}\text{-rank}\,G = 1$ was given by Saper and Stern [40, Appendix].

The theory of $\mathcal{L}$-modules applies to all D and to sheaves other than intersection cohomology, so for future applications we work in a more general context and do not assume D to be Hermitian or equal-rank except as needed.

The paper is divided into four parts: I. $\mathcal{L}$-modules and Micro-support; II. A Global Vanishing Theorem for $\mathcal{L}$-modules; III. Micro-support Calculations; and IV. Satake Compactifications and Functoriality of Micro-support. After Part I, Parts II-IV may be read in any order provided one is willing to follow up on a few references; the last section of the paper, however, uses everything that precedes it. We now briefly outline the main results of the paper, after which we summarize the notation we will use; see also the semi-expository article [46].

0.1. **$\mathcal{L}$-modules on the Reductive Borel-Serre Compactification.** (§§1–6) The construction of the reductive Borel-Serre compactification $\widehat{X} = \coprod_P X_P$ is briefly indicated in §§1–2. Let i_P and $\hat{\imath}_P$ denote respectively the inclusion of a stratum X_P and its closure $\widehat{X}_P$ into $\widehat{X}$. Let $\mathbf{D}^b_{\mathcal{X}}(\widehat{X})$ be the derived category whose objects are bounded complexes of sheaves $\mathcal{S}$ on $\widehat{X}$ whose cohomology sheaves $\mathcal{H}(i_P^!\mathcal{S})$ supported on each stratum X_P are locally constant, say induced from graded representations $\mathcal{E}_P$ of Γ_{L_P}. We would like to consider such objects equipped with the extra structure that $\mathcal{E}_P$ is induced by restriction from a graded representation E_P of L_P.

In fact it is simpler to start with the family of graded representations E_P of L_P and construct from it an object of $\mathbf{D}^b_{\mathcal{X}}(\widehat{X})$. However we need something to "glue" these representations together. Consider two strata X_P and X_Q such that $X_P \subseteq \widehat{X}_Q$, the closure of X_Q; this means $P \subseteq Q$ (after perhaps replacing Q by a Γ-conjugate). Let $N_P \supseteq N_Q$ be the corresponding unipotent radicals. The topological link of the stratum X_P when intersected with X_Q is homotopically equivalent to an arithmetic quotient of $N_P^Q(\mathbb{R}) = N_P(\mathbb{R})/N_Q(\mathbb{R})$. A theorem of Nomizu and van Est states that the cohomology of this nilmanifold (with coefficients in $\mathbb{E}_Q$, the locally constant sheaf associated to E_Q) is isomorphic to the Lie algebra cohomology $H(\mathfrak{n}_P^Q; E_Q)$; furthermore this is an isomorphism of Γ_{L_P}-modules.

Thus in §3 we are led to consider combinatorial data $\mathcal{M} = (E_\cdot, f_{\cdot\cdot})$ consisting of a family of graded representations E_P of L_P, together with "dual attaching" morphisms $f_{PQ}\colon H(\mathfrak{n}_P^Q; E_Q) \xrightarrow{[1]} E_P$ for all $P \subseteq Q$. If $\mathcal{M}$ is required to satisfy a certain differential type condition (3.3.1) it is called an $\mathcal{L}$-*module*. From an $\mathcal{L}$-module $\mathcal{M}$ one can construct in a natural way an object $\mathcal{S}_{\widehat{X}}(\mathcal{M})$ of $\mathbf{D}^b_{\mathcal{X}}(\widehat{X})$ such that $\mathcal{H}(i_P^! \mathcal{S}_{\widehat{X}}(\mathcal{M})) = \mathbb{E}_P$.

In fact we may define a category of $\mathcal{L}$-modules $\mathfrak{Mod}(\mathcal{L}_W)$ for any locally closed union of strata W of $\widehat{X}$. If $k\colon W \hookrightarrow Z$ is an inclusion of such spaces, we have variants for $\mathcal{L}$-modules of the usual functors k_*, $k_!$, k^*, and $k^!$, as well as truncation functors by degree and by weight. In §4 we see there exists a realization functor $\mathcal{S}_W\colon \mathfrak{Mod}(\mathcal{L}_W) \to \mathbf{D}^b_{\mathcal{X}}(W)$ and via $\mathcal{S}_W$ the functors k_*, etc. on $\mathcal{L}$-modules correspond to the usual derived functors on $\mathbf{D}^b_{\mathcal{X}}(W)$. The point is that these derived functors preserve the extra structure.

There are a number of different incarnations of $\mathcal{S}_W(\mathcal{M})$ as a complex of sheaves. In the one we find simplest, a global section of $\mathcal{S}_W(\mathcal{M})$ corresponds to a family $\omega = (\omega_P)$, where ω_P is a special differential form [22] on $X_P \subseteq W$ with coefficients in $\mathbb{E}_P$. The differential on $\mathcal{S}_W(\mathcal{M})$ consists of the exterior derivative on each ω_P together with interaction terms based on the f_{PQ}.

If $\mathcal{M}$ is an $\mathcal{L}$-module on $\widehat{X}$, we define its cohomology $H(\widehat{X};\mathcal{M})$ to be the hypercohomology of the associated complex of sheaves $\mathcal{S}_{\widehat{X}}(\mathcal{M})$. The complexes of sheaves underlying many cohomology theories on $\widehat{X}$ can be lifted to $\mathcal{L}$-modules. For example, let E be a regular representation of G. Then there exist $\mathcal{L}$-modules $i_{G*}E$, $\mathcal{I}_p\mathcal{C}(E)$, and $\mathcal{W}^\eta\mathcal{C}(E)$ which correspond respectively to the ordinary cohomology $H(X;\mathbb{E}) = H(\Gamma; E)$, the intersection cohomology $I_pH(\widehat{X};\mathbb{E})$, and the weighted cohomology $W^\eta H(\widehat{X};\mathbb{E})$ (see [22]). The latter two $\mathcal{L}$-modules are constructed in §§5 and 6.

0.2. **Micro-support.** (§§7–9) To an $\mathcal{L}$-module $\mathcal{M}$ we associate its *micro-support* $\operatorname{SS}(\mathcal{M})$. The micro-support is a subset of the set of all irreducible representations of L_P for all $P \in \mathcal{P}$. As we will see below, the micro-support of $\mathcal{M}$ limits the range of degrees in which $H(\widehat{X};\mathcal{M})$ can be nonzero. (Micro-support here is vaguely analogous to the micro-support of a sheaf introduced by Kashiwara and Schapira [27], hence the name. There is no formal correspondence however.)

In order to define $\operatorname{SS}(\mathcal{M})$, let V be an irreducible L_P-module and let ξ_V denote the character by which the maximal $\mathbb{Q}$-split torus S_P in the center of L_P acts on V. Also let $V|_{M_P}$ denote the restriction of V to $M_P = {}^0L_P$, the natural complement to S_P. The closed strata $\widehat{X}_Q$ containing X_P correspond to subsets Δ_P^Q of Δ_P, the simple roots of S_P acting on the Lie algebra of the unipotent radical of P. Let $\widehat{X}_Q \supseteq X_P$ be a closed stratum such that

$$\{\, \alpha \mid (\xi_V + \rho, \alpha) < 0 \,\} \subseteq \Delta_P^Q \subseteq \{\, \alpha \mid (\xi_V + \rho, \alpha) \le 0 \,\}$$

where as usual ρ denotes one half the sum of the positive roots. Then V belongs to $\operatorname{SS}(\mathcal{M})$ if and only if

$$(V|_{M_P})^* \cong \overline{V|_{M_P}}, \text{ and} \tag{0.2.1}$$

$$H(i_P^* \hat{\imath}_Q^! \mathcal{M})_V \neq 0 \tag{0.2.2}$$

for some such Q.

0.3. **Vanishing Theorem for $\mathcal{L}$-modules.** (§§10–15) For V in $\mathrm{SS}(\mathcal{M})$, denote the least and greatest degrees for which (0.2.2) can be nonzero by $c(V;\mathcal{M}) \leq d(V;\mathcal{M})$. Set

$$c(\mathcal{M}) = \inf_{V\in\mathrm{SS}(\mathcal{M})} \tfrac{1}{2}(\dim D_P - \dim D_P(V)) + c(V;\mathcal{M}) \ , \text{ and}$$

$$d(\mathcal{M}) = \sup_{V\in\mathrm{SS}(\mathcal{M})} \tfrac{1}{2}(\dim D_P + \dim D_P(V)) + d(V;\mathcal{M}) \ ,$$

where $D_P(V)$ is the symmetric space associated to the reductive $\mathbb{R}$-subgroup of L_P whose roots (with respect to a fundamental Cartan subalgebra) are orthogonal to the highest weight of V. (This space depends on a θ-stable ordering; we choose one for which $\dim D_P(V)$ is maximized.) Then we have the

Theorem 10.4. *Let $\mathcal{M}$ be an $\mathcal{L}$-module on $\widehat{X}$. Then $H^i(\widehat{X};\mathcal{M}) = 0$ for $i \notin [c(\mathcal{M}), d(\mathcal{M})]$.*

The proof of the vanishing theorem proceeds by a combinatorial generalization of the analytic arguments from [47] together with a spectral sequence argument; this reduces the problem to Raghunathan's vanishing theorem for ordinary L^2-cohomology, Theorem 14.1. In §11 we recall ordinary L^2-cohomology and a version in which restriction to boundary faces is well defined. Now we need to represent $H(\widehat{X};\mathcal{M})$ by a variant of L^2-cohomology for our forms $(\omega_P) \in \mathcal{S}_{\widehat{X}}(\mathcal{M})$. However even though each $\widehat{X}_P$ is compact this is not entirely straightforward: the L^2-norm on ω_P with respect to the natural locally symmetric metric on $\widehat{X}_P$ is not appropriate because, being a complete metric on a noncompact space, it would impose L^2-growth conditions on ω_P which we do not want. The solution in §12 is to replace $\overline{X}$ by a naturally diffeomorphic compact domain $\overline{X}_t$ within X (this was constructed in [43]), and use the induced metric; we then work on the corresponding $\widehat{X}_t$ instead of $\widehat{X}$.

0.4. **Calculations of Micro-support.** (§§16–20) For simplicity, in this introduction we only give the results for the *essential micro-support* $\mathrm{SS}_{\mathrm{ess}}(\mathcal{M})$; this is the subset of $\mathrm{SS}(\mathcal{M})$ of V for which (0.2.2) is nonvanishing for all possible Q (in a compatible fashion). The essential micro-support determines $\mathrm{SS}(\mathcal{M})$ and may be used in calculating $c(\mathcal{M})$ and $d(\mathcal{M})$.

For weighted cohomology one can explicitly calculate $H(i_P^* \hat{\imath}_Q^! \mathcal{M})$ and one finds the

Theorem 16.3. *Let E be an irreducible regular G-module and let η be a middle weight profile. The micro-support is nonempty if and only if $(E|_{{}^0G})^* \cong \overline{E|_{{}^0G}}$; if this condition is satisfied then $\mathrm{SS}_{\mathrm{ess}}(\mathcal{W}^\eta\mathcal{C}(E)) = \{E\}$ and $c(E;\mathcal{W}^\eta\mathcal{C}(E)) = d(E;\mathcal{W}^\eta\mathcal{C}(E)) = 0$.*

The situation for intersection cohomology is far more delicate. We do not have an explicit closed formula for $H(i_P^* \hat{\imath}_Q^! \mathcal{M})$ (or even for the local cohomology $H(i_P^*\mathcal{M})$) and we are forced to use an inductive argument. However condition (0.2.1) is usually not preserved under passage to a larger parabolic $\mathbb{Q}$-subgroup. Nonetheless we have the following result:

Corollary 17.2. *Assume the irreducible components of the $\mathbb{Q}$-root system of G are of type A_n, B_n, C_n, BC_n, or G_2. Let E be an irreducible regular G-module and* *let p be a middle perversity. If* $(E|_{{}^0G})^* \cong \overline{E|_{{}^0G}}$, *then* $\mathrm{SS}_{\mathrm{ess}}(\mathcal{I}_p\mathcal{C}(E)) = \{E\}$ *with* $c(E;\mathcal{I}_p\mathcal{C}(E)) = d(E;\mathcal{I}_p\mathcal{C}(E)) = 0$.

Remarks. Unlike the situation for weighted cohomology, the micro-support can be nonempty even if $(E|_{{}^0G})^* \ncong \overline{E|_{{}^0G}}$; see Theorem 17.1 for details.

The hypothesis on the $\mathbb{Q}$-root system ensures that the the corresponding $\mathbb{Q}$-split group for each almost $\mathbb{Q}$-simple factor of G has a quasi-minuscule representation whose weights are linearly ordered (see the proof of Lemma 19.4). We conjecture the results on $\mathrm{SS}(\mathcal{I}_p\mathcal{C}(E))$ continue to hold without this hypothesis. The hypothesis is automatically satisfied if D is Hermitian or has a Satake compactification with all real boundary components equal-rank.

Finally the micro-support of $i_{G*}E$ is easy to calculate and is treated in [46]. As an application of the vanishing theorem it is proved in [46] that that if D is Hermitian or equal-rank and E has regular highest weight, then $H^i(X;\mathbb{E}) = 0$ for $i < \dim_{\mathbb{C}} X$. This answers a question posed by Tilouine.

0.5. **Satake Compactifications.** (§21) We recall the theory of Satake compactifications. Briefly, given a faithful irreducible finite dimensional representation $\sigma\colon G \to \mathrm{GL}(V)$, Satake [49], [50] constructed a compactification ${}_{\mathbb{R}}D^*_\sigma = \coprod_R D_{R,h}$ of D, where the strata $D_{R,h}$ are called the *real boundary components* and are indexed by their normalizers R. The union of just the so-called *rational boundary components* yields a space D^*_σ which, under certain conditions on σ, yields a compactification $X^*_\sigma = \Gamma\backslash D^*_\sigma$ of $X = \Gamma\backslash D$. The strata $F_R = \Gamma_{L_{R,h}}\backslash D_{R,h}$ are indexed by a subset $\mathcal{P}^*_\sigma \subseteq \mathcal{P}$. In the case that D is Hermitian symmetric, X^*_σ for a certain σ is the Baily-Borel-Satake compactification. It has a projective algebraic structure [2] and its strata are indexed by Γ-conjugacy classes of maximal parabolic $\mathbb{Q}$-subgroups.

Zucker showed [57] that there is a natural quotient map $\pi\colon \widehat{X} \to X^*_\sigma$. For each stratum X_P of $\widehat{X}$ there exists a stratum $F_{P^\dagger}$ of X^*_σ such that $\pi|_{X_P}\colon X_P \to F_{P^\dagger}$ is a flat bundle with typical fiber $X_{P,\ell}$. In fact, there is a connected normal $\mathbb{Q}$-subgroup $L_{P,\ell} \subseteq L_P$ so that $X_{P,\ell}$ is an arithmetic quotient of the symmetric space associated with $D_{P,\ell}$, while $F_{P^\dagger}$ is an arithmetic quotient of $L_{P,h} = L_P/L_{P,\ell}$. The inverse image $\pi^{-1}(F_R)$ of a stratum F_R of X^*_σ is the union of those X_P for which $P^\dagger = R$; we denote it $X_R(L_{R,\ell})$ since it is a partial compactification of X_R in the vertical ("$L_{R,\ell}$") directions. The restriction $\pi|_{X_R(L_{R,\ell})}\colon X_R(L_{R,\ell}) \to F_R$ is again a flat bundle with typical fiber $\widehat{X}_{R,\ell}$. Let $\hat{\imath}_{R,\ell}\colon \widehat{X}_{R,\ell} \hookrightarrow X_R(L_{R,\ell})$ denote the inclusion of a typical fiber. In §3.5 we define functors $\hat{\imath}^*_{R,\ell}$ and $\hat{\imath}^!_{R,\ell}$ on $\mathcal{L}$-modules; these commute with their analogues on the derived category.

0.6. **Functoriality of Micro-support.** (§§22–26) We need to understand how micro-support is affected when we apply functorial operations such as k^* or $k^!$ to $\mathcal{M}$. For brevity we only deal with k^* in this introduction. A few general results are presented in §22. For the case where k is a closed embedding, however, we only have a result for the *weak micro-support* $\mathrm{SS}_w(\mathcal{M})$: this is defined similarly to $\mathrm{SS}(\mathcal{M})$ but omits condition (0.2.1). Briefly if an irreducible L_P-module $V \in \mathrm{SS}_w(k^*\mathcal{M})$, then there exists an irreducible $L_{\tilde{P}}$-module $\tilde{V} \in \mathrm{SS}_w(\mathcal{M})$ such that V is an irreducible constituent of $H^\ell(\mathfrak{n}_P^{\tilde{P}};\tilde{V})$ with

$$(\xi_V + \rho)|_{\mathfrak{a}_P^{\tilde{P}}} \in {}^+\mathfrak{a}_P^{\tilde{P}*}\ , \tag{0.6.1}$$

where ${}^+\mathfrak{a}_P^{\tilde{P}*}$ is the real convex cone generated by $\Delta_P^{\tilde{P}}$.

The problem of $\mathrm{SS}_w(k^*\mathcal{M})$ versus $\mathrm{SS}(k^*\mathcal{M})$ is eliminated if k is the embedding of $X_R(L_{R,\ell})$ for a Satake compactification in which all rational boundary components are equal-rank; see §23. The point is that if $L_P = \widetilde{L_{P,h}}L_{P,\ell}$ is an almost direct product with the symmetric space of $\widetilde{L_{P,h}}$ being equal-rank, then $(V|_{\widetilde{L_{P,h}}})^* \cong \overline{V|_{\widetilde{L_{P,h}}}}$ is automatic.

We find that $c(k^*\mathcal{M})$ and $d(k^*\mathcal{M})$ may be estimated in terms of $c(\mathcal{M})$ and $d(\mathcal{M})$, together with the degrees ℓ from above. These are the lengths $\ell(w)$ of certain Weyl group elements w and (0.6.1) gives information on the geometry of the corresponding Weyl chamber. A basic lemma in §24 translates this geometry into an estimate on $\ell(w)$. In the case that $\#\Delta_P^{\tilde{P}} = 1$, the estimate is $\ell(w) \leq \frac{1}{2}\dim \mathfrak{n}_P^{\tilde{P}} - \mathfrak{n}_P(V)$. See §24.1 for the definition of $\mathfrak{n}_P(V)$; in §25 we estimate $\mathfrak{n}_P(V)$ in more geometric terms—this requires the stronger condition that all *real* boundary components are equal-rank.

The end result, after restricting to a typical fiber $\widehat{X}_{R,\ell} \subseteq X_R(L_{R,\ell})$, is

Corollary 26.2. *Let X_σ^* be a Satake compactification of X and assume that all the real boundary components $D_{R,h}$ of the associated Satake compactification ${}_{\mathbb{R}}D_\sigma^*$ are equal-rank. Let $\mathcal{M}$ be an $\mathcal{L}$-module with $\mathrm{SS}_{\mathrm{ess}}(\mathcal{M}) = \{E\}$ for some irreducible regular G-module E and $c(E;\mathcal{M}) = d(E;\mathcal{M}) = 0$. Then for every stratum F_R of X_σ^*,*

$$d(\hat{\imath}_{R,\ell}^*\mathcal{M}) \leq \frac{1}{2}\operatorname{codim} F_R - \dim \mathfrak{a}_R^G \qquad \text{and} \qquad c(\hat{\imath}_{R,\ell}^!\mathcal{M}) \geq \frac{1}{2}\operatorname{codim} F_R + \dim \mathfrak{a}_R^G.$$

0.7. **The Conjecture of Rapoport and Goresky-MacPherson.** (§27) Let $\mathcal{M}$ be an $\mathcal{L}$-module. We wish to identify $\pi_*\mathcal{S}_{\widehat{X}}(\mathcal{M})$ as an element of the derived category on X_σ^*. If $x \in F_R \subseteq X_\sigma^*$, then $\pi^{-1}(x) = \widehat{X}_{R,\ell}$, a fiber of $\pi|_{X_R(L_{R,\ell})}$. Thus the local cohomology at x of $\pi_*\mathcal{S}_{\widehat{X}}(\mathcal{M})$ is simply $H(\hat{\imath}_{R,\ell}^*\mathcal{M})$, and the local cohomology supported at x is $H(\hat{\imath}_{R,\ell}^!\mathcal{M})$. For $\mathcal{M} = \mathcal{I}_p\mathcal{C}(E)$, we consequently see from Theorem 10.4, Corollary 17.2, and Corollary 26.2 that $\pi_*\mathcal{S}_{\widehat{X}}(\mathcal{M})$ satisfies the local characterization of intersection cohomology. This proves the conjecture of Rapoport and Goresky-MacPherson. By replacing Corollary 17.2 with Theorem 16.3 we obtain the following generalization of the main theorem of [22]:

Theorem 27.2. *Under the assumptions of Theorem* 27.1*, let η be a middle weight profile and let p be a middle perversity. Then there is a natural quasi-isomorphism $\pi_*\mathcal{W}^\eta\mathcal{C}(\widehat{X};\mathbb{E}) \cong \mathcal{I}_p\mathcal{C}(X^*;\mathbb{E})$.*

0.8. **Work to be Done.** In [45] the theory of $\mathcal{L}$-modules is generalized to allow E_P to be a locally regular L_P-module and hence admit the possibility of non-Hausdorff local cohomology; this allowed us to realize L^2-cohomology as an $\mathcal{L}$-module and calculate its micro-support. There are many other developments and applications of the theory of $\mathcal{L}$-modules that would be interesting to pursue. We mention for example the following topics: $\mathbb{Q}$-structures on an $\mathcal{L}$-module and its cohomology (see §4.10); $\mathcal{L}$-modules on Satake compactifications (for which we use the family $(L_{R,h})_{R\in\mathcal{P}_\sigma^*}$ and an appropriate link cohomology functor—see §3.10); the removal of the hypothesis on the $\mathbb{Q}$-root system in Theorem 17.1; the action of Hecke correspondences on $\mathcal{L}$-modules and their cohomology; the "homotopy category" of $\mathcal{L}$-modules (see §3.9) and its localization with respect to the null system of $\mathcal{L}$-modules with empty micro-support; a "micro-support" spectral sequence for the cohomology of an $\mathcal{L}$-module; further investigation of the structure of $H(\Gamma; E)$ using $\mathcal{L}$-modules and the relation of this approach to that using Eisenstein series as initiated by Harder (see in particular the work of Franke [20]).

0.9. **Acknowledgments.** I would like to thank Mark Goresky for introducing me to this problem and for discussing it with me and Mark McConnell many times during the academic year 1988-89 while I visited Harvard University. I would also like to thank Michael Rapoport for reintroducing me to the problem in a letter to Stern and myself in summer 1991, and encouraging us to publish our proof of the $\mathbb{Q}$-rank 1 case. I am grateful for stimulating discussions with many people which aided me in this research, in particular, A. Borel, R. Bryant, R. Hain, M. Goresky, G. Harder, J.-P. Labesse, G. Lawler, M. Rapoport, J. Rohlfs, J. Schwermer, J. Tilouine, and N. Wallach. Finally I would like to thank the referee for many valuable suggestions and corrections, and in particular for suggesting the simplified proof of Proposition 16.2 that appears here.

0.10. **Notation.** We now summarize the notation that we will use throughout the paper.

0.10.1. The cardinality of a finite set T will be denoted $\#T$. For z in a group Z and $S \subseteq Z$ a subset, let ${}^{z}S = S^{z^{-1}}$ denote the conjugate zSz^{-1}. If Z is a topological space, the topological closure of a subset $S \subseteq Z$ will be denoted by $\mathrm{cl}(S)$. The notation id_Z indicates the identity map of Z.

0.10.2. If $\mathcal{C}$ is a category we will write $E \in \mathcal{C}$ to mean E is an object of $\mathcal{C}$. If $\mathcal{C}$ is an additive category, let $\mathbf{C}(\mathcal{C})$ denote the category of complexes of objects of $\mathcal{C}$ and let $\mathbf{Gr}(\mathcal{C})$ denote full subcategory of complexes with zero differential, that is, graded objects of $\mathcal{C}$. The full subcategory of $\mathbf{C}(\mathcal{C})$ or $\mathbf{Gr}(\mathcal{C})$ consisting of bounded complexes will be denoted with a superscript b.

Let $\mathcal{C}$ be an additive category. For $k \in \mathbb{Z}$, the *shift* $C[k]$ of an object $C \in \mathbf{C}(\mathcal{C})$ is defined by $C[k]^i = C^{i+k}$ and $d_{C[k]} = (-1)^k d_C$. Given a functor $F\colon \mathcal{C} \to \mathbf{C}(\mathcal{C}')$, where $\mathcal{C}$ and $\mathcal{C}'$ are additive categories, we shall often implicitly extend it to a functor $F\colon \mathbf{Gr}(\mathcal{C}) \to \mathbf{C}(\mathcal{C}')$ by setting $F(C) = \oplus_i F(C^i)[-i]$. Under this convention we have the equality $F(C)[k] = F(C[k])$. In a few instances we will further extend F to a functor $\mathbf{C}(\mathcal{C}) \to \mathbf{C}(\mathcal{C}')$ by taking the associated total complex.

0.10.3. For a topological space W, let $\mathfrak{Sh}(W)$ denote the category of sheaves of vector spaces over $\mathbb{C}$ on W. We write $\mathbf{C}(W)$ for $\mathbf{C}(\mathfrak{Sh}(W))$. Suppose W is equipped with a stratification $\mathcal{X}$. We will call a complex of sheaves $\mathcal{S}$ *constructible*[1] if for all strata $X_P \in \mathcal{X}$, the restriction $H(\mathcal{S})|_{X_P}$ of the local cohomology sheaves is locally constant. Let $\mathbf{C}_{\mathcal{X}}(W)$ denote the category of complexes of sheaves on W which are constructible. Let $\mathbf{D}_{\mathcal{X}}(W)$ be the associated derived category, obtained by formally inverting quasi-isomorphisms in the corresponding homotopy category. We denote the usual derived functors on $\mathbf{D}_{\mathcal{X}}(W)$ without the prefix "R", for example, i_* instead of Ri_*.

---

[1] In the terminology of [10, V, 3.3(ii)] this is $\mathcal{X}$-cohomologically locally constant, or $\mathcal{X}$-clc.

0.10.4. All references to manifolds, smooth differential forms, fiber bundles, locally constant sheaves, etc. should be taken in the sense of V-manifolds [48] (also called a orbifolds [54]).

For Y a manifold with (possibly empty) boundary or corners and $\mathbb{E}$ a locally constant sheaf on Y, let $\mathbb{E} \mapsto \mathcal{A}(Y;\mathbb{E})$ denote the de Rham resolution functor consisting of sheaves of smooth differential forms on Y with values in $\mathbb{E}$. The differential is exterior differentiation, denoted $d = d_Y = d_{Y,\mathbb{E}}$. Note that if $\mathbb{E}$ is graded we follow the sign convention of §0.10.2. The complex of global sections is denoted $A(Y;\mathbb{E})$ and the subcomplex of forms with compact support is denoted $A_c(Y;\mathbb{E})$.

0.10.5. For any algebraic group defined over $\mathbb{R}$, we denote the Lie algebra of its real points by the corresponding Gothic letter, for example, $\mathfrak{g} = \operatorname{Lie} G(\mathbb{R})$. The complexification is denoted by a subscript $\mathbb{C}$, for example $\mathfrak{g}_{\mathbb{C}}$. Now let Z be an algebraic group defined over $\mathbb{Q}$. The group of characters of Z defined over $\mathbb{Q}$ is denoted $X(Z)$. The one-dimensional representation space associated to a character χ is denoted $\mathbb{C}_\chi$. Let ${}^0Z = \bigcap_{\chi\in X(Z)} \operatorname{Ker}\chi^2$ be the intersection of the kernels of the squares of characters of Z. The group Z is an *almost direct product* of $\mathbb{Q}$-subgroups H and K if the product morphism $H \times K \to Z$ is surjective with finite kernel.

0.10.6. For a connected reductive $\mathbb{Q}$-group L, let $\mathfrak{Mod}(L)$ denote the category of regular representations of L. We write $\mathbf{C}(L)$ and $\mathbf{Gr}(L)$ instead of $\mathbf{C}(\mathfrak{Mod}(L))$ and $\mathbf{Gr}(\mathfrak{Mod}(L))$. If $M \subseteq L$ is a reductive $\mathbb{Q}$-subgroup and $E \in \mathfrak{Mod}(L)$, let $E|_M$ denote the regular representation of M obtained from E by restriction.

Let $\mathfrak{Irr}(L)$ denote the isomorphism classes of irreducible regular L-modules. For any $E \in \mathfrak{Mod}(L)$ we have an *isotypical decomposition*

$$E = \bigoplus_{V\in\mathfrak{Irr}(L)} E_V$$

where $E_V = \operatorname{Hom}_L(V,E) \otimes V$. For any irreducible representation $V \in \mathfrak{Irr}(L)$, we let ξ_V denote the character by which S_L acts on V. Since S_L is $\mathbb{Q}$-split, ξ_V is defined over $\mathbb{Q}$. Although ξ_V may not extend to a character of L, there exists a natural number k such that ξ_V^k extends to a character of L defined over $\mathbb{Q}$.

0.10.7. Let Z be an algebraic group and let $\sigma\colon Z \to \operatorname{GL}(V)$ be a regular representation. Recall the contragredient representation σ^* on V^* is defined by $(\sigma^*(g)\psi)(v) = \psi(\sigma(g^{-1})v)$. Define the complex vector space $\overline{V}$ to have vectors $\overline{v}$ indexed by $v \in V$, with operations $\overline{v}+\overline{w} = \overline{v+w}$ and $\lambda\overline{v} = \overline{\overline{\lambda}v}$. If Z is defined over $\mathbb{R}$ there is a complex conjugate representation $\overline{\sigma}$ on $\overline{V}$ defined by $\overline{\sigma}(g)\overline{v} = \overline{\sigma(\overline{g})v}$.

0.10.8. In this paper G will be a connected reductive algebraic group defined over $\mathbb{Q}$, and $\Gamma \subseteq G(\mathbb{Q})$ will be an arithmetic subgroup. (We briefly depart from this convention on G in §§21.2, 21.3.) We let D denote the associated symmetric space $G(\mathbb{R})/KA_G$, where K is a maximal compact subgroup of $G(\mathbb{R})$ and A_G is the connected component of the group of real points of the maximal $\mathbb{Q}$-split torus S_G in the center of G. Let $X = \Gamma\backslash D$ be the arithmetic quotient. If E is a regular representation of G, the induced locally constant sheaf on X will be denoted by the corresponding blackboard bold letter, thus $\mathbb{E} = D \times_\Gamma E$.

Since we do not assume Γ to be neat, X (and the various other arithmetic quotients that we will consider) need not be manifolds, but rather V-manifolds.

Alternatively the reader may assume that Γ is neat; any arithmetic group contains a neat subgroup with finite index.

0.10.9. Let P be a parabolic $\mathbb{Q}$-subgroup of G, with unipotent radical N_P and Levi quotient $L_P = P/N_P$. Set $M_P = {}^0L_P$. Let S_P denote the maximal $\mathbb{Q}$-split torus in the center of L_P and set $A_P = S_P(\mathbb{R})^0$. Restriction of characters yields an injective morphism $X(L_P) \to X(S_P)$ whose image has finite index. In terms of real points we have $L_P(\mathbb{R}) = M_P(\mathbb{R}) \times A_P$.

If $P \subseteq Q$ are parabolic $\mathbb{Q}$-subgroups, P/N_Q is a parabolic $\mathbb{Q}$-subgroup of L_Q with unipotent radical $N_P^Q = N_P/N_Q$. One can naturally identify S_Q as a subgroup of S_P and $X(L_Q)$ as a subgroup of $X(L_P)$. Namely consider the diagram

$$\begin{tikzcd} L_P & P/N_Q \arrow[l, two heads, "j"'] \arrow[r, hook, "i"] & L_Q \end{tikzcd} ;$$

since P/N_Q is parabolic in L_Q, i identifies its center with that of L_Q, while j identifies it with a subgroup of the center of L_P. Furthermore

$$\begin{tikzcd} X(L_P) \arrow[r, "\sim"] & X(P/N_Q) & X(L_Q) \arrow[l, hook', "i^*"'] \end{tikzcd} .$$

Set $S_P^Q = \left(\bigcap_{\chi \in X(L_Q)} \operatorname{Ker} \chi \cap S_P\right)^0$. Then S_P is an almost direct product

$$S_P = S_Q \cdot S_P^Q .$$

If we write $A_P^Q = S_P^Q(\mathbb{R})^0$, then A_P is an honest direct product

$$A_P = A_Q \times A_P^Q . \tag{0.10.1}$$

0.10.10. Let P be a parabolic $\mathbb{Q}$-subgroup of G. Let $\mathfrak{a}_P$ be the Lie algebra of A_P and let $\langle\ ,\ \rangle$ denote the natural pairing $\mathfrak{a}_P^* \times \mathfrak{a}_P \to \mathbb{R}$. We may identify $X(S_P) \otimes_{\mathbb{Z}} \mathbb{R} \cong \mathfrak{a}_P^*$. Consequently we will view elements of $X(S_P)$ both as characters on S_P and as linear functionals on $\mathfrak{a}_P$.

The group L_P acts via conjugation on N_P (and hence on $\mathfrak{n}_{P\mathbb{C}}$) via any lift $\widetilde{L_P} \subseteq P$. Although this action depends on the lift, the characters $\alpha \in X(S_P)$ by which S_P acts are independent of the lift. We denote these characters by $\Phi(\mathfrak{n}_P, S_P)$ or $\Phi(\mathfrak{n}_P, \mathfrak{a}_P)$. As usual we call them "roots" even though in general for P nonminimal they are not the positive roots of a root system. They do, however, have the property that every such root is a unique nonnegative integral linear combination of so-called simple roots; we let Δ_P denote these simple roots.

In the case that $P = P_0$ is a minimal parabolic $\mathbb{Q}$-subgroup, Δ_{P_0} is a basis of the $\mathbb{Q}$-root system ${}_{\mathbb{Q}}\Phi$ of G. In this case we omit the subscript P_0 and simply write Δ, S, $\mathfrak{a}$, etc.

0.10.11. If $P \subseteq Q$ are parabolic $\mathbb{Q}$-subgroups, the *type of Q with respect to P* is the subset Δ_P^Q of Δ_P consisting of roots restricting to 1 on S_Q. Parabolic $\mathbb{Q}$-subgroups Q containing P are in one-to-one correspondence with subsets of Δ_P via $Q \leftrightarrow \Delta_P^Q$.

Let

$$\mathfrak{a}_P = \mathfrak{a}_Q \oplus \mathfrak{a}_P^Q$$

be induced from (0.10.1). By §0.10.9 we have a natural inclusion $\mathfrak{a}_Q^* \hookrightarrow \mathfrak{a}_P^*$. Then $\mathfrak{a}_Q$ is the subspace of $\mathfrak{a}_P$ annihilated by Δ_P^Q, whereas $\mathfrak{a}_P^Q$ is the subspace annihilated by Δ_Q together with a basis of $X(S_G)$. We also have a dual decomposition

$$\mathfrak{a}_P^* = \mathfrak{a}_Q^* \oplus \mathfrak{a}_P^{Q*} .$$

Given $\alpha \in \mathfrak{a}_P^*$ we write $\alpha = \alpha_Q + \alpha^Q$ for its corresponding decomposition and likewise for elements of $\mathfrak{a}_P$.

0.10.12. For each $\gamma \in \Delta$, let $\gamma^\vee \in \mathfrak{a}$ be the corresponding coroot. (We shall occasionally identify $\mathfrak{a}$ with $\mathfrak{a}^*$ via a Weyl group invariant inner product $(\ ,\ )$, in which case $\gamma^\vee = \frac{2}{(\gamma,\gamma)}\gamma$.) To define coroots where P is an arbitrary parabolic $\mathbb{Q}$-subgroup, we proceed as in [1]. Namely for $\alpha \in \Delta_P$, let $\gamma \in \Delta \setminus \Delta^P$ be the unique root restricting to α on $\mathfrak{a}_P$ and let $\alpha^\vee = (\gamma^\vee)_P$ be the projection of $\gamma^\vee$ to $\mathfrak{a}_P$.

If $P \subseteq Q$, then $\mathfrak{a}_P^Q$ has a basis of coroots, $\{\alpha^\vee\}_{\alpha\in\Delta_P^Q}$. Let $\widehat{\Delta}_P^Q = \{\beta_\alpha^Q\}_{\alpha\in\Delta_P^Q}$ be the dual basis of $\mathfrak{a}_P^{Q*}$. On the other hand, Δ_P^Q is a basis for $\mathfrak{a}_P^{Q*}$. For $\beta \in \widehat{\Delta}_P^Q$, define $\beta^\vee \in \mathfrak{a}_P^Q$ by $\langle \alpha, \beta^\vee\rangle = \langle \beta, \alpha^\vee\rangle$ for all $\alpha \in \Delta_P^Q$. Then $\{\beta^\vee\}_{\beta\in\widehat{\Delta}_P^Q}$ is the basis of $\mathfrak{a}_P^Q$ dual to Δ_P^Q.

0.10.13. Let ${}^+\mathfrak{a}_P^*$ denote the real convex cone spanned by Δ_P and let $\mathfrak{a}_P^{*+}$ denote the real convex cone spanned by $\widehat{\Delta}_P$. Then $\mathfrak{a}_P^{*+}$ is the dominant cone

$$\{\, \mu \in \mathfrak{a}_P^* \mid \langle \mu, \alpha^\vee\rangle \geq 0 \text{ for all } \alpha \in \Delta_P \,\}$$

and $\mathfrak{a}_P^{*+} \subseteq {}^+\mathfrak{a}_P^*$. Similarly define ${}^+\mathfrak{a}_P^{Q*}$ and $\mathfrak{a}_P^{Q*+}$.

0.10.14. Beginning in §21 we will need to consider parabolic $\mathbb{R}$-subgroups. The discussion in §§0.10.9–0.10.13 applies with $\mathbb{Q}$ replaced by $\mathbb{R}$. We will add a left subscript $\mathbb{R}$ to the notation in this case, for example, ${}_{\mathbb{R}}S_P$, ${}_{\mathbb{R}}A_P$, ${}_{\mathbb{R}}\Delta_P$, etc.

0.10.15. Let $P \subseteq Q$ be parabolic $\mathbb{Q}$-subgroups. Let (P, Q) denote the parabolic $\mathbb{Q}$-subgroup "complementary" to Q with respect to P with type $\Delta_P \setminus \Delta_P^Q$. When (P, Q) occurs as a subscript or a superscript, we omit the parentheses, for example, $\Delta_P^{P,Q}$. Thus we have a disjoint decomposition

$$\Delta_P = \Delta_P^{P,Q} \amalg \Delta_P^Q \ .$$

Restriction to $\mathfrak{a}_Q$ sends Δ_P^Q to zero and defines a bijection $\Delta_P^{P,Q} \xrightarrow{\sim} \Delta_Q$.

Note that this decomposition of Δ_P induces a direct sum decomposition

$$\mathfrak{a}_P = \mathfrak{a}_Q \oplus \mathfrak{a}_{P,Q}^G \ ,$$

however in general $\mathfrak{a}_{P,Q}^G \neq \mathfrak{a}_P^Q$.

0.10.16. Let

$$\mathcal{P} = \{\, \Gamma\text{-conjugacy classes of parabolic } \mathbb{Q}\text{-subgroups of } G \,\} \ ;$$

notationally we do not distinguish a parabolic $\mathbb{Q}$-subgroup from its Γ-conjugacy class. Define a partial order on $\mathcal{P}$ by

$$P \leq Q \quad \iff \quad P \subseteq Q^\gamma \text{ for some } \gamma \in \Gamma.$$

(When $P \leq Q$ we will always feel free to replace Q by a Γ-conjugate so as to arrange $P \subseteq Q$ as needed.) For P and Q in $\mathcal{P}$, let $P \vee Q$ denote the least upper bound of P and Q; let $P \cap Q$ denote the greatest lower bound when it exists.

Given $P \leq R$, let $[P, R]$ denote the interval $\{\, Q \in \mathcal{P} \mid P \leq Q \leq R \,\}$. This is a complemented lattice: for every $Q \in [P, R]$ there exists $Q' \in [P, R]$ such that

$Q \cap Q' = P$ and $Q \vee Q' = R$, namely $Q' = (P, Q) \cap R$. We represent this situation by the diagram

$$\begin{tikzcd} & R \arrow[dl, dash] \arrow[dr, dash] & \\ Q' \arrow[dr, dash] & & Q \arrow[dl, dash] \\ & P & \end{tikzcd} \quad . \tag{0.10.2}$$

0.10.17. Let P be a parabolic $\mathbb{Q}$-subgroup and let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{l}_P$. Write $\mathfrak{h} = \mathfrak{b}_P \oplus \mathfrak{a}_P$ where $\mathfrak{b}_P$ is a Cartan subalgebra of $\mathfrak{m}_P$. (In §25 we shall also need $\mathfrak{h} = {}_{\mathbb{R}}\mathfrak{b}_P \oplus {}_{\mathbb{R}}\mathfrak{a}_P$ for P a parabolic $\mathbb{R}$-subgroup, where ${}_{\mathbb{R}}\mathfrak{b}_P$ is a Cartan subalgebra of ${}_{\mathbb{R}}\mathfrak{m}_P$.) The corresponding roots will be denoted $\Phi(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ and a positive system will be denoted $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. (In the case $Q = G$ we shall often simply write Φ and Φ^+.)

Via a lift, $\mathfrak{h}$ acts on $\mathfrak{n}_P$ via the adjoint action; the corresponding roots are independent of the lift and will be denoted $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{P\mathbb{C}})$. Likewise via a lift, $\mathfrak{h}$ may be considered as a Cartan subalgebra of $\mathfrak{l}_Q$ for any parabolic $\mathbb{Q}$-subgroup $Q \supseteq P$; the decomposition $\mathfrak{h} = \mathfrak{b}_Q + \mathfrak{a}_Q$ and the roots $\Phi(\mathfrak{l}_{Q\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ are independent of the lift. We can always choose a positive system $\Phi^+(\mathfrak{l}_{Q\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ containing $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ and $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{P\mathbb{C}})$.

0.10.18. Let $\rho = \rho^P + \rho_P$ be one-half the sum of roots in $\Phi^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$, where ρ^P (resp. ρ_P) is one-half the sum of the roots in $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ (resp. $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$). We may identify $2\rho_P$ with the $\mathbb{Q}$-rational character of L_P by which L_P (via any lift $\widetilde{L}_P$) acts on $\bigwedge^{\dim \mathfrak{n}_P} \mathfrak{n}_P$. Thus ρ_P is supported on $\mathfrak{a}_P$ and $\rho = \rho^P + \rho_P$ corresponds to the decomposition induced by restriction to the summands in $\mathfrak{h} = \mathfrak{b}_P \oplus \mathfrak{a}_P$.

0.10.19. Let P be a parabolic $\mathbb{Q}$-subgroup and let E be a regular G-module. The *nilpotent Lie algebra cohomology functor* $E \mapsto H(\mathfrak{n}_P; E)$ from $\mathfrak{Mod}(G)$ to $\mathbf{Gr}(L_P)$ will play a crucial role in what follows. As a vector space, $H(\mathfrak{n}_P; E)$ is defined as the cohomology of the complex $C(\mathfrak{n}_P; E) = \operatorname{Hom}(\bigwedge \mathfrak{n}_P, E)$ equipped with the usual differential (see for example [13, I, §1.1]). The coadjoint representation induces an action of P on $C(\mathfrak{n}_P; E)$ which commutes with the differential, and hence there is an action of P on $H(\mathfrak{n}_P; E)$. The unipotent radical N_P acts trivially [13, I, §1.1(5)] so this descends to a representation of L_P. Finally, a morphism of G-modules $f \colon E \to E'$ induces a morphism of P-modules $C(\mathfrak{n}_P; E) \to C(\mathfrak{n}_P; E')$ which commutes with the differentials and hence induces an L_P-module morphism $H(\mathfrak{n}_P; E) \to H(\mathfrak{n}_P; E')$ which we denote $H(\mathfrak{n}_P; f)$.

0.10.20. Let W denote the Weyl group of Φ. For $w \in W$ let $\ell(w)$ denote the length of $w \in W$ with respect to the reflections in simple roots and set $\Phi_w = \{\, \gamma \in \Phi^+ \mid w^{-1}\gamma < 0 \,\}$. It is well known that $\ell(w) = \#\Phi_w$.

For a parabolic $\mathbb{Q}$-subgroup P let $W^P \subseteq W$ be the subgroup corresponding to the Weyl group of $\Phi(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ and let $W_P \subseteq W$ be the set of minimal length coset representatives of $W^P \backslash W$.[2] We have $w \in W_P$ if and only if $\Phi_w \subseteq \Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. There is a factorization

$$W = W^P W_P.$$

---

[2] Our use of subscript versus superscript here is opposite from the usual convention, but this way is more compatible with the subscripts and superscripts on A and Δ.

The corresponding groups and coset representatives for the $\mathbb{Q}$-root system will be used in §19, and will be denoted there ${}_{\mathbb{Q}}W$, etc.

0.10.21. *Kostant's Theorem.* Let E be a regular G-module. The complement W_P plays an important role in Kostant's theorem [31], which states that $H(\mathfrak{n}_P; E)$ has a decomposition as an L_P-module indexed by W_P:

$$H(\mathfrak{n}_P; E) = \bigoplus_{w \in W_P} H(\mathfrak{n}_P; E)_w \ .$$

Furthermore, $H(\mathfrak{n}_P; E)_w$ is concentrated in degree $\ell(w)$ and, if E is irreducible with highest weight λ, then $H(\mathfrak{n}_P; E)_w$ is the irreducible L_P-module having highest weight $w(\lambda + \rho) - \rho$.

0.10.22. Let $P \subseteq Q$ be parabolic $\mathbb{Q}$-subgroups. We have

$$W = W^Q W_Q = (W^P W_P^Q) W_Q$$

and one may check that

$$W_P = W_P^Q W_Q.$$

For $w \in W_P$ write $w = w^Q w_Q$ accordingly. We have the important equality

$$\ell(w) = \ell(w^Q) + \ell(w_Q)$$

and in fact

$$\Phi_w = \Phi_{w^Q} \coprod w^Q \Phi_{w_Q} \ . \tag{0.10.3}$$

It immediately follows from Kostant's theorem that

$$H(\mathfrak{n}_P; E)_w \cong H(\mathfrak{n}_P^Q; H(\mathfrak{n}_Q; E)_{w_Q})_{w^Q}$$

and consequently [22, (11.8)], [51, 4.10]

$$H(\mathfrak{n}_P; E) \cong H(\mathfrak{n}_P^Q; H(\mathfrak{n}_Q; E)) \ . \tag{0.10.4}$$

Part I. $\mathcal{L}$-modules and Micro-support

1. Geodesic Action and Related Bundles

Recall that D is the symmetric space $G(\mathbb{R})/KA_G$ associated to G and $X = \Gamma\backslash D$ is an arithmetic quotient. In this section we briefly recall the geodesic retraction and nilmanifold fibration associated to a parabolic $\mathbb{Q}$-subgroup P [11, §3], [56, §3], [22, §§3, 4, 7]. This section also serves to establish our notation for the various spaces and bundles that will enter into the construction of the reductive Borel-Serre compactification in §2. One novelty of our presentation is that we introduce a "geodesic action" for the entire Levi quotient $L_P(\mathbb{R})$ as opposed to merely A_P; this is convenient for describing the flat connection on the nilmanifold fibration.

1.1. **Geodesic action.** Given any basepoint $x \in D$, let $L_{P,x}$ denote the unique lift of L_P to P which is stable under the Cartan involution associated to x. Since $P(\mathbb{R})$ acts transitively on D, we may express $y \in D$ as px for $p \in P(\mathbb{R})$; since $P = N_P \rtimes L_{P,x}$ we may write $p = ur$ with $u \in N_P(\mathbb{R})$ and $r \in L_{P,x}(\mathbb{R})$. Define the *geodesic action* of $L_P(\mathbb{R})$ on D by

$$z \mathbf{o}\, y = u z_x r x, \tag{1.1.1}$$

where $z_x \in L_{P,x}(\mathbb{R})$ denotes the lift of z. Note that at the basepoint x the geodesic action of an element of L_P agrees with the ordinary action of its lift in $L_{P,x}$. Clearly the restriction of this action to A_P agrees with the geodesic action as defined in [11].

Lemma. *The geodesic action $(z, y) \mapsto z \mathbf{o}\, y$ of $L_P(\mathbb{R})$ on D is well-defined* (*that is, $z \mathbf{o}\, y$ is independent of the choice of p and x above*). *It satisfies*

$$q(z \mathbf{o}\, y) = {}^q z \mathbf{o}\, qy \qquad (q \in P(\mathbb{R})), \tag{1.1.2}$$

where $z \mapsto {}^q z$ denotes the action of $P(\mathbb{R})$ by conjugation on $L_P(\mathbb{R})$.

Proof. Consider first the choice of p. If $y = p'x$ for $p' \in P(\mathbb{R})$, then $p' = pk$ for some $k \in K_x \cap P \subseteq L_{P,x}$ and hence $p' = ur'$ with $r' = rk$. Thus $uz_x r' x = u z_x r x$ and so the right hand side of (1.1.1) depends only on x, y, and z.

Now consider another basepoint x'. Write $x' = qx = vlx$ with $v \in N_P(\mathbb{R})$ and $l \in L_{P,x}(\mathbb{R})$. Then $L_{P,x'} = v L_{P,x} v^{-1}$. We may express $y = p'x'$ where $p' = pq^{-1} = url^{-1}v^{-1} = (uv^{-1})(vrl^{-1}v^{-1}) = u'r'$. Thus we compute $u' z_{x'} r' x' = (uv^{-1})(v z_x v^{-1})(vrl^{-1}v^{-1}) qx = u z_x r x$, which shows that (1.1.1) is independent of x.

To prove (1.1.2), calculate $q(z \mathbf{o}\, y) = q u z_x r x = {}^q u\, {}^q(z_x)\, {}^q r\, qx$. However ${}^q(z_x)$ is the lift of ${}^q z$ to $L_{P,x'}$, where $x' = qx$. Thus $q(z \mathbf{o}\, y) = {}^q u ({}^q z)_{x'}\, {}^q r x' = {}^q z \mathbf{o}\, ({}^q u\, {}^q r x') = {}^q z \mathbf{o}\, qy$. ☐

Corollary. *The geodesic action of $L_P(\mathbb{R})$ and the usual action of $N_P(\mathbb{R})$ commute. The geodesic action of $A_P \subseteq L_P(\mathbb{R})$ commutes with the usual action of $P(\mathbb{R})$.*

Proof. By the lemma we need to show that ${}^q z = z$ for $q \in N_P(\mathbb{R})$ and $z \in L_P(\mathbb{R})$ (and likewise for $q \in P(\mathbb{R})$ and $z \in A_P$). Write $z = z_x N_P(\mathbb{R})$ for some basepoint x. Then ${}^q z = q z_x q^{-1} N_P(\mathbb{R}) = z_x q^{z_x} q^{-1} N_P(\mathbb{R})$, so we need to show that $q^{z_x} q^{-1} \in N_P(\mathbb{R})$. This follows in the first case since $q \in N_P(\mathbb{R})$ and $N_P(\mathbb{R})$ is normal. In the second case, write $q = vl$ with $v \in N_P(\mathbb{R})$ and $l \in L_{P,x}(\mathbb{R})$. Then $q^{z_x} q^{-1} = v^{z_x} l^{z_x} l^{-1} v^{-1} = v^{z_x} v^{-1}$, since $z_x \in A_{P,x}$ is in the center of $L_P(\mathbb{R})$. Now proceed as in the first case. ☐

1.2. **Comparison of geodesic actions.** Consider now two parabolic $\mathbb{Q}$-subgroups $P \subseteq Q$; as in §0.10.9 the split component A_Q may be viewed as a subgroup of A_P. Thus it has a geodesic action viewed as a subgroup of either $L_Q(\mathbb{R})$ or $L_P(\mathbb{R})$; these two geodesic actions agree [11, 3.10].

For the extended geodesic action we have the

Lemma. *The geodesic action of $N_P^Q(\mathbb{R}) = N_P(\mathbb{R})/N_Q(\mathbb{R}) \subseteq L_Q(\mathbb{R})$ commutes with the geodesic action of $L_P(\mathbb{R})$.*

Proof. Write $y \in D$ as $y = urx$, where $u \in N_P(\mathbb{R})$, $r \in L_{P,x}(\mathbb{R})$. Since $N_P \subseteq Q = N_Q \rtimes L_{Q,x}$ we can decompose $N_P = N_Q \rtimes N_{P,x}^Q$ where $N_{P,x}^Q \subseteq L_{Q,x}$ is the lift of

N_P^Q. Thus we can write $u = vw$, where $v \in N_Q(\mathbb{R})$ and $w \in N_{P,x}^Q(\mathbb{R})$. The geodesic action of $n \in N_P^Q(\mathbb{R})$ is now $n \mathbin{\mathbf{o}} y = vn_x wrx$, which clearly commutes with the geodesic action of $z \in L_P(\mathbb{R})$, $z \mathbin{\mathbf{o}} y = uz_x rx = vwz_x rx$. $\square$

1.3. **Geodesic retraction.** By Corollary 1.1 the geodesic action of A_P and the ordinary action of ${}^0P(\mathbb{R})$ commute. The consequent action of $A_P \times {}^0P(\mathbb{R})$ on D is transitive and the stabilizer of any point x is $\{A_G\} \times {}^0P(\mathbb{R}) \cap K_x$, where $K_x \subseteq G(\mathbb{R})$ is the maximal compact subgroup fixing x. Set[3] $\mathcal{A}_P^G = {}^0P(\mathbb{R})\backslash D$ and $e_P = A_P\backslash D$. The preceding implies there is an isomorphism

$$D \cong \mathcal{A}_P^G \times e_P \tag{1.3.1}$$

of $(A_P^G \times {}^0P(\mathbb{R}))$-homogeneous spaces. The quotient map

$$D \longrightarrow e_P = A_P\backslash D$$

obtained by collapsing the orbits of the geodesic action of A_P is a trivial principal A_P^G-bundle with canonical trivializing sections given by the orbits of ${}^0P(\mathbb{R})$. Note that after choosing a basepoint $x \in D$ and hence $\bar{x} \in \mathcal{A}_P^G$, there is an isomorphism of A_P^G-homogeneous spaces $A_P^G \xrightarrow{\sim} \mathcal{A}_P^G$ given by $a \mapsto a \mathbin{\mathbf{o}} \bar{x}$. Thus $\mathcal{A}_P^G$ is an affine version of A_P^G.

The action of $\Gamma_P = \Gamma \cap P$ on $D \cong \mathcal{A}_P^G \times e_P$ operates only on the second factor, so the principal A_P^G-bundle structure persists after taking the quotient:

$$r_P \colon \Gamma_P\backslash D \longrightarrow Y_P = \Gamma_P\backslash e_P.$$

The map r_P is called *geodesic retraction.* Since $\Gamma_P \subset {}^0P(\mathbb{R})$, the quotients by Γ_P of the orbits of ${}^0P(\mathbb{R})$ still yield canonical trivializing sections of r_P.

If $P \subseteq Q$ this map descends to yield a canonically trivial A_P^Q-bundle (also called geodesic retraction)

$$r_P \colon \Gamma_P\backslash e_Q \longrightarrow Y_P.$$

1.4. **Nilmanifold fibrations.** The geodesic action of ${}^0L_P(\mathbb{R})$ descends to e_P and by Corollary 1.1 this action commutes with the $N_P(\mathbb{R})$-action. The resulting action of $N_P(\mathbb{R}) \times {}^0L_P(\mathbb{R})$ on e_P is transitive and the stabilizer of any point is contained in $\{1\} \times {}^0L_P(\mathbb{R})$. Set $\mathcal{N}_P(\mathbb{R}) = {}^0L_P(\mathbb{R})\backslash e_P$ and $D_P = N_P(\mathbb{R})\backslash e_P$; the space $\mathcal{N}_P(\mathbb{R})$ is an affine version of $N_P(\mathbb{R})$ and D_P is the generalized symmetric space associated to L_P. Then there is an isomorphism

$$e_P \cong \mathcal{N}_P(\mathbb{R}) \times D_P$$

of $(N_P(\mathbb{R}) \times {}^0L_P(\mathbb{R}))$-homogeneous spaces. The principal $N_P(\mathbb{R})$-bundle

$$\mu \colon e_P \longrightarrow D_P$$

obtained by collapsing the orbits of the action of $N_P(\mathbb{R})$ has canonical trivializing sections given by the orbits of the geodesic action of ${}^0L_P(\mathbb{R})$. Given a basepoint $x \in D$ and hence $\bar{x} \in \mathcal{N}_P(\mathbb{R})$, there is an isomorphism $n_x \colon N_P(\mathbb{R}) \xrightarrow{\sim} \mathcal{N}_P(\mathbb{R})$ of $N_P(\mathbb{R})$-homogeneous spaces defined by $n_x(u) = u\bar{x}$.

The action of ${}^0P(\mathbb{R})$ on e_P descends to actions on both $\mathcal{N}_P(\mathbb{R})$ and D_P, and the isomorphism $e_P \cong \mathcal{N}_P(\mathbb{R}) \times D_P$ is an isomorphism of ${}^0P(\mathbb{R})$-homogeneous spaces. Set $\mathcal{N}_P(\mathbb{R})' = \Gamma_{N_P}\backslash\mathcal{N}_P(\mathbb{R})$ and note that the action of Γ_P on $\mathcal{N}_P(\mathbb{R})$ descends to an action of $\Gamma_{L_P} = \Gamma_P/\Gamma_{N_P}$ on $\mathcal{N}_P(\mathbb{R})'$. The previous decomposition descends to an isomorphism $\Gamma_{N_P}\backslash e_P \cong \mathcal{N}_P(\mathbb{R})' \times D_P$ of Γ_{L_P}-homogeneous spaces and thus there is a canonically trivial $\mathcal{N}_P(\mathbb{R})'$-bundle

---

[3] In [11] e_P is denoted $e(P)$.

$$\mu' \colon \Gamma_{N_P}\backslash e_P \to D_P \ .$$

Set[4] $N_P(\mathbb{R})' = \Gamma_{N_P}\backslash N_P(\mathbb{R})$. Given a basepoint $x \in D$, the isomorphism n_x descends to $n'_x \colon N_P(\mathbb{R})' \xrightarrow{\sim} \mathcal{N}_P(\mathbb{R})'$.

Finally take the quotient by Γ_{L_P} of μ'; we obtain a flat bundle (with fibers noncanonically diffeomorphic to $\mathcal{N}_P(\mathbb{R})'$)

$$p \colon Y_P \longrightarrow X_P = \Gamma_{L_P}\backslash D_P.$$

This is simply the flat bundle $D_P \times_{\Gamma_{L_P}} \mathcal{N}_P(\mathbb{R})' \to X_P$ associated to the Γ_{L_P}-space $\mathcal{N}_P(\mathbb{R})'$. It is called the *nilmanifold fibration*.

1.5. For later use, we calculate [22, (7.8)] the action of Γ_{L_P} on $N_P(\mathbb{R})'$ that is needed in order to make $n'_x \colon N_P(\mathbb{R})' \xrightarrow{\sim} \mathcal{N}_P(\mathbb{R})'$ an isomorphism of Γ_{L_P}-spaces. Let $\gamma\Gamma_{N_P} \in \Gamma_{L_P}$ and let $\Gamma_{N_P}u \in N_P(\mathbb{R})'$; write $\gamma = vl$ where $v \in N_P(\mathbb{R})$ and $l \in L_{P,x}(\mathbb{R})$. Then

$$\gamma\Gamma_{N_P} \cdot n'_x(\Gamma_{N_P}u) = \Gamma_{N_P}\gamma.u.\bar{x} = \Gamma_{N_P}v.l.u.\bar{x} = \Gamma_{N_P}v.{}^{l}u.\bar{x} = n'_x(\Gamma_{N_P}v.{}^{l}u) \ .$$

Thus the induced action of γ on $N_P(\mathbb{R})'$ is $\Gamma_{N_P}u \mapsto \Gamma_{N_P}v.{}^{l}u = \Gamma_{N_P}{}^{\gamma}u.v$.

2. The Reductive Borel-Serre Compactification

Let $X = \Gamma\backslash D$ be a locally symmetric space. We describe the Borel-Serre compactification $\overline{X}$ [11] and its quotient introduced by Zucker [56], the reductive Borel-Serre compactification $\widehat{X}$ [22, §8].

2.1. **The Borel-Serre compactification.** Let P be a parabolic $\mathbb{Q}$-subgroup of G. There is an isomorphism $A_P^G \xrightarrow{\sim} (\mathbb{R}^{>0})^{\Delta_P}$ defined by $a \mapsto (a^\alpha)_{\alpha\in\Delta_P}$ and thus A_P^G may be embedded into a semigroup $\bar{A}_P^G \xrightarrow{\sim} (\mathbb{R}^{>0} \cup \{\infty\})^{\Delta_P}$.

The orbits of the action of A_P^G on $\bar{A}_P^G$ yield a stratification indexed by the parabolic $\mathbb{Q}$-subgroups $Q \supseteq P$. Namely, let $o_Q \in \bar{A}_P^G$ be defined by $\alpha(o_Q) = \infty$ for $\alpha \in \Delta_P^{P,Q}$, and $\alpha(o_Q) = 1$ for $\alpha \in \Delta_P^Q$. Then we have the stratification

$$\bar{A}_P^G = \coprod_{Q\supseteq P} A_P^G \cdot o_Q. \tag{2.1.1}$$

Recall that the quotient map $D \to e_P$ by the geodesic action of A_P^G is a principal A_P^G-bundle. Let $D(P) = D \times_{A_P^G} \bar{A}_P^G$ be the associated $\bar{A}_P^G$-bundle. The orbits of $A_P^G \times {}^0P(\mathbb{R})$ yield a stratification

$$D(P) = \coprod_{Q\supseteq P} e_Q \ .$$

If $P \subseteq Q$ there is a natural identification $D(Q) \subseteq D(P)$ as an open union of strata.

Given the above identifications, set $\overline{D} = \bigcup_P D(P)$. This is a manifold-with-corners whose stratification by faces is

$$\overline{D} = \coprod_P e_P \ .$$

---

[4] In [22] the group $N_P(\mathbb{R})$ is denoted $\mathscr{U}_P$ and the nilmanifold $N_P(\mathbb{R})'$ is denoted N_P. What we denote here as $\mathcal{N}_P(\mathbb{R})$ and $\mathcal{N}_P(\mathbb{R})'$ are not explicitly mentioned.

The construction of $\overline{D}$ may be generalized to apply to e_P (being a space of type $S - \mathbb{Q}$ [11, 2.3, 3.9]) and the resulting $\overline{e_P}$ may be identified with the closure $\mathrm{cl}(e_P)$ of e_P in $\overline{D}$.

The action of $G(\mathbb{Q})$ extends to $\overline{D}$ and the quotient $\overline{X} = \Gamma\backslash\overline{D}$ is compact. This is the *Borel-Serre compactification* of X. On each stratum e_P the equivalence relation induced by Γ is that of $\Gamma_P = \Gamma \cap P$, and two strata e_P and $e_{P'}$ become identified in the quotient if and only if $P' = {}^\gamma P$ for some $\gamma \in \Gamma$. Thus the strata of $\overline{X}$ are indexed by the Γ-conjugacy classes $\mathcal{P}$ of parabolic $\mathbb{Q}$-subgroups and we have

$$\overline{X} = \coprod_{P \in \mathcal{P}} Y_P \ ,$$

where recall $Y_P = \Gamma_P\backslash e_P$. The bijection $P \mapsto Y_P$ from $\mathcal{P}$ to the strata of $\overline{X}$ is an isomorphism of partially ordered sets, where $Y_P \leq Y_Q$ if and only if $Y_P \subseteq \mathrm{cl}(Y_Q)$. Note that $\mathrm{cl}(Y_Q)$ may be identified with the Borel-Serre compactification $\overline{Y}_Q$ of Y_Q.

2.2. **Local structure of $\overline{X}$.** Set $\bar{\mathcal{A}}_P^G = \mathcal{A}_P^G \times_{\mathcal{A}_P^G} \bar{A}_P^G$. The factorization (1.3.1) extends to $D(P) = \bar{\mathcal{A}}_P^G \times e_P$. It follows from reduction theory that near a point of the stratum e_P of $\overline{D}$, the equivalence relation induced by Γ is that of Γ_P. Thus $\overline{X}$ is locally homeomorphic along Y_P to $\bar{\mathcal{A}}_P^G \times Y_P$ (where we embed $Y_P \subseteq \bar{\mathcal{A}}_P^G \times Y_P$ as $\{o_P\} \times Y_P$). Consequently if Γ is torsion-free then $\overline{X}$ is a smooth manifold-with-corners. In general, since any arithmetic group has a torsion-free subgroup of finite index there may be finite quotient singularities and $\overline{X}$ is a "V-manifold-with-corners".

The description of a neighborhood of the entire closed stratum $\overline{Y}_P$ is much more subtle; see [43, §8].

2.3. **Reductive Borel-Serre compactification.** Let $\widehat{X}$ be the stratified *set*

$$\widehat{X} = \coprod_{P \in \mathcal{P}} X_P$$

and define a stratified map $p \colon \overline{X} \to \widehat{X}$ by setting $p|_{Y_P}$ to be the flat nilmanifold fibration $Y_P \to X_P$ with fiber $\mathcal{N}_P(\mathbb{R})'$. We give $\widehat{X}$ the quotient topology induced by p; this is the *reductive Borel-Serre compactification* of X. (Note that $X = X_G = Y_G$.) Again $P \mapsto X_P$ is an isomorphism of $\mathcal{P}$ with the partially ordered set of strata of $\widehat{X}$. The closure $\mathrm{cl}(X_Q)$ is identified with the reductive Borel-Serre compactification $\widehat{X}_Q$ of X_Q.

We denote the inclusion map of a stratum X_P by $i_P \colon X_P \hookrightarrow \widehat{X}$ and the inclusion of its closure by $\hat{\imath}_P \colon \widehat{X}_P \hookrightarrow \widehat{X}$. Let $U_P = \coprod_{R \geq P} X_P$ be the open star neighborhood of a stratum X_P; we denote the inclusion of the deleted star neighborhood by $j_P \colon U_P \setminus X_P \hookrightarrow \widehat{X}$ and the inclusion of the complement of $\widehat{X}_P$ by $\hat{\jmath}_P \colon \widehat{X} \setminus \widehat{X}_P \hookrightarrow \widehat{X}$.

2.4. **Local structure of $\widehat{X}$.** Note that the identification $A_P^G \cong \mathcal{A}_P^G$ from §1.3 (which depends on the choice of a basepoint) extends to an identification $\bar{\mathcal{A}}_P^G \cong \bar{A}_P^G$. The stratification (2.1.1) of $\bar{A}_P^G$ thus induces one of $\bar{\mathcal{A}}_P^G$ and this stratification is independent of the basepoint.

Set $\mathcal{Z}_P = (\bar{\mathcal{A}}_P^G \times \mathcal{N}_P(\mathbb{R})')/{\sim}$, where $(a, \Gamma_{N_P} n_1) \sim (a, \Gamma_{N_P} n_2)$ if and only if $n_2 = u n_1$ for $u \in N_Q(\mathbb{R})$ and a belongs to the Q-stratum. The action of Γ_{L_P} on $\mathcal{N}_P(\mathbb{R})'$ induces an action on $\mathcal{Z}_P$. Then $\widehat{X}$ is locally homeomorphic along X_P to the bundle $D_P \times_{\Gamma_{L_P}} \mathcal{Z}_P$.

Note that if Γ is neat then the strata X_P are smooth, however in general they are only V-manifolds. Also a description of a neighborhood of the entire closed stratum $\widehat{X}_P$ may easily be deduced from [43, §8].

3. $\mathcal{L}$-MODULES

3.1. **Notation.** Let $W \subseteq \widehat{X}$ be a union of strata, that is, $W = \bigcup_{P \in \mathcal{P}(W)} X_P$ for some subset $\mathcal{P}(W) \subseteq \mathcal{P}$. We say W is an *admissible* space if it is locally closed, or equivalently if P, $Q \in \mathcal{P}(W)$ imply that $[P,Q] \subseteq \mathcal{P}(W)$. We will reuse the notations i_P, $\hat{\imath}_P$, j_P, and $\hat{\jmath}_P$ from §2.3 in order to denote the analogous inclusions into W.

3.2. Let W be an admissible space. Let $\mathcal{L}_W$ denote the system consisting of

$$\begin{cases} \text{the reductive algebraic groups } L_P & \text{for all } P \in \mathcal{P}(W), \text{ and} \\ \text{the functors } H(\mathfrak{n}_P^Q;\cdot)\colon \mathfrak{Mod}(L_Q) \longrightarrow \mathbf{Gr}(L_P) & \text{for all } P \leq Q \in \mathcal{P}(W). \end{cases}$$

Note that the functors $H(\mathfrak{n}_P^Q;\cdot)$ are degree nondecreasing and there are natural isomorphisms (in view of (0.10.4))

$$H(\mathfrak{n}_P^R;\cdot) \cong H(\mathfrak{n}_P^Q; H(\mathfrak{n}_Q^R;\cdot)) \qquad \text{for all } P \leq Q \leq R. \tag{3.2.1}$$

3.3. **The category of $\mathcal{L}$-modules.** Let $\mathcal{L} = \mathcal{L}_W$ for a fixed admissible space W. An *$\mathcal{L}$-module* $\mathcal{M} = (E_\cdot, f_{\cdot\cdot})$ consists of

$$\begin{cases} \text{an object } E_P \text{ of } \mathbf{Gr}(L_P) & \text{for all } P \in \mathcal{P}(W), \text{ and} \\ \text{degree 1 morphisms } f_{PQ}\colon H(\mathfrak{n}_P^Q; E_Q) \xrightarrow{[1]} E_P & \text{for all } P \leq Q \in \mathcal{P}(W) \end{cases}$$

satisfying the condition that

$$\sum_{Q \in [P,R]} f_{PQ} \circ H(\mathfrak{n}_P^Q; f_{QR}) = 0 \qquad \text{for all } P \leq R \in \mathcal{P}(W). \tag{3.3.1}$$

(Equation (3.2.1) shows this formula makes sense.)

Note that (3.3.1) implies in particular that (E_P, f_{PP}) is a complex which we denote $i_P^!\mathcal{M}$. (See §3.4 below for a general definition of $k^!$.)

An *$\mathcal{L}$-morphism* $\mathcal{M} \to \mathcal{M}'$ is a family $\phi = (\phi_{\cdot\cdot})$, where

$$\phi_{PQ}\colon H(\mathfrak{n}_P^Q; E_Q) \to E'_P \qquad \text{for all } P \leq Q \in \mathcal{P}(W),$$

such that the following condition is satisfied:

$$\sum_{Q \in [P,R]} \phi_{PQ} \circ H(\mathfrak{n}_P^Q; f_{QR}) = \sum_{Q \in [P,R]} f'_{PQ} \circ H(\mathfrak{n}_P^Q; \phi_{QR}) \qquad \text{for all } P \leq R \in \mathcal{P}(W). \tag{3.3.2}$$

The composition of two $\mathcal{L}$-morphisms $\phi\colon \mathcal{M} \to \mathcal{M}'$ and $\phi'\colon \mathcal{M}' \to \mathcal{M}''$ is defined by

$$(\phi' \circ \phi)_{PR} = \sum_{Q \in [P,R]} \phi'_{PQ} \circ H(\mathfrak{n}_P^Q; \phi_{QR}) \qquad \text{for all } P \leq R \in \mathcal{P}(W). \tag{3.3.3}$$

The *category of $\mathcal{L}$-modules* $\mathfrak{Mod}(\mathcal{L})$ has for objects the $\mathcal{L}$-modules and for morphisms the $\mathcal{L}$-morphisms. The full subcategory $\mathfrak{Mod}^b(\mathcal{L})$ consists of $\mathcal{L}$-modules $\mathcal{M}$ for which all E_P lie in $\mathbf{Gr}^b(L_P)$. Note that if $W = X_P$ consists of a single stratum, then $\mathfrak{Mod}(\mathcal{L}_{X_P}) = \mathbf{C}(L_P)$, however in general $\mathfrak{Mod}(\mathcal{L})$ is not the category of complexes $\mathbf{C}(\mathcal{C})$ for some category $\mathcal{C}$.

3.4. **Standard functors on $\mathcal{L}$-modules.** Given an inclusion $k\colon Z \hookrightarrow W$ of admissible spaces one can define functors k^*, k_*, $k^!$, and $k_!$ between the respective categories of $\mathcal{L}$-modules that are motivated by the usual functors on the derived category of sheaves.

The functor $k^!$ associates to an $\mathcal{L}_W$-module $\mathcal{M}$ an $\mathcal{L}_Z$-module with the same data but with P and Q restricted to belong to $\mathcal{P}(Z)$. Condition (3.3.1) continues to hold since Z is locally closed. One special case is $i^!_P\mathcal{M} = (E_P, f_{PP})$, the *local cohomology complex at P with supports.* The functor k_* associates to an $\mathcal{L}_Z$-module $\mathcal{M}$ an $\mathcal{L}_W$-module with the same data but extended by $E_P = 0$ and $f_{PQ} = 0$ if one of the subscripts is outside of $\mathcal{P}(Z)$.

We only define k^* in the cases we need, namely when Z is open in W (in which case $k^* = k^!$ has been defined above) and when Z has a unique maximal face. Suppose X_T is a maximal face of Z; that is, $T \in \mathcal{P}(Z)$ and $Z \subseteq \widehat{X}_T$. For $\mathcal{M} = (E_\cdot, f_{\cdot\cdot})$ an $\mathcal{L}_W$-module, define $k^*\mathcal{M} = (E'_\cdot, f'_{\cdot\cdot})$ by

$$\begin{cases} E'_P = \displaystyle\bigoplus_{R\in[P,(P,T)]} H(\mathfrak{n}_P^R; E_R), \\ f'_{PQ} = \displaystyle\sum_{\substack{R\cap T=P\\ S\cap T=Q\\ R\le S}} H(\mathfrak{n}_P^R; f_{RS}), \end{cases}$$

for all $P \le Q$ with P, $Q \in \mathcal{P}(Z)$. (Recall that (P,T) denotes the parabolic $\mathbb{Q}$-subgroup opposite to T relative to P.) The reader can verify this is an $\mathcal{L}_Z$-module. As a special case, the *local cohomology complex at P* is

$$i^*_P\mathcal{M} = \Bigl(\bigoplus_{P\le R} H(\mathfrak{n}_P^R; E_R), \sum_{R\in[P,S]} H(\mathfrak{n}_P^R; f_{RS})\Bigr).$$

It should always be understood in these and similar formulas that R and S are restricted to belong to $\mathcal{P}(W)$.

We only define $k_!$ in the case we need, namely when Z is closed in W, in which case $k_! = k_*$ has been defined above.

Proposition. *Assume in the following that j and k are inclusions of admissible spaces and that the indicated functors have been defined above.*

(i) *There are identities*:

$$\begin{aligned} j^! \circ k^! &= (k\circ j)^! , & k_* \circ j_* &= (k\circ j)_* , \\ j^* \circ k^* &= (k\circ j)^* , & k_! \circ j_! &= (k\circ j)_! . \end{aligned}$$

(ii) *There are adjoint relations*:

$$\begin{aligned} \operatorname{Hom}_{\mathfrak{Mod}(\mathcal{L}_W)}(\mathcal{M}, k_*\mathcal{M}') &\cong \operatorname{Hom}_{\mathfrak{Mod}(\mathcal{L}_Z)}(k^*\mathcal{M}, \mathcal{M}'), \\ \operatorname{Hom}_{\mathfrak{Mod}(\mathcal{L}_W)}(\mathcal{M}, k^!\mathcal{M}') &\cong \operatorname{Hom}_{\mathfrak{Mod}(\mathcal{L}_Z)}(k_!\mathcal{M}, \mathcal{M}'). \end{aligned} \tag{3.4.1}$$

(iii) *Consider the commutative diagram of inclusions of admissible spaces,*

$$\begin{tikzcd} Z \arrow[r, "k_2"] \arrow[d, "j_1"'] & Z_2 \arrow[d, "j_2"'] \\ Z_1 \arrow[r, "k_1"] & W . \end{tikzcd}$$

where $Z = Z_1 \cap Z_2$. Then

$$k_1^! \circ j_{2*} = j_{1*} \circ k_2^! \ . \tag{3.4.2}$$

Now assume that Z_1 has a unique maximal face X_T, that Z has a unique maximal face X_{T_0}, and furthermore that Z_1 and Z_2 intersect "transversely" in the sense that

$$U \cap Z_2 = U \cap W \cap \widehat{X}_{(T_0,T)}, \tag{3.4.3}$$

where $U = \bigcup_{P\in\mathcal{P}(Z)} \bigcup_{Q\geq P} X_Q$ is the star neighborhood of Z. Then

$$j_1^! \circ k_1^* = k_2^* \circ j_2^!.$$

Proof. Parts (i) and (ii) and the first part of (iii) are left to the reader. For the rest of (iii), note that $E'_P = \bigoplus_Q H(\mathfrak{n}_P^Q; E_Q)$ for both $j_1^! k_1^* \mathcal{M}$ and $k_2^* j_2^! \mathcal{M}$, where in the first case the sum is over $Q \in [P,(P,T)] \cap \mathcal{P}(W)$, while in the second case the sum is over $Q \in [P,(P,T_0)] \cap \mathcal{P}(Z_2)$. The sums are equal by (3.4.3) since $(P,T) = (P,T_0) \cap (T_0,T)$. ☐

Elsewhere we will define k^* and $k_!$ more generally and prove an analogue of the proposition in a suitable homotopy category of $\mathcal{L}$-modules (see §3.9).

3.5. **Pullback of an $\mathcal{L}_W$-module to a fiber.** There is one other functor we will need to consider beginning in §22. Suppose G has a connected normal $\mathbb{Q}$-subgroup which we denote G_ℓ. Let $\tilde{G}_h$ be a complementary connected normal $\mathbb{Q}$-subgroup so that $G = \tilde{G}_h G_\ell$ is an almost direct product. Set $G_h = G/G_\ell = \tilde{G}_h/(G_\ell \cap \tilde{G}_h)$ and let $\Gamma_{G_\ell} = \Gamma \cap G_\ell$ and $\Gamma_{G_h} = \Gamma/\Gamma_{G_\ell}$ be the induced arithmetic subgroups. Parabolic $\mathbb{Q}$-subgroups of G correspond to pairs of a parabolic $\mathbb{Q}$-subgroup of G_ℓ and a parabolic $\mathbb{Q}$-subgroup of G_h: given $P \subseteq G$ we associate $P_\ell = P \cap G_\ell$ and $P_h = P/P_\ell$ and conversely given P_ℓ and P_h we associate $P = \tilde{P}_h P_\ell$ (where $\tilde{P}_h \subseteq \tilde{G}_h$ is a lift of P_h).

The geometric picture is as follows. There is a factorization of symmetric spaces $D = D_h \times D_\ell$ and a flat bundle of arithmetic quotients $X \to X_h$ with fibers X_ℓ. (The action of Γ_{G_h} is induced from the action of the finite quotient $\Gamma/\Gamma_\ell(\Gamma \cap \tilde{G}_h)$, so this bundle becomes trivial over a finite cover of X_h.) Let $X(G_\ell) \subseteq \widehat{X}$ be the partial compactification of X obtained by replacing the fibers by $\widehat{X}_\ell$. Specifically, note that the map $P \mapsto P_h$ induces a surjection $\mathcal{P}(\widehat{X}) \to \mathcal{P}(\widehat{X}_h)$. Then set $X(G_\ell)$ to be the union of strata X_P for all $P \in \mathcal{P}(\widehat{X})$ satisfying $P_h = G_h$. The above fibration extends to a fibration

$$\pi \colon X(G_\ell) \longrightarrow X_h$$

with fibers $\widehat{X}_\ell$. Let $\hat{\imath}_{G,\ell} \colon \widehat{X}_\ell \hookrightarrow X(G_\ell)$ be the inclusion of a generic fiber.

The map $P_\ell \mapsto P = \tilde{G}_h P_\ell$ induces a surjection $\mathcal{P}(\widehat{X}_\ell) \to \mathcal{P}(X(G_\ell))$. (It may not be injective if the finite group $\Gamma/\Gamma_\ell(\Gamma \cap \tilde{G}_h)$ is not trivial.) Define the connected normal $\mathbb{Q}$-subgroup $L_{P,\ell} = P_\ell/N_P \subseteq L_P$. Let $\mathcal{M} = (E_\cdot, f_{\cdot\cdot})$ be an $\mathcal{L}_{X(G_\ell)}$-module. Define an $\mathcal{L}_{\widehat{X}_\ell}$-module $\hat{\imath}_{G,\ell}^* \mathcal{M} = (E'_\cdot, f'_{\cdot\cdot})$ by

$$\begin{cases} E'_{P_\ell} = \operatorname{Res}^{L_P}_{L_{P,\ell}} E_P, \\ f'_{P_\ell Q_\ell} = \operatorname{Res}^{L_P}_{L_{P,\ell}} f_{PQ}, \end{cases}$$

for all $P_\ell \le Q_\ell \in \mathcal{P}(\widehat{X}_\ell)$. (Here $\operatorname{Res}^{L_P}_{L_{P,\ell}}$ denotes the restriction of a representation of L_P to a representation of $L_{P,\ell}$.) This makes sense since there is a natural isomorphism

$$\operatorname{Res}^{L_P}_{L_{P,\ell}} H(\mathfrak{n}_P^Q; E_Q) \cong H(\mathfrak{n}_{P_\ell}^{Q_\ell}; \operatorname{Res}^{L_Q}_{L_{Q,\ell}} E_Q) \ .$$

Similarly define $\hat{\imath}^!_{G,\ell}\mathcal{M}$ except that in this case $E'_{P_\ell} = \operatorname{Res}^{L_P}_{L_{P,\ell}} E_P[-\dim D_h]$.

More generally for $R \in \mathcal{P}$ we may define $\hat{\imath}^*_{R,\ell}$ and $\hat{\imath}^!_{R,\ell}$ when we are given a connected normal $\mathbb{Q}$-subgroup $L_{R,\ell}$ of L_R. In this case we have a fibration $X_R(L_{R,\ell}) \to X_{R,h}$ with fiber $\widehat{X}_{R,\ell}$ and a surjection $\mathcal{P}(\widehat{X}_{R,\ell}) \to \mathcal{P}(X_R(L_{R,\ell}))$ given by $P_\ell \mapsto P$ where $P/N_R = \widetilde{L_{R,h}}P_\ell$. We note the following identities for $P_\ell \in \mathcal{P}(\widehat{X}_{R,\ell})$:

$$\begin{aligned} i^*_{P_\ell} \circ \hat{\imath}^*_{R,\ell} &= i^*_{P_\ell} \circ \hat{\imath}^*_{P,\ell} \circ i^*_P \ , & i^*_{P_\ell} \circ \hat{\imath}^!_{R,\ell} &= i^*_{P_\ell} \circ \hat{\imath}^!_{P,\ell} \circ i^*_P \ , \\ i^!_{P_\ell} \circ \hat{\imath}^*_{R,\ell} &= i^*_{P_\ell} \circ \hat{\imath}^*_{P,\ell} \circ i^!_P \ , & i^!_{P_\ell} \circ \hat{\imath}^!_{R,\ell} &= i^*_{P_\ell} \circ \hat{\imath}^!_{P,\ell} \circ i^!_P \ . \end{aligned} \tag{3.5.1}$$

(The initial $i^*_{P_\ell}$ on the right-hand side of these equalities would not be needed if $\Gamma/\Gamma_\ell(\Gamma \cap \tilde{G}_h)$ were trivial.)

3.6. **Local cohomology with supports.** Let $\mathcal{M}$ be an $\mathcal{L}_W$-module, where W is an admissible space. It is easy to verify that there is a short exact sequence

$$0 \to i^!_Q\mathcal{M} \to i^*_Q\mathcal{M} \to i^*_Q j_{Q*} j^*_Q \mathcal{M} \to 0 \tag{3.6.1}$$

relating the two types of local cohomology complexes at Q. There is also a short exact sequence

$$0 \to \hat{\imath}^!_Q\mathcal{M} \to \hat{\imath}^*_Q\mathcal{M} \to \hat{\imath}^*_Q \hat{\jmath}_{Q*} \hat{\jmath}^*_Q \mathcal{M} \to 0 \ . \tag{3.6.2}$$

More generally, for $P \le Q \in \mathcal{P}(W)$ define the *local cohomology complex of* $\mathcal{M}$ *at* P *supported on* Q to be

$$i^*_P \hat{\imath}^!_Q \mathcal{M} = \Bigl(\bigoplus_{R \in [P,Q]} H(\mathfrak{n}_P^R; E_R), \sum_{R \le S \in [P,Q]} H(\mathfrak{n}_P^R; f_{RS}) \Bigr) \ ;$$

this will play an essential role in the definition of micro-support of $\mathcal{M}$ later in §7. If $Q \le Q'$ there is a short exact sequence comparing $i^*_P \hat{\imath}^!_Q \mathcal{M}$ and $i^*_P \hat{\imath}^!_{Q'} \mathcal{M}$, namely

$$0 \to i^*_P \hat{\imath}^!_Q \mathcal{M} \to i^*_P \hat{\imath}^!_{Q'} \mathcal{M} \to i^*_P \hat{\jmath}_{Q*} \hat{\jmath}^*_Q \hat{\imath}^!_{Q'} \mathcal{M} \to 0 \ ; \tag{3.6.3}$$

this follows from (3.6.2) by replacing $\mathcal{M}$ with $\hat{\imath}^!_{Q'}\mathcal{M}$ and applying i^*_P. The corresponding long exact sequence is

$$\cdots \longrightarrow H^i(i^*_P \hat{\imath}^!_Q \mathcal{M}) \longrightarrow H^i(i^*_P \hat{\imath}^!_{Q'} \mathcal{M}) \longrightarrow H^i(i^*_P \hat{\jmath}_{Q*} \hat{\jmath}^*_Q \hat{\imath}^!_{Q'} \mathcal{M}) \longrightarrow \cdots \ . \tag{3.6.4}$$

We wish to study $H(i^*_P \hat{\jmath}_{Q*} \hat{\jmath}^*_Q \hat{\imath}^!_{Q'} \mathcal{M})$. Set $P' = (P,Q) \cap Q'$ so that $P = P' \cap Q$ and $Q' = P' \vee Q$:

$$\begin{tikzcd} & Q' & \\ P' \arrow[ur, dash] & & Q \arrow[ul, dash] \\ & P \arrow[ul, dash] \arrow[ur, dash] & \end{tikzcd} \tag{3.6.5}$$

When $\#\Delta_P^{P'} = 1$ the last term of (3.6.3) is equal to

$$i^*_P i_{P'*} i^*_{P'} \hat{\imath}^!_{Q'} \mathcal{M} = H(\mathfrak{n}_P^{P'}; i^*_{P'} \hat{\imath}^!_{Q'} \mathcal{M}) \ ;$$

since the functor $H(\mathfrak{n}_P^{\tilde P};\cdot)$ is exact it commutes with taking cohomology and so

$$H(i_P^* i_{P'*} i_{P'}^* \hat\imath_{Q'}^! \mathcal{M}) = H(\mathfrak{n}_P^{P'}; H(i_{P'}^* \hat\imath_{Q'}^! \mathcal{M}))\ . \tag{3.6.6}$$

In general we have the

3.7. Lemma. *Given* $P \le Q \le Q'$, *set* $P' = (P,Q)\cap Q'$. *There are two spectral sequences abutting to* $H(i_P^* \hat\jmath_{Q*} \hat\jmath_Q^* \hat\imath_{Q'}^! \mathcal{M})$. *The first* (*the Fary spectral sequence*) *has*

$$E_1^{-p,\cdot} = \bigoplus_{\substack{P<\tilde P\le P'\\ \#\Delta_P^{\tilde P}=p}} H(\mathfrak{n}_P^{\tilde P}; H(i_{\tilde P}^* \hat\imath_{\tilde Q}^! \mathcal{M}))[-p]\ , \tag{3.7.1}$$

where $\tilde Q = \tilde P \vee Q$. *The second* (*the Mayer-Vietoris spectral sequence*) *has*

$$E_1^{p,\cdot} = \bigoplus_{\substack{P<\tilde P\le P'\\ \#\Delta_P^{\tilde P}=p+1}} H(\mathfrak{n}_P^{\tilde P}; H(i_{\tilde P}^* \hat\imath_{Q'}^! \mathcal{M}))\ . \tag{3.7.2}$$

Proof. Any $R \in [P,Q']$ belongs to $[\tilde P, \tilde Q]$ for a unique $\tilde P \in [P,P']$ (namely $\tilde P = R \cap P'$). The situation is represented by the following diagram, in which each parallelogram has the analogous meaning to (0.10.2):

(3.7.3)



.

(Of course this figure is not meant to suggest that the parabolic $\mathbb{Q}$-subgroups lying between P and P' are linearly ordered.)

For the first spectral sequence, decompose $i_P^* \hat\jmath_{Q*} \hat\jmath_Q^* \hat\imath_{Q'}^! \mathcal{M}$ (as an L_P-*module*, not as a complex) as

$$\bigoplus_{P<\tilde P\le P'} i_P^* i_{\tilde P*} i_{\tilde P}^* \hat\imath_{\tilde Q}^! \mathcal{M} = \bigoplus_{P<\tilde P\le P'} H(\mathfrak{n}_P^{\tilde P}; i_{\tilde P}^* \hat\imath_{\tilde Q}^! \mathcal{M})\ . \tag{3.7.4}$$

The corresponding sum in which $\tilde P$ is restricted to those with $\#\Delta_P^{\tilde P} \le p$ is a subcomplex and defines an increasing filtration on $i_P^* \hat\jmath_{Q*} \hat\jmath_Q^* \hat\imath_{Q'}^! \mathcal{M}$ as p varies. The associated graded complex is exactly equal to (3.7.4). In view of (3.6.6), the E_1 term of the resulting spectral sequence has the desired form, (3.7.1).

For the Mayer-Vietoris spectral sequence, consider the long exact sequence

$$0 \to i_P^* \hat\jmath_{Q*} \hat\jmath_Q^* \hat\imath_{Q'}^! \mathcal{M} \to \bigoplus_{\#\Delta_P^{\tilde P}=1} i_P^* i_{\tilde P*} i_{\tilde P}^* \hat\imath_{Q'}^! \mathcal{M} \to \bigoplus_{\#\Delta_P^{\tilde P}=2} i_P^* i_{\tilde P*} i_{\tilde P}^* \hat\imath_{Q'}^! \mathcal{M} \to \cdots$$

in which $\tilde P$ satisfies $P < \tilde P \le P'$. Thus $H(i_P^* \hat\jmath_{Q*} \hat\jmath_Q^* \hat\imath_{Q'}^! \mathcal{M})$ is the cohomology of a double complex; the spectral sequence (3.7.2) corresponds to the "first filtration" $\#\Delta_P^{\tilde P} \ge p+1$. □

3.8. The *mapping cone* of an $\mathscr{L}$-morphism $\phi\colon \mathcal{M}\to\mathcal{M}'$ is defined by

$$M(\phi)=\left(E_\cdot[1]\oplus E'_\cdot\,,\begin{pmatrix}-f_{\cdot\cdot} & 0\\ \phi_{\cdot\cdot} & f'_{\cdot\cdot}\end{pmatrix}\right).$$

This is an $\mathscr{L}$-module; condition (3.3.1) follows from the corresponding condition for $\mathcal{M}$ and $\mathcal{M}'$ and from (3.3.2). The mapping cone defines a functor from the category of diagrams $\phi\colon \mathcal{M}\to\mathcal{M}'$ to $\mathfrak{Mod}(\mathscr{L})$. Define the *shift* of an $\mathscr{L}$-module by $\mathcal{M}[1]=(E_\cdot[1],-f_{\cdot\cdot})$. There are natural $\mathscr{L}$-morphisms $\alpha(\phi)\colon\mathcal{M}'\to M(\phi)$ and $\beta(\phi)\colon M(\phi)\to\mathcal{M}[1]$ given by

$$\alpha(\phi)_{PP}=\begin{pmatrix}0\\ \mathrm{id}_{E'_P}\end{pmatrix}\qquad\text{for all } P,$$
$$\beta(\phi)_{PP}=\begin{pmatrix}\mathrm{id}_{E_P[1]} & 0\end{pmatrix}\qquad\text{for all } P.$$

(Set $\alpha(\phi)_{PQ}=0$ for $P\lneqq Q$ and similarly for $\beta(\phi)$.)

3.9. One may define a notion of homotopy for $\mathscr{L}$-morphisms and set $\mathbf{K}(\mathscr{L})$ to be the category with $\mathscr{L}$-modules as objects and homotopy classes of $\mathscr{L}$-morphisms as morphisms. With the definition of mapping cone above, $\mathbf{K}(\mathscr{L})$ may be given the structure of a *triangulated category* [27, §§1.4, 1.5]. A *quasi-isomorphism* of $\mathscr{L}$-modules is an $\mathscr{L}$-morphism that induces quasi-isomorphisms on local cohomology complexes. Since $\mathfrak{Mod}(L_P)$ is a semisimple abelian category for all P one can show that any quasi-isomorphism of $\mathscr{L}$-modules has a homotopy inverse. Thus it is reasonable to notate $\mathbf{K}(\mathscr{L})$ also as $\mathbf{D}(\mathscr{L})$. We will discuss this in more detail elsewhere.

3.10. **Generalizations.** The concept of $\mathscr{L}$-modules may be greatly generalized. For example, in §3.2 one could start with a general family of reductive $\mathbb{Q}$-groups $(L_\sigma)_\sigma$ indexed by a partially ordered set and replace $H(\mathfrak{n}_P^Q;\cdot)$ by some "link cohomology" functors $\mathcal{M}_{\sigma\tau}\colon \mathfrak{Mod}(L_\tau)\longrightarrow\mathbf{Gr}(L_\sigma)$ satisfying $\mathcal{M}_\sigma^\upsilon\cong\mathcal{M}_\sigma^\tau\circ\mathcal{M}_\tau^\upsilon$. One particularly simple situation will be introduced in §12.10 in which all the groups L_σ will be identical. One could also drop the condition that the groups are reductive algebraic and replace $\mathfrak{Mod}(L_\sigma)$ by a suitable category of representations. We will not consider such generalizations further here.

4. REALIZATION OF $\mathscr{L}$-MODULES

For an admissible space W let $\mathcal{X}$ denote the stratification $W=\coprod_{P\in\mathscr{P}(W)}X_P$. Recall that $\mathbf{C}_{\mathcal{X}}(W)$ denotes the category of constructible complexes of sheaves on W and that $\mathbf{D}_{\mathcal{X}}(W)$ denotes the corresponding derived category. Note that a regular L_P-module E_P induces by restriction a Γ_{L_P}-module and hence a locally constant sheaf $\mathbb{E}_P=E_P\times_{\Gamma_P}D_P$ on X_P. Thus we obtain a functor $\mathfrak{Mod}(L_P)\to\mathbf{D}_{\mathcal{X}}(X_P)$. In this section we generalize this to $\mathscr{L}$-modules in order to obtain functors $\mathcal{S}_W\colon\mathfrak{Mod}(\mathscr{L}_W)\to\mathbf{D}_{\mathcal{X}}(W)$.

4.1. **Theorem.**

(i) *There exists a family of functors*

$$\mathcal{A}_P\colon\mathfrak{Mod}(L_P)\to\mathbf{C}_{\mathcal{X}}(\widehat{X}_P)\qquad\textit{for all } P\in\mathscr{P}$$

and natural morphisms for all $P\le Q\in\mathscr{P}$,

$$k_{PQ}\colon \hat{\imath}_{Q*}\mathcal{A}_Q(E_Q)\longrightarrow\hat{\imath}_{P*}\mathcal{A}_P(H(\mathfrak{n}_P^Q;E_Q))\qquad\textit{for all } E_Q\in\mathfrak{Mod}(L_Q).\tag{4.1.1}$$

Each $\mathcal{A}_P$ *is an incarnation of the functor* $E_P \mapsto i_{P*}\mathbb{E}_P$, $R(L_P) \to \mathbf{D}_{\mathcal{X}}(\widehat{X}_P)$, *and* $\hat{\imath}_P^*(k_{PQ})$ *becomes a natural isomorphism in* $\mathbf{D}_{\mathcal{X}}(\widehat{X}_P)$.

(ii) *A family of functors as in* (i) *determines functors*

$$\mathcal{A}_W\colon \mathfrak{Mod}(\mathfrak{L}_W) \to \mathbf{C}_{\mathcal{X}}(W) \qquad \textit{for all admissible spaces } W, \textit{ and}$$

corresponding functors $\mathcal{S}_W$ *into* $\mathbf{D}_{\mathcal{X}}(W)$. *If* $k\colon Z \hookrightarrow W$ *is an inclusion of admissible spaces, then* $k_* \circ \mathcal{S}_Z = \mathcal{S}_W \circ k_*$ *and similarly for the other functors defined in* §§3.4 *and* 3.5.

(iii) *If two families of functors as in* (i) *are naturally quasi-isomorphic compatibly with* (4.1.1), *then the corresponding functors* $\mathcal{S}_W$ *are naturally isomorphic.*

Remark. A functor $\mathcal{S}_W$ as in the theorem with be called a *realization* of the category of $\mathfrak{L}_W$-modules. Given a realization functor and an $\mathfrak{L}_W$-module $\mathcal{M}$, the *cohomology* $H(W;\mathcal{M})$ *of* $\mathcal{M}$ is defined to be the hypercohomology $H(W;\mathcal{S}_W(\mathcal{M}))$. More generally the cohomology $H(Y;\mathcal{M})$ of any subset $Y \subseteq W$ with coefficients in $\mathcal{M}$ is defined to be the hypercohomology of $\mathcal{S}_W(\mathcal{M})|_Y$. In this paper we will use the incarnation $\mathcal{A}_W(\mathcal{M})$ constructed below using special differential forms. This is a complex of fine sheaves so $H(\widehat{X};\mathcal{M})$ may be computed from the complex of global sections of $\mathcal{A}_{\widehat{X}}(\mathcal{M})$.

4.2. **Proof of Theorem 4.1(ii)(iii).** We assume that a family of functors $\{\mathcal{A}_P\}_{P\in\mathcal{P}}$ exists as in (i). Let $\mathcal{M} = (E_\cdot, f_{\cdot\cdot})$ be an $\mathfrak{L}_W$-module. Define

$$\begin{cases} \mathcal{A}_W(\mathcal{M}) = \displaystyle\bigoplus_{P\in\mathcal{P}(W)} \hat{\imath}_{P*}\mathcal{A}_P(E_P), \\ d_{\mathcal{A}_W(\mathcal{M})} = \displaystyle\sum_{P\in\mathcal{P}(W)} d_P + \sum_{P\le Q\in\mathcal{P}(W)} \mathcal{A}_P(f_{PQ}) \circ k_{PQ}\ , \end{cases} \tag{4.2.1}$$

where d_P is the differential of $\mathcal{A}_P(E_P)$. (Note that in applying $\mathcal{A}_P$ to E_P we are using the sign convention of §0.10.2.) This is a complex of sheaves. For $\phi = (\phi_{\cdot\cdot})$ an $\mathfrak{L}_W$-morphism define

$$\mathcal{A}_W(\phi) = \sum_{P\le Q} \mathcal{A}_P(\phi_{PQ}) \circ k_{PQ}\ .$$

This is the desired functor for (ii); we let $\mathcal{S}_W$ be the corresponding functor into $\mathbf{D}_{\mathcal{X}}(W)$. It is easy to verify that $\mathcal{S}_W$ commutes with the usual functors defined in §§3.4 and 5.3 and that (iii) holds.

4.3. **Special differential forms.** We now prepare for the proof of Theorem 4.1(i) which will be at the end of this section.

Let E be a regular representation of G and let $\mathbb{E}$ denote the corresponding locally constant sheaf on $\overline{X}$, or on any stratum of $\overline{X}$. For any stratum Y_P of $\overline{X}$, consider the diagram

$$\overline{X} = \Gamma\backslash\overline{D} \xleftarrow{q} \Gamma_P\backslash D(P) \xrightarrow{r_P} Y_P,$$

where $q\colon \Gamma_P\backslash D(P) \hookrightarrow \Gamma_P\backslash\overline{D} \to \Gamma\backslash\overline{D}$ is the inclusion followed by the quotient and r_P is the geodesic retraction. Reduction theory implies that if $y \in Y_P$ there exists a sufficiently small neighborhood $U \subseteq \overline{X}$ of y such that $q\colon q^{-1}(U) \xrightarrow{\sim} U$ is a diffeomorphism. An $\mathbb{E}$-valued differential form ω on $\overline{X}$ is *locally lifted from the boundary* [6, §8] if for all strata Y_P and all $y \in Y_P$, there exists U as above so that $q^*(\omega|_U) = r_P^*\psi$ where ψ is a smooth form on $U \cap Y_P$. A form on a stratum $Y_P = \Gamma_P\backslash e_P$ is called $N_P(\mathbb{R})$-*invariant* if its lift to e_P is invariant under the action of $N_P(\mathbb{R})$; let $\mathcal{A}_{\mathrm{inv}}(Y_P;\mathbb{E})$ denote the sheaf of $N_P(\mathbb{R})$-invariant $\mathbb{E}$-valued forms on Y_P and let $A_{\mathrm{inv}}(Y_P;\mathbb{E})$ denote the global sections.

Let $\mathcal{A}_{\mathrm{sp}}(\overline{X};\mathbb{E}) \subseteq \mathcal{A}(\overline{X};\mathbb{E})$ denote the subcomplex of forms which are locally lifted from the boundary and whose restriction to each stratum Y_P is in $\mathcal{A}_{\mathrm{inv}}(Y_P;\mathbb{E})$. The sheaf complex of *special differential forms* on $\widehat{X}$ is [22, (13.2)]

$$\mathcal{A}_{\mathrm{sp}}(\widehat{X};\mathbb{E}) = p_*\mathcal{A}_{\mathrm{sp}}(\overline{X};\mathbb{E}) \ ,$$

the pushforward under the quotient map $p\colon \overline{X} \to \widehat{X}$. Let $A_{\mathrm{sp}}(\overline{X};\mathbb{E})$ and $A_{\mathrm{sp}}(\widehat{X};\mathbb{E})$ denote the corresponding complexes of global sections.

4.4. **Restriction to boundary strata.** Recall from §1.4 that the nilmanifold fibration $p\colon Y_P \to X_P$ is the flat bundle associated to the Γ_{L_P}-space $\mathcal{N}_P(\mathbb{R})'$, and that given a basepoint $x \in D$ there is an isomorphism $n'_x\colon N_P(\mathbb{R})' \xrightarrow{\sim} \mathcal{N}_P(\mathbb{R})'$. The Γ_{L_P}-action on $\mathcal{N}_P(\mathbb{R})'$ induces the structure of an Γ_{L_P}-module on $A_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E})$, the $N_P(\mathbb{R})$-invariant forms; let $\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E})$ denote[5] the associated complex of locally constant sheaves on X_P. The fiber $\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E})_x$ is the complex of $N_P(\mathbb{R})$-invariant forms on $p^{-1}(x)$.

A special differential form ω on $\widehat{X}$ determines an $N_P(\mathbb{R})$-invariant form on Y_P which we denote $\omega|_{Y_P}$; this form can be viewed as a differential form on X_P with values in $\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E})$ which we denote $\omega|_{X_P}$. In fact there is a natural isomorphism of complexes [22, 12.6, 13.4(2)]

$$i_P^*\mathcal{A}_{\mathrm{sp}}(\widehat{X};\mathbb{E}) \xrightarrow{\sim} \mathcal{A}(X_P;\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E})), \tag{4.4.1}$$

where the complex on the right is the associated total complex. This operation extends to an isomorphism

$$\hat{\imath}_P^*\mathcal{A}_{\mathrm{sp}}(\widehat{X};\mathbb{E}) \xrightarrow{\sim} \mathcal{A}_{\mathrm{sp}}(\widehat{X}_P;\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E}))$$

and hence there is a natural morphism, denoted $\omega \mapsto \omega|_{\widehat{X}_P}$,

$$\mathcal{A}_{\mathrm{sp}}(\widehat{X};\mathbb{E}) \longrightarrow \hat{\imath}_{P*}\mathcal{A}_{\mathrm{sp}}(\widehat{X}_P;\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E})) \ . \tag{4.4.2}$$

4.5. A theorem of Nomizu [36] and van Est [55] implies that the natural inclusion of complexes of differential forms induces an isomorphism

$$H(A_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E})) \cong H(\mathcal{N}_P(\mathbb{R})';\mathbb{E});$$

from this, the Poincaré lemma, and (4.4.1) one sees that

Lemma. ([22, §13]) *There is a natural isomorphism* $\mathcal{A}_{\mathrm{sp}}(\widehat{X};\mathbb{E}) \xrightarrow{\sim} i_{G*}\mathbb{E}$ *in* $\mathbf{D}_{\mathcal{X}}(\widehat{X})$ *for* $E \in \mathfrak{Mod}(G)$.

4.6. Recall that $C(\mathfrak{n}_P;E)$ denotes the complex $\operatorname{Hom}(\bigwedge \mathfrak{n}_P;E)$ with differential as in [13, I, §1.1]. Its cohomology $H(\mathfrak{n}_P;E)$ is a graded L_P-module; let $\mathbb{H}(\mathfrak{n}_P;E)$ denote the corresponding graded locally constant sheaf on X_P.

4.7. **Lemma.** (compare [22, §12]) *There is a natural quasi-isomorphism on* X_P:

$$h_P\colon \mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})';\mathbb{E}) \longrightarrow \mathbb{H}(\mathfrak{n}_P;E).$$

---

[5] In [22] this complex is denoted $\mathbf{C}^{\cdot}(N_P;\mathbf{E})$.

Proof. Choose a basepoint $x \in D$ and let $L_{P,x} \subseteq P$ denote the lift of L_P stable under the Cartan involution associated to x. L_P acts on $C(\mathfrak{n}_P; E)$ through the coadjoint action of $L_{P,x}$; we denote by $C(\mathfrak{n}_P; E)_x$ the resulting complex of L_P-modules and by $\mathbb{C}(\mathfrak{n}_P; E)_x$ the corresponding complex of locally constant sheaves.

The quasi-isomorphism h_P is given by a composition of quasi-isomorphisms

$$\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})'; \mathbb{E}) \xrightarrow{h_x} \mathbb{C}(\mathfrak{n}_P; E)_x \longrightarrow \mathbb{H}(\mathfrak{n}_P; E)\ . \tag{4.7.1}$$

The first map is induced by an isomorphism [22, (12.13)] of the underlying Γ_{L_P}-modules. Namely, transfer a form from $\mathcal{N}_P(\mathbb{R})'$ to $N_P(\mathbb{R})'$ via the isomorphism n'_x and evaluate the form at the identity; one may use §1.5 to verify this respects the Γ_{L_P}-action. For the second map of (4.7.1), note that if E were irreducible, a theorem of Kostant [31] says that each irreducible component of $H(\mathfrak{n}_P; E)$ occurs in both $H(\mathfrak{n}_P; E)$ and $C(\mathfrak{n}_P; E)_x$ with multiplicity one. Thus for general regular E there is a unique map $C(\mathfrak{n}_P; E)_x \to H(\mathfrak{n}_P; E)$ in $\mathbf{C}(L_P)$ inducing the identity on cohomology. This induces the second map in (4.7.1).

If x' is an another basepoint then $L_{P,x'} = vL_{P,x}v^{-1}$ for some $v \in N_P(\mathbb{R})$. Thus there is an isomorphism $\mathrm{Ad}^*(v)\colon C(\mathfrak{n}_P; E)_x \xrightarrow{\sim} C(\mathfrak{n}_P; E)_{x'}$ in $\mathbf{C}(L_P)$. One may check from the definition of h_x that the left triangle of

$$\begin{tikzcd}
 & \mathbb{C}(\mathfrak{n}_P; E)_x \arrow[dd, "\mathrm{Ad}^*(v)"'] \arrow[dr] & \\
\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P(\mathbb{R})'; \mathbb{E}) \arrow[ur, "h_x"] \arrow[dr, "h_{x'}"'] & & \mathbb{H}(\mathfrak{n}_P; E) \\
 & \mathbb{C}(\mathfrak{n}_P; E)_{x'} \arrow[ur] &
\end{tikzcd}$$

commutes; since $\mathrm{Ad}^*(v)$ induces the identity on cohomology, the right triangle commutes by the uniqueness noted above. Thus h_P is independent of the basepoint x.

The verification of naturality for each map in (4.7.1) is left to the reader. □

4.8. This lemma together with equation (4.4.2) yields the

Corollary. *There is a natural morphism*

$$k_P \colon \mathcal{A}_{\mathrm{sp}}(\widehat{X}; \mathbb{E}) \longrightarrow \hat{\imath}_{P*}\mathcal{A}_{\mathrm{sp}}(\widehat{X}_P; \mathbb{H}(\mathfrak{n}_P; E))$$

given by $\omega \mapsto \mathcal{A}_{\mathrm{sp}}(\widehat{X}_P; h_P)(\omega|_{\widehat{X}_P})$ *such that* $\hat{\imath}_P^*(k_P)$ *is a quasi-isomorphism.*

4.9. **Proof of Theorem 4.1(i).** Apply Lemma 4.5, Lemma 4.7, and Corollary 4.8 with X and G replaced by X_Q and L_Q respectively. We obtain for all $P \leq Q$ and all $E_Q \in \mathfrak{Mod}(L_Q)$ natural quasi-isomorphisms

$$\mathcal{A}_{\mathrm{sp}}(\widehat{X}_Q; \mathbb{E}_Q) \longrightarrow i_{Q*}\mathbb{E}_Q\ ,$$
$$h_{PQ} \colon \mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P^Q(\mathbb{R})'; \mathbb{E}_Q) \longrightarrow \mathbb{H}(\mathfrak{n}_P^Q; E_Q)\ ,$$

and a natural morphism

$$k_{PQ} \colon \hat{\imath}_{Q*}\mathcal{A}_{\mathrm{sp}}(\widehat{X}_Q; \mathbb{E}_Q) \longrightarrow \hat{\imath}_{P*}\mathcal{A}_{\mathrm{sp}}(\widehat{X}_P; \mathbb{H}(\mathfrak{n}_P^Q; E_Q))\ ,$$

where $k_{PQ}(\omega) = \mathcal{A}_{\mathrm{sp}}(\widehat{X}_P; h_{PQ})(\omega|_{\widehat{X}_P})$ and $\hat{\imath}_P^*(k_{PQ})$ is a quasi-isomorphism. The proposition follows if we set $\mathcal{A}_P(E_P) = \mathcal{A}_{\mathrm{sp}}(\widehat{X}_P; \mathbb{E}_P)$. □

4.10. There are other choices of $\{\mathcal{A}_P\}_{P\in\mathcal{P}}$ that yield different incarnations of the same realization $\mathcal{S}_W$. Consider for simplicity just $P = G$. Then to avoid the boundary conditions of special differential forms at the expense of complicating the combinatorics, one may set $\mathcal{A}_G(E)$ equal to the resolution of $i_{G*}\mathbb{E}$ by *combinatorial forms*

$$\mathcal{A}_{\mathrm{comb}}(\widehat{X};\mathbb{E}) = \bigoplus_{P\leq Q} \mathcal{A}((\Gamma_P/\Gamma_{N_Q})\backslash(\overline{N_Q(\mathbb{R})\backslash e_P}); \mathbb{A}_{\mathrm{inv}}(\mathcal{N}_Q(\mathbb{R})';\mathbb{E})).$$

As another example, if E has a $\mathbb{Q}$-structure one may set $\mathcal{A}_G(E)$ equal to the resolution $i_{G*}\mathbf{I}(\mathbb{E})$ defined in [22, (27.9)]; in this way one obtains a incarnation $\mathcal{A}_W$ defined over $\mathbb{Q}$ for $\mathcal{L}$-modules defined over $\mathbb{Q}$. We will not go into further details here.

5. Example: the Intersection Cohomology $\mathcal{L}$-module

Let $W \subseteq \widehat{X}$ be an admissible subspace.

5.1. Recall that there are truncation functors on $\mathbf{C}(L_P)$ given by

$$\begin{aligned} \tau^{\leqslant n}C\colon &\quad \cdots \to C^{n-1} \to \operatorname{Ker} d^n_C \to 0 \to \cdots\ , \\ \tau^{>n}C\colon &\quad \cdots \to 0 \to C^{n+1}/\operatorname{Im} d^n_C \to C^{n+2} \to \cdots\ . \end{aligned}$$

There is a natural short exact sequence $0 \to \tau^{\leqslant n}C \to C \to \tau^{>n}C \to 0$ whose morphisms induce isomorphisms

$$H^i(\tau^{\leqslant n}C) = \begin{cases} H^i(C) & i \leq n, \\ 0 & i > n, \end{cases} \qquad \text{and} \qquad H^i(\tau^{>n}C) = \begin{cases} 0 & i \leq n, \\ H^i(C) & i > n. \end{cases}$$

The exact sequence also implies there are natural quasi-isomorphisms [27, 1.7.5]

$$M(\tau^{\leqslant n}C \to C) \xrightarrow{\sim} \tau^{>n}C \qquad \text{and} \qquad \tau^{\leqslant n}C \xrightarrow{\sim} M(C \to \tau^{>n}C)[-1] \tag{5.1.1}$$

which suggests how to extend these functors to $\mathfrak{Mod}(\mathcal{L}_W)$.

5.2. For $Q \in \mathcal{P}(W)$ consider the composition $\mathcal{M} \to i_{Q*}i_Q^*\mathcal{M} \to i_{Q*}\tau^{>n}i_Q^*\mathcal{M}$ of natural maps (the first being the adjunction morphism implied by (3.4.1)). Define a functor $\tau_Q^{\leqslant n}$ on $\mathfrak{Mod}(\mathcal{L}_W)$ by

$$\tau_Q^{\leqslant n}\mathcal{M} = M(\mathcal{M} \to i_{Q*}\tau^{>n}i_Q^*\mathcal{M})[-1]\ ;$$

§3.8 shows there is a natural morphism $\tau_Q^{\leqslant n}\mathcal{M} \to \mathcal{M}$. By (5.1.1) there is a natural quasi-isomorphism

$$i_P^* \circ \tau_Q^{\leqslant n} \cong \begin{cases} \tau^{\leqslant n} \circ i_Q^* & \text{if } P = Q, \\ i_P^* & \text{if } P \nleq Q. \end{cases} \tag{5.2.1}$$

If $P < Q$ a simple formula like (5.2.1) does not hold. Instead there is a short exact sequence

$$0 \to H(\mathfrak{n}_P^Q; \tau^{>n}i_Q^*\mathcal{M}[-1]) \to i_P^*\tau_Q^{\leqslant n}\mathcal{M} \to i_P^*\mathcal{M} \to 0\ ;$$

since $H(\mathfrak{n}_P^Q;\cdot)$ is degree nondecreasing we at least have a quasi-isomorphism

$$\tau^{\leqslant n} \circ i_P^* \circ \tau_Q^{\leqslant n} \cong \tau^{\leqslant n} \circ i_P^*\ . \tag{5.2.2}$$

It is also clear that

$$\tau_P^{\leqslant n} \circ \tau_Q^{\leqslant n} = \tau_Q^{\leqslant n} \circ \tau_P^{\leqslant n} \quad \text{if } P \not< Q \text{ and } Q \not< P. \tag{5.2.3}$$

5.3. **Degree truncation of $\mathcal{L}$-modules.** The functor $\tau^{\leqslant n}$ on $\mathfrak{Mod}(\mathcal{L}_W)$ is defined as

$$\tau^{\leqslant n} = \tau_{Q_1}^{\leqslant n} \circ \tau_{Q_2}^{\leqslant n} \circ \cdots \circ \tau_{Q_N}^{\leqslant n} ,$$

where we write $\mathcal{P}(W) = \{Q_1, \dots, Q_N\}$ so that $Q_i \leq Q_j$ implies $i \leq j$; the functor is independent of the choice of ordering on $\mathcal{P}(W)$ by (5.2.3). There is a natural morphism $\tau^{\leqslant n}\mathcal{M} \to \mathcal{M}$. From (5.2.1)–(5.2.3) it follows that there is a quasi-isomorphism

$$i_Q^* \circ \tau^{\leqslant n} \cong \tau^{\leqslant n} \circ i_Q^* .$$

By applying i_Q^* and using (5.1.1) it is easy to verify the

Lemma. *Let W be an admissible space and let $\mathcal{S}_W \colon \mathfrak{Mod}(\mathcal{L}_W) \to \mathbf{D}_{\mathcal{X}}(W)$ be a realization functor from Theorem 4.1. Then $\mathcal{S}_W \circ \tau^{\leqslant n} \cong \tau^{\leqslant n} \circ \mathcal{S}_W$.*

Remark. The isomorphism does not hold on the level of complexes. The point is that $\tau^{\leqslant n}$ on $\mathcal{L}$-modules is defined "externally", via a mapping cone, whereas $\tau^{\leqslant n}$ on complexes of sheaves is defined "internally", as a sub-object.

5.4. **Intersection cohomology $\mathcal{L}$-module.** Let E be a regular G-module and let p be an ordinary perversity, that is, a function $p \colon \{2, \dots, \dim X\} \to \mathbb{Z}$ such that $p(2) = 0$, and $p(k+1) = p(k)$ or $p(k) + 1$. The *intersection cohomology $\mathcal{L}_{\widehat{X}}$-module $\mathcal{I}_p\mathcal{C}(E)$ (with perversity p and coefficients E)* is defined as follows. Set

$$\mathcal{M}_X = E$$

For an admissible space $W \subseteq \widehat{X}$ strictly containing X let $P \in \mathcal{P}(W)$ be minimal both with respect to the partial ordering on $\mathcal{P}(W)$ and to $\dim X_P$. Let $j_P \colon W \setminus X_P \hookrightarrow W$ denote the inclusion. Define the $\mathcal{L}_W$-module $\mathcal{M}_W$ inductively by setting

$$\mathcal{M}_W = \tau^{\leqslant p(\operatorname{codim}_{\widehat{X}} X_P)} j_{P*}\mathcal{M}_{W \setminus X_P} . \tag{5.4.1}$$

Finally set $\mathcal{I}_p\mathcal{C}(E) = \mathcal{M}_{\widehat{X}}$; there is a natural morphism $\mathcal{I}_p\mathcal{C}(E) \to i_{G*}E$. Note that since p is nondecreasing one may replace $\tau^{\leqslant p(\operatorname{codim}_{\widehat{X}} X_P)}$ in (5.4.1) by $\tau_P^{\leqslant p(\operatorname{codim}_{\widehat{X}} X_P)}$.

Let $\mathbb{E} = D \times_\Gamma E$ be the locally constant sheaf on X induced by E and let $\mathcal{I}_p\mathcal{C}(\widehat{X}; \mathbb{E})$ be the corresponding intersection cohomology sheaf [23] in $\mathbf{D}_{\mathcal{X}}(\widehat{X})$. The following proposition is immediate from the definitions and the fact that $\mathcal{S}_{\widehat{X}}$ commutes with the standard functors and with truncation (Theorem 4.1(ii) and Lemma 5.3).

Proposition. *There is a natural isomorphism $\mathcal{S}_{\widehat{X}}(\mathcal{I}_p\mathcal{C}(E)) \cong \mathcal{I}_p\mathcal{C}(\widehat{X}; \mathbb{E})$.*

5.5. **Local intersection cohomology.** We wish to give a formula for the local cohomology at P of $\mathcal{I}_p\mathcal{C}(E)$. There are two ingredients to the formula. The first is the nilpotent cohomology $H(\mathfrak{n}_P; E)$ and its decomposition $\bigoplus_{w \in W_P} H(\mathfrak{n}_P; E)_w$ from Kostant's theorem as in §0.10.21. The second ingredient is a certain combinatorial invariant $I_{p_w}H(c(|\Delta_P|))$, which we now describe.

Let $|\Delta_P|$ denote the geometric simplex with vertices indexed by the elements of Δ_P. It has a natural stratification given by the decomposition into open faces. The strata correspond to parabolic $\mathbb{Q}$-subgroups $Q \gneqq P$ (that is, to nonempty subsets $\Delta_P^Q \subseteq \Delta_P$) and are denoted $|\Delta_P^Q|^\circ$. We give the cone $c(|\Delta_P|)$ the induced stratification—the strata consist of the open cones on $|\Delta_P^Q|^\circ$ for $Q \gneqq P$ together with the cone point (indexed by P). Any constructible subset (that is, any union of open faces) of $|\Delta_P|$ or $c(|\Delta_P|)$ is given the induced stratification.

Given an ordinary perversity p and an element $w \in W$, define $p_w \colon \mathcal{P} \setminus \{G\} \to \mathbb{Z}$ by

$$p_w(Q) = p(\dim \mathfrak{n}_Q + \#\Delta_Q) - \ell(w_Q),$$

where w_Q denotes the projection of w to the second factor of $W = W^Q W_Q$. This defines an integer-valued function on the singular strata of $c(|\Delta_P|)$ or $|\Delta_P|$ (a perversity in the sense of [4]). Let $I_{p_w}H(U)$ denote the corresponding intersection cohomology group over $\mathbb{Z}$ for any open constructible subset U of $|\Delta_P|$ or $c(|\Delta_P|)$. (Unlike the intersection cohomology for an ordinary perversity p, the group here depends on a choice of stratification; we always use the stratification fixed above.) Note that if $P \leq Q$, then $w_Q = (w_P)_Q$ and so p_w and p_{w_P} agree on the strata of $|\Delta_P|$.

Proposition. *$H(i_P^* \mathcal{I}_p \mathcal{C}(E)) \cong \bigoplus_{w \in W_P} H(\mathfrak{n}_P; E)_w \otimes I_{p_w} H(c(|\Delta_P|))$ where L_P acts trivially on the second factor.*

Proof. There is a Mayer-Vietoris spectral sequence that abuts to the link cohomology $H(i_P^* j_{P*} j_P^* \mathcal{I}_p \mathcal{C}(E))$ (Lemma 3.7 with $Q' = G$ and $Q = P$) with

$$E_1^{p,\cdot} = \bigoplus_{\#\Delta_P^{\tilde{P}} = p+1} H(\mathfrak{n}_P^{\tilde{P}}; H(i_{\tilde{P}}^* \mathcal{I}_p \mathcal{C}(E))) \ .$$

By induction we may assume the proposition is true for $\tilde{P} > P$ and thus we can compute

$$\begin{aligned} E_1^{p,\cdot} &\cong \bigoplus_{\#\Delta_P^{\tilde{P}} = p+1} H(\mathfrak{n}_P^{\tilde{P}}; \bigoplus_{w_{\tilde{P}} \in W_{\tilde{P}}} H(\mathfrak{n}_{\tilde{P}}; E)_{w_{\tilde{P}}} \otimes I_{p_{w_{\tilde{P}}}} H(c(|\Delta_{\tilde{P}}|))) \\ &\cong \bigoplus_{\#\Delta_P^{\tilde{P}} = p+1} \bigoplus_{w_{\tilde{P}} \in W_{\tilde{P}}} \bigoplus_{w_P^{\tilde{P}} \in W_P^{\tilde{P}}} H(\mathfrak{n}_P^{\tilde{P}}; H(\mathfrak{n}_{\tilde{P}}; E)_{w_{\tilde{P}}})_{w_P^{\tilde{P}}} \otimes I_{p_{w_{\tilde{P}}}} H(c(|\Delta_{\tilde{P}}|)) \\ &\cong \bigoplus_{w \in W_P} H(\mathfrak{n}_P; E)_w \otimes \bigoplus_{\#\Delta_P^{\tilde{P}} = p+1} I_{p_w} H(c(|\Delta_{\tilde{P}}|)) \ . \end{aligned} \tag{5.5.1}$$

On the other hand, there is an analogous Mayer-Vietoris spectral sequence abutting to $I_{p_w}H(|\Delta_P|)$. Namely cover $|\Delta_P|$ by the open stars U_α of the vertices $\alpha \in \Delta_P$. For a parabolic $\mathbb{Q}$-subgroup $\tilde{P} > P$, the intersection $U_{\tilde{P}} = \bigcap_{\alpha \in \Delta_P^{\tilde{P}}} U_\alpha$ is the open star of the open face $|\Delta_P^{\tilde{P}}|^\circ$. For this spectral sequence

$$\widetilde{E}_1^{p,\cdot} = \bigoplus_{\#\Delta_P^{\tilde{P}} = p+1} I_{p_w} H(U_{\tilde{P}}) \cong \bigoplus_{\#\Delta_P^{\tilde{P}} = p+1} I_{p_w} H(c(|\Delta_{\tilde{P}}|)) \ . \tag{5.5.2}$$

Comparing (5.5.1) and (5.5.2) we see that $E_1^{p,\cdot} \cong \bigoplus_{w \in W_P} H(\mathfrak{n}_P; E)_w \otimes \widetilde{E}_1^{p,\cdot}$. This isomorphism induces isomorphisms at all stages of the spectral sequences and we find that

$$H(i_P^* j_{P*} j_P^* \mathcal{I}_p \mathcal{C}(E)) \cong \bigoplus_{w \in W_P} H(\mathfrak{n}_P; E)_w \otimes I_{p_w} H(|\Delta_P|) \ .$$

In order to obtain $H(i_P^* \mathcal{I}_p \mathcal{C}(E))$, this link cohomology is to be truncated in degrees greater than $p(\operatorname{codim}_{\widehat{X}} X_P) = \ell(w) + p_w(P)$, which corresponds exactly to replacing the second factor by $I_{p_w} H(c(|\Delta_P|))$. ☐

6. Example: the Weighted Cohomology $\mathcal{L}$-module

6.1. The reference for this subsection is [22, §9] though we give a slightly different description.

Let $P \in \mathcal{P}$ and let $X(S_P^G)$ denote the lattice of rational characters on S_P^G. For $\alpha \in \Delta_P$, let $R_\alpha \geq P$ be the maximal parabolic $\mathbb{Q}$-subgroup having type $\Delta_P \setminus \{\alpha\}$ with respect to P. We may canonically identify $X(S_{R_\alpha}^G) \cong \mathbb{Z}$ so that the unique element of Δ_{R_α} corresponds to a positive integer. Consider the map $X(S_P^G) \hookrightarrow \prod_{\alpha\in\Delta_P} X(S_{R_\alpha}^G) \cong \mathbb{Z}^{\Delta_P}$ defined by $\chi \mapsto (\chi|_{S_{R_\alpha}^G})_{\alpha\in\Delta_P}$. The image is a sublattice of finite index; after tensoring with $\mathbb{R}$ we have an isomorphism $\phi_P \colon \mathfrak{a}_P^{G*} = X(S_P^G) \otimes_{\mathbb{Z}} \mathbb{R} \xrightarrow{\sim} \mathbb{R}^{\Delta_P}$. For $P = P_0$ minimal we omit the subscript.

Define a partial ordering on $\mathfrak{a}_P^{G*}$ by declaring $\zeta \geq \eta$ if and only if $\phi_P(\zeta) \geq \phi_P(\eta)$ component-wise. Thus if ${}^+\mathfrak{a}_P^{G*}$ denotes the real convex cone generated by all $\alpha \in \Delta_P$, we have

$$\begin{aligned} \zeta \geq \eta \qquad &\iff \qquad \zeta - \eta \in {}^+\mathfrak{a}_P^{G*} \\ &\iff \qquad \langle \zeta, \beta^\vee \rangle \geq \langle \eta, \beta^\vee \rangle \qquad \text{for all } \beta \in \widehat{\Delta}_P. \end{aligned} \tag{6.1.1}$$

If $P \leq Q$ and $\zeta \in \mathfrak{a}_P^{G*}$, let $\zeta_Q = \zeta|_{\mathfrak{a}_Q^G}$ denote the restriction. We have a commutative diagram

$$\begin{tikzcd} \mathfrak{a}_P^{G*} \arrow[r, "\phi_P"] \arrow[d] & \mathbb{R}^{\Delta_P} \arrow[d] \\ \mathfrak{a}_Q^{G*} \arrow[r, "\phi_Q"] & \mathbb{R}^{\Delta_Q} \end{tikzcd}$$

where the left-hand vertical map is restriction and the right-hand vertical map is the projection $\mathbb{R}^{\Delta_P} = \mathbb{R}^{\Delta_P^Q} \times \mathbb{R}^{\Delta_P \setminus \Delta_P^Q} \to \mathbb{R}^{\Delta_P \setminus \Delta_P^Q} \cong \mathbb{R}^{\Delta_Q}$.

For a regular L_P-module E we write

$$E = \bigoplus_{\chi \in X(S_P^G)} E_\chi$$

where E_χ is the submodule on which S_P^G acts via χ. Given $\eta \in \mathfrak{a}^{G*}$, there are *weight truncation functors* on $\mathfrak{Mod}(L_P)$ defined by

$$\tau^{\geqslant\eta} E = \bigoplus_{\chi \geq \eta_P} E_\chi \quad \text{and} \quad \tau^{\not\geqslant\eta} E = \bigoplus_{\chi \not\geq \eta_P} E_\chi \ ;$$

the element η will be called a *weight profile*. (In [22] a weight profile is required to satisfy $\phi(\eta) \in (\mathbb{Z} + \frac{1}{2})^\Delta$; this ensures that any truncation functor $\tau^{\geqslant\eta}$ arises from a unique weight profile η. We will not assume this, but we will feel to replace η by η' provided $\tau^{\geqslant\eta}$ is unchanged.) Clearly we have a split short exact sequence $0 \to \tau^{\geqslant\eta} E \to E \to \tau^{\not\geqslant\eta} E \to 0$.

6.2. **Weight truncation of $\mathcal{L}$-modules.** Consider $P \leq Q \in \mathcal{P}$ and let E be a regular L_Q-module. The subtorus $S_Q^G \subseteq S_P^G$ acts on $H(\mathfrak{n}_P^Q; E_\chi)$ via the character χ. Thus

$$\tau^{\geqslant\eta} H(\mathfrak{n}_P^Q; E) \subseteq H(\mathfrak{n}_P^Q; \tau^{\geqslant\eta} E) \ . \tag{6.2.1}$$

Given a weight profile η we define local weight truncation functors $\tau_Q^{\geqslant\eta}$ on $\mathfrak{Mod}(\mathscr{L}_W)$ via a mapping cone

$$\tau_Q^{\geqslant\eta}\mathcal{M} = M(\mathcal{M} \to i_{Q*}\tau^{\not\geqslant\eta} i_Q^*\mathcal{M})[-1] \ ; \tag{6.2.2}$$

these are analogous to the local degree truncation functors in §5.2. The only change is in proving the analogue of (5.2.2). Instead of the fact that $H(\mathfrak{n}_P^Q;\cdot)$ is degree nondecreasing, we use that it is weight nonincreasing by (6.2.1); we also use that $\tau^{\geqslant\eta}$ is an exact functor.

The weight truncation functor $\tau^{\geqslant\eta}$ on $\mathfrak{Mod}(\mathscr{L}_W)$ is defined as a composition of these local truncation functors as in §5.3. There is a natural morphism $\tau^{\geqslant\eta}\mathcal{M} \to \mathcal{M}$ and a quasi-isomorphism

$$i_Q^* \circ \tau^{\geqslant\eta} \cong \tau^{\geqslant\eta} \circ i_Q^* \ . \tag{6.2.3}$$

6.3. **Weighted cohomology $\mathscr{L}$-module.** Let E be a regular G-module and let η be a weight profile. The *weighted cohomology* $\mathscr{L}_{\widehat{X}}$*-module* $\mathcal{W}^\eta\mathcal{C}(E)$ (*with weight profile* η *and coefficients* E) is defined as

$$\mathcal{W}^\eta\mathcal{C}(E) = \tau^{\geqslant\eta} i_{G*}E \ .$$

There is a natural morphism $\mathcal{W}^\eta\mathcal{C}(E) \to i_{G*}E$. Now let $\mathcal{W}^\eta\mathcal{C}(\widehat{X};\mathbb{E})$ be the corresponding weighted cohomology sheaf [22] in $\mathbf{D}_{\mathcal{X}}(\widehat{X})$.

Proposition. *There is a natural isomorphism* $\mathcal{S}_{\widehat{X}}(\mathcal{W}^\eta\mathcal{C}(E)) \cong \mathcal{W}^\eta\mathcal{C}(\widehat{X};\mathbb{E})$.

6.4. **Local weighted cohomology.** The formula for the local cohomology at P of $\mathcal{W}^\eta\mathcal{C}(E)$ is considerably simpler than that for $\mathcal{I}_p\mathcal{C}(E)$ since the former is defined by a simple truncation operation as opposed to the inductive truncation procedure of the latter. The following proposition is an immediate consequence of (6.2.3) and the definitions.

Proposition. $H(i_P^*\mathcal{W}^\eta\mathcal{C}(E)) \cong \tau^{\geqslant\eta}H(i_P^* i_{G*}E) \cong \tau^{\geqslant\eta}H(\mathfrak{n}_P;E)$.

6.5. *Remark.* In order to rephrase this in a form analogous to Proposition 5.5, let E be irreducible and have highest weight λ with respect to a Cartan subalgebra of $\mathfrak{l}_P$ and a positive system $\Phi^+(\mathfrak{g}_{\mathbb{C}},\mathfrak{h}_{\mathbb{C}})$ containing $\Phi(\mathfrak{n}_{P\mathbb{C}},h_{\mathbb{C}})$. Then by Kostant's theorem §0.10.21 we have

$$H(i_P^*\mathcal{W}^\eta\mathcal{C}(E)) \cong \bigoplus_{w\in W_P} H(\mathfrak{n}_P;E)_w \otimes \tau^{\geqslant\eta_P-(w(\lambda+\rho)-\rho)_P}\mathbb{C} \tag{6.5.1}$$

the second factor is either 0 or $\mathbb{C}$ with the trivial action of L_P. As Rapoport pointed out to me, the second factor here depends on both w and E, unlike $I_{p_w}H(c(|\Delta_P|))$ of Proposition 5.5 which is independent of E.

7. Micro-support of an $\mathscr{L}$-module

In this section we define the micro-support of an $\mathscr{L}$-module. This is an analogue of the notion for sheaves [27]; the analogy is not precise, partially because it is most natural here to restrict certain modules to be isomorphic to the complex conjugate of their contragredient.

7.1. **Notation.** Let W be an admissible space possessing a unique maximal stratum X_R and set

$$\mathfrak{Irr}(\mathcal{L}_W) = \coprod_{P\in\mathcal{P}(W)} \mathfrak{Irr}(L_P)\ .$$

For an L_P-module $V \in \mathfrak{Irr}(\mathcal{L}_W)$, define parabolic $\mathbb{Q}$-subgroups $Q'^W_V \geq Q^W_V \geq P$ by

$$\begin{aligned} \Delta_P^{Q^W_V} &= \{\, \alpha \in \Delta_P^R \mid \langle \xi_V + \rho, \alpha^\vee \rangle < 0 \,\}\ , \\ \Delta_P^{Q'^W_V} &= \{\, \alpha \in \Delta_P^R \mid \langle \xi_V + \rho, \alpha^\vee \rangle \leq 0 \,\}\ . \end{aligned}$$

Note that Q^W_V and Q'^W_V depend on W through the parabolic $\mathbb{Q}$-subgroup R indexing its maximal stratum. When $W = \widehat{X}$ we simply write Q_V and Q'_V.

7.2. **Definition.** The *weak micro-support* $\mathrm{SS}_w(\mathcal{M})$ of an $\mathcal{L}_W$-module $\mathcal{M}$ is the set of all $V \in \mathfrak{Irr}(\mathcal{L}_W)$ such that

(i) $H(i_P^* \hat{\imath}_Q^! \mathcal{M})_V \neq 0$ for some $Q \in [Q^W_V, Q'^W_V]$.

The *micro-support* $\mathrm{SS}_w(\mathcal{M})$ consists of those $V \in \mathrm{SS}_w(\mathcal{M})$ which satisfy in addition

(ii) $(V|_{M_P})^* \cong \overline{V|_{M_P}}$.

The *essential micro-support* $\mathrm{SS}_{\mathrm{ess}}(\mathcal{M})$ is the set of those $V \in \mathrm{SS}(\mathcal{M})$ such that

$$H(i_P^* \hat{\imath}_{Q^W_V}^! \mathcal{M})_V \longrightarrow H(i_P^* \hat{\imath}_{Q'^W_V}^! \mathcal{M})_V \tag{7.2.1}$$

is nonzero.

In §8 we will discuss condition (ii) further; this will lead in §9 to alternate forms of condition (i) and a clarification of the relationship between $\mathrm{SS}(\mathcal{M})$ and $\mathrm{SS}_{\mathrm{ess}}(\mathcal{M})$.

8. Conjugate Self-contragredient Modules and Fundamental Weyl Elements

We recall several results due to Borel and Casselman [9]. The first is an alternate form of Definition 7.2(ii) in terms of highest weights. The second relates this condition on certain irreducible components V of $H(\mathfrak{n}_P; E)$ to the condition on E.

8.1. **The involution $\boldsymbol{\tau_P}$.** Let P be a parabolic $\mathbb{R}$-subgroup and let V be an irreducible regular L_P-module. We say that V is *conjugate self-contragredient* if

$$V^* \cong \overline{V}$$

as representations of L_P. Clearly this is the case if and only if there exists a nondegenerate L_P-invariant sesquilinear form on V, however we will not use this characterization. Instead let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{l}_P$ and fix a positive system $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})$ of roots. Borel and Casselman [9, §1] construct an involution $\tau_P \colon \mathfrak{h}^*_\mathbb{C} \to \mathfrak{h}^*_\mathbb{C}$ that sends the highest weight of a representation of L_P to its complex conjugate contragredient. Thus V is conjugate self-contragredient if and only if

$$\tau_P(\mu) = \mu.$$

where μ is the highest weight of V.

To define τ_P, first let $\mathfrak{h}^*_\mathbb{R}$ be the real form of $\mathfrak{h}^*_\mathbb{C}$ spanned by the roots $\Phi(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})$ and the differentials of rational characters of the center of L_P. Let $s^P \in W^P$ be the element of the Weyl group of $\mathfrak{l}_P$ satisfying $s^P(-\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})) = \Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})$; similarly if c denotes complex conjugation of $\mathfrak{h}^*_\mathbb{C}$ with respect to the original real form $\mathfrak{h}^*$ let $s^c \in W^P$ satisfy $s^c(c\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})) = \Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})$. The automorphism $-s^P$ on

$\mathfrak{h}^*_{\mathbb{R}}$ transforms the highest weight of an irreducible L_P-module to that of its contragredient, while $s^c c$ on $\mathfrak{h}^*_{\mathbb{R}}$ transforms the highest weight of an irreducible L_P-module to that of its complex conjugate representation. Thus the desired automorphism of $\mathfrak{h}^*_{\mathbb{R}}$ is

$$\tau_P = (-s^P) \circ (s^c c) = (s^c c) \circ (-s^P)$$

and we extend this $\mathbb{C}$-linearly to $\mathfrak{h}^*_{\mathbb{C}}$. We let τ_P also denote the dual involution of $\mathfrak{h}_{\mathbb{C}}$.

The involution τ_P depends on the choice of the Cartan subalgebra $\mathfrak{h}$ and the positive system of roots. If $\tilde{\Phi}^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) = w\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ for $w \in W^P$, then the corresponding involution $\tilde{\tau}_P$ satisfies

$$\tilde{\tau}_P = w\tau_P w^{-1} \ . \tag{8.1.1}$$

8.2. **An alternate form of Definition 7.2(ii).** Now assume that P is defined over $\mathbb{Q}$. We may write $\mathfrak{h} = \mathfrak{b}_P + \mathfrak{a}_P$ where $\mathfrak{b}_P$ is a Cartan subalgebra of $\mathfrak{m}_P$. Note that τ_P acts by negation on $\mathfrak{a}_P$ and leaves $\mathfrak{b}_{P\mathbb{C}}$ invariant. Let $\mathfrak{b}_{P\mathbb{C}} = \mathfrak{b}_{P,1} + \mathfrak{b}_{P,-1}$ be the decomposition into the $+1$ and -1 eigenspaces of τ_P.

The following conditions are equivalent (where μ denotes the highest weight of V):

$$(V|_{M_P})^* \cong \overline{V|_{M_P}}, \text{ that is, } V|_{M_P} \text{ is conjugate self-contragredient,} \tag{8.2.1}$$

$$\tau_P(\mu|_{\mathfrak{b}_P}) = \mu|_{\mathfrak{b}_P}, \tag{8.2.2}$$

$$\mu|_{\mathfrak{b}_{P,-1}} = 0. \tag{8.2.3}$$

(In (8.2.2) and elsewhere we attempt to lighten the notation by writing $\mu|_{\mathfrak{b}_P}$ for $\mu|_{\mathfrak{b}_{P\mathbb{C}}}$.)

For later use we set $\tau'_P = -\tau_P$.

8.3. **Example.** Recall that $\rho = \rho^P + \rho_P$, where ρ^P is one half the sum of roots in $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ and ρ_P is one half the sum of roots in $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. The set $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ is W^P-invariant and thus $\rho_P|_{\mathfrak{b}_P} = 0$. On the other hand, $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ is τ_P-invariant and thus $\tau_P(\rho^P) = \rho^P$. Together these imply

$$\tau_P(\rho|_{\mathfrak{b}_P}) = \rho|_{\mathfrak{b}_P} \ . \tag{8.3.1}$$

8.4. **Fundamental Cartan subalgebras.** The familiar situation is when $\tau_P = \theta$ for some Cartan involution θ of $\mathfrak{l}_P$ and hence τ'_P acts on roots as complex conjugation. For a fixed P, this can always be arranged by choosing $\mathfrak{h}$ and $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ appropriately. Namely $\tau_P = \theta$ if and only if $\mathfrak{h}$ is a θ-stable Cartan subalgebra of $\mathfrak{l}_P$ and $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ is θ-stable. A θ-stable positive system exists if and only if there exists a regular element in the compact part of $\mathfrak{h}$ and hence if and only if the compact part of $\mathfrak{h}$ has maximal dimension. A Cartan subalgebra with this property for some Cartan involution is called *fundamental* (or *maximally compact*) for $\mathfrak{l}_P$; fundamental Cartan subalgebras always exist.

In the case that $\tau_P = \theta$ the decomposition $\mathfrak{b}_{P\mathbb{C}} = \mathfrak{b}_{P,1} + \mathfrak{b}_{P,-1}$ is defined over $\mathbb{R}$ and corresponds to the Cartan decomposition into compact and $\mathbb{R}$-split parts; the condition that $V|_{M_P}$ is conjugate self-contragredient is equivalent to μ vanishing on the $\mathbb{R}$-split part of $\mathfrak{b}_P$.

8.5. **Equal-rank groups.** A reductive algebraic group Z defined over $\mathbb{R}$ will be called *equal-rank* if $\mathbb{C}$-rank $Z = \operatorname{rank} K_Z$, where K_Z is a maximal compact subgroup of $Z(\mathbb{R})$. This holds if and only if a fundamental Cartan subalgebra is compact and thus if and only if every regular representation of Z is conjugate self-contragredient.

8.6. **Fundamental parabolic subgroups.** A Cartan subalgebra $\mathfrak{h}$ for $\mathfrak{l}_P$ may of course be identified with a Cartan subalgebra for $\mathfrak{g}$ via a lift; we will always assume this has been done. Consider the condition on a parabolic $\mathbb{R}$-subgroup P that a fundamental Cartan subalgebra $\mathfrak{h}$ for $\mathfrak{l}_P$ is also fundamental for $\mathfrak{g}$. Such parabolic $\mathbb{R}$-subgroups exist and a minimal one is called *fundamental*. If P_0 is a fundamental parabolic $\mathbb{R}$-subgroup, then $L_{P_0}/_{\mathbb{R}}S_{P_0}$ is equal-rank; $P_0 \neq G$ if and only if $G/_{\mathbb{R}}S_G$ is not equal-rank. Various conditions equivalent to P containing a fundamental parabolic $\mathbb{R}$-subgroup are given in [9, 1.8]; here are some others more suitable for our purposes.

Lemma. *Let P be a parabolic $\mathbb{R}$-subgroup of G and fix a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{l}_P$. Let $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ be a positive system for $\mathfrak{l}_{P\mathbb{C}}$ and extend it to a positive system $\Phi^+ = \Phi^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ containing $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. Then the following conditions are equivalent*:

- (i) *P contains a fundamental parabolic $\mathbb{R}$-subgroup*;
- (ii) *There exists $w \in W_P$ such that $w\tau_G w^{-1} = \tau_P$*;
- (iii) *There exists $w \in W_P$ such that τ_P' interchanges Φ_w and $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) \setminus \Phi_w$, where $\Phi_w = \{\, \gamma \in \Phi^+ \mid w^{-1}\gamma < 0 \,\}$*;
- (iv) *There exists $w \in W_P$ such that $w\Phi^+$ is τ_P-stable.*

The conditions in (ii), (iii) *and* (iv) *for a given w are equivalent.*

Proof. Let τ_P and τ_G be defined as in §8.1 with respect to $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ and Φ^+ respectively. We choose a Cartan involution θ for $\mathfrak{g}$ which induces one on $\mathfrak{l}_P$ via a θ-stable lift. We can assume that $\mathfrak{h}$ is a fundamental θ-stable Cartan subalgebra for $\mathfrak{l}_P$ and (by conjugating by an element of W^P) that $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ is θ-stable and thus that $\tau_P = \theta$.

(i)⇒(ii): Suppose that P contains a fundamental parabolic $\mathbb{R}$-subgroup. Then $\mathfrak{h}$ is a fundamental Cartan subalgebra for $\mathfrak{g}$ as well so there exists a θ-stable positive system $\tilde{\Phi}^+$ for $\mathfrak{g}$ which contains $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. Let $\tilde{\tau}_G$ denote the operator of §8.1 with respect to $\tilde{\Phi}^+$; since $\tilde{\Phi}^+$ is θ-stable it satisfies $\tilde{\tau}_G = \theta = \tau_P$, while if $\tilde{\Phi}^+ = w\Phi^+$ for $w \in W$ then $\tilde{\tau}_G = w\tau_G w^{-1}$. Since both positive systems contain $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$, w must lie in W_P.

(ii)⇒(iii): Suppose $w\tau_G w^{-1} = \tau_P$ for some $w \in W_P$. Then $\tilde{\tau}_G = \tau_P$ where $\tilde{\tau}_G$ is with respect to $\tilde{\Phi}^+ = w\Phi^+$. Decompose $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) = \Phi_w \cup (\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) \setminus \Phi_w)$. Observe that these subsets correspond to those roots that are negative (respectively positive) with respect to $\tilde{\Phi}^+$. However τ_P' preserves $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ and since $\tau_P' = \tilde{\tau}_G'$ it must interchange the two subsets.

(iii)⇒(iv): If w is as in (iii), the positive system $w\Phi^+ = \Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) \cup -\Phi_w \cup (\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) \setminus \Phi_w)$ is τ_P-stable.

(iv)⇒(i): Since $w\Phi^+$ is θ-stable, $\mathfrak{h}$ is fundamental for $\mathfrak{g}$. □

8.7. **Fundamental Weyl elements.** An element $w \in W_P$ satisfying $w\tau_G w^{-1} = \tau_P$ as in Lemma 8.6(ii) will be called a *fundamental Weyl element* (for P in G). More generally, if $P \leq R$ an element $w \in W_P^R$ is fundamental for P in R if $w\tau_R w^{-1} = \tau_P$

(equivalently it is fundamental for P/N_R in L_R). Besides the equivalent formulations given in Lemma 8.6(iii) and (iv), here are two basic facts about fundamental Weyl elements:

(i) A fundamental Weyl element $w \in W_P$ satisfies $\ell(w) = \frac{1}{2}\dim \mathfrak{n}_P$.

(ii) If $P \leq R$ and $w = w^R w_R \in W_P^R W_R = W_P$, then w is fundamental for P in G if and only if w^R is fundamental for P in R and w_R is fundamental for R in G.

Fact (i) is clear from Lemma 8.6(iii). For (ii), note that τ_P leaves $\Phi(\mathfrak{l}_{R\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ invariant and thus if $w\Phi^+$ is τ_P-stable, so is $w^R\Phi^+(\mathfrak{l}_{R\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) = w\Phi^+ \cap \Phi(\mathfrak{l}_{R\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. Hence by Lemma 8.6(iv), $w\tau_G w^{-1} = \tau_P$ implies $w^R\tau_R(w^R)^{-1} = \tau_P$ and consequently also $w_R\tau_G w_R^{-1} = \tau_R$. The converse is clear.

8.8**. Lemma.** *Let E be an irreducible regular G-module and let P be a parabolic subgroup of G. The following conditions are equivalent*:

(i) *P contains a fundamental parabolic $\mathbb{R}$-subgroup and $(E|_{{}^0G})^* \cong \overline{E|_{{}^0G}}$;*

(ii) *There exists an irreducible constituent V of $H(\mathfrak{n}_P; E)$ satisfying $(\xi_V + \rho)|_{\mathfrak{a}_P^G} = 0$ and $(V|_{M_P})^* \cong \overline{V|_{M_P}}$.*

If either condition holds, $V = H^{\ell(w)}(\mathfrak{n}_P; E)_w$ for $w \in W_P$ a fundamental Weyl element for P in G and (ii) *holds for any such V.*

Proof. Let $\mathfrak{h}$ be a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{l}_P$ and fix positive orderings for the roots of $\mathfrak{l}_{P\mathbb{C}}$ and $\mathfrak{g}_{\mathbb{C}}$ as in Lemma 8.6. Let E have highest weight λ; by Kostant's theorem, for all $w \in W_P$ the L_P-module $V = H^{\ell(w)}(\mathfrak{n}_P; E)_w$ has highest weight $\mu = w(\lambda+\rho) - \rho$. One may use §§8.2 and 8.3 to translate (i) and (ii) into assertions regarding w and λ; their equivalence in this form is proved in [9, 3.6(iii)(iv)] (one must replace λ in [9] by $\lambda + \rho$ to agree with our notation). For the final assertion, note that since τ_P preserves the decomposition $\mathfrak{b}_G = \mathfrak{b}_P + \mathfrak{a}_P^G$ and acts as -1 on the second factor, (ii) holds for a given w if and only if $\tau_P w(\lambda+\rho)|_{\mathfrak{b}_G} = w(\lambda+\rho)|_{\mathfrak{b}_G}$ and hence if and only if $w^{-1}\tau_P w((\lambda+\rho)|_{\mathfrak{b}_G}) = (\lambda+\rho)|_{\mathfrak{b}_G}$. Since $w^{-1}\tau_P w$ and τ_G are members of W after composing with $-c$ and they agree on a regular element, they must be equal. (In fact this argument may be completed to directly prove the equivalence of (i) and (ii).) □

9. Alternate Forms of Definition 7.2(i)

Let W be an admissible space possessing a unique maximal stratum X_R.

9.1. **A partial ordering on $\mathfrak{Irr}(\mathcal{L}_W)$.** Let V, $\tilde{V} \in \mathfrak{Irr}(\mathcal{L}_W)$ be irreducible L_P- and $L_{\tilde{P}}$-modules respectively. Set $V \preccurlyeq_0 \tilde{V}$ if the following three conditions are fulfilled:

(i) $P \leq \tilde{P}$;

(ii) $V = H^{\ell(w)}(\mathfrak{n}_P^{\tilde{P}}; \tilde{V})_w$ for an element $w \in W_P^{\tilde{P}}$;

(iii) $(\xi_V + \rho)|_{\mathfrak{a}_P^{\tilde{P}}} = 0$.

From §0.10.22 it is easy to see that this is a partial order. If $V \preccurlyeq_0 \tilde{V}$ we set $[\tilde{V} : V] = \ell(w)$, where w is as in (ii). (We will consider the partial ordering obtained by relaxing (iii) later in §22.3.)

Lemma 8.8 immediately implies the following

Lemma. *If $V \preccurlyeq_0 \tilde{V}$ and $(V|_{M_P})^* \cong \overline{V|_{M_P}}$ then $(\tilde{V}|_{M_{\tilde{P}}})^* \cong \overline{\tilde{V}|_{M_{\tilde{P}}}}$, the Weyl element w from* (ii) *above is fundamental for P in $\tilde{P}$, and $[\tilde{V} : V] = \frac{1}{2}\dim \mathfrak{n}_P^{\tilde{P}}$.*

9.2**. Proposition.** *Let $\mathcal{M}$ be an $\mathfrak{L}_W$-module and let $V \in \mathfrak{Irr}(\mathfrak{L}_W)$ be an irreducible L_P-module. The following conditions on V are equivalent:*

(i) $H(i_P^* \hat{\imath}_Q^! \mathcal{M})_V \neq 0$ *for some* $Q \in [Q_V^W, Q_V'^W]$.

(ii) $H(i_P^* i_{\tilde{P}*} i_{\tilde{P}}^* \hat{\imath}_{\tilde{Q}}^! \mathcal{M})_V \neq 0$ *for some* $\tilde{Q} \in Q_V^W, Q_V'^W]$ *and* $\tilde{P} = (P, Q_V^W) \cap \tilde{Q}$.

(iii) *There exists an $L_{\tilde{P}}$-module $\tilde{V} \succcurlyeq_0 V$ such that $H(i_{\tilde{P}}^* \hat{\imath}_{Q_{\tilde{V}}^W}^! \mathcal{M})_{\tilde{V}} \neq 0$.*

(iv) *There exists an $L_{\tilde{\tilde{P}}}$-module $\tilde{\tilde{V}} \succcurlyeq_0 V$ such that*

$$\operatorname{Im}\bigl(H(i_{\tilde{\tilde{P}}}^* \hat{\imath}_{Q_{\tilde{\tilde{V}}}^W}^! \mathcal{M})_{\tilde{\tilde{V}}} \longrightarrow H(i_{\tilde{\tilde{P}}}^* \hat{\imath}_{Q_{\tilde{\tilde{V}}}'^W}^! \mathcal{M})_{\tilde{\tilde{V}}}\bigr) \neq 0 \ .$$

If these conditions hold it can be arranged that $\tilde{P}$ in (ii) *and* (iii) *is the same and that $\tilde{\tilde{P}} \geq \tilde{P}$. Furthermore, let $[c_i(\cdot;\mathcal{M}), d_i(\cdot;\mathcal{M})]$ $(i = 1, \dots, 4)$ denote the range of degrees in which the expressions in* (i)–(iv) *respectively can be nonvanishing, where in the parentheses we indicate the module and, if desired, the parabolic $\mathbb{Q}$-subgroup to which we wish to restrict. Then*

$$\begin{aligned} &[c_1(V;\mathcal{M}), d_1(V;\mathcal{M})] \subseteq [c_2(V;\mathcal{M}), d_2(V;\mathcal{M})] \ , \\ &[c_2(V,\tilde{P};\mathcal{M}), d_2(V,\tilde{P};\mathcal{M})] = [c_3(\tilde{V};\mathcal{M}), d_3(\tilde{V};\mathcal{M})] + \tfrac{1}{2}\dim \mathfrak{n}_P^{\tilde{P}} \ , \\ &[c_3(\tilde{V};\mathcal{M}), d_3(\tilde{V};\mathcal{M})] \subseteq [c_4(\tilde{\tilde{V}};\mathcal{M}), d_4(\tilde{\tilde{V}};\mathcal{M}) + \dim \mathfrak{a}_{\tilde{P}}^{\tilde{\tilde{P}}}] + \tfrac{1}{2}\dim \mathfrak{n}_{\tilde{P}}^{\tilde{\tilde{P}}} \ . \end{aligned} \tag{9.2.1}$$

Proof. (i)$\Longleftrightarrow$(ii): Assume (i) holds and consider the long exact sequence

$$\cdots \longrightarrow H^j(i_P^* \hat{\imath}_{Q_V^W}^! \mathcal{M})_V \xrightarrow{\beta} H^j(i_P^* \hat{\imath}_Q^! \mathcal{M})_V \longrightarrow H^j(i_P^* \hat{\jmath}_{Q_V^W *} \hat{\jmath}_{Q_V^W}^* \hat{\imath}_Q^! \mathcal{M})_V \longrightarrow \cdots \tag{9.2.2}$$

as in (3.6.4). If β is surjective, then $H(i_P^* \hat{\imath}_{Q_V^W}^! \mathcal{M})_V \neq 0$ and (ii) holds with $\tilde{Q} = Q_V^W$ and $\tilde{P} = P$. Otherwise the last term of (9.2.2) is nonzero which implies by the Fary spectral sequence of Lemma 3.7 and equation (3.6.6) that (ii) holds for some $\tilde{Q} \leq Q$ with $\tilde{P} > P$. Conversely suppose (ii) holds for some $\tilde{Q}$ and consider the analogous long exact sequence to (9.2.2) in which Q has been replaced by $\tilde{Q}$. By (ii) the Fary spectral sequence for the last term has E_1 nonzero at the top level. Thus either the last term is nonzero, in which case (i) holds for Q_V^W or $\tilde{Q}$, or else the last term is zero, in which case (ii) holds also for some smaller $\tilde{Q}$ and we may repeat the argument.

(ii)$\Longleftrightarrow$(iii): Assume that (ii) holds. Then by (3.6.6) we see that $H(i_{\tilde{P}}^* \hat{\imath}_{\tilde{Q}}^! \mathcal{M})_{\tilde{V}} \neq 0$ for some irreducible $L_{\tilde{P}}$-module $\tilde{V}$ such that V is an irreducible constituent of $H(\mathfrak{n}_P^{\tilde{P}}; \tilde{V})$. By the definition of $\tilde{P}$ we see that $(\xi_V + \rho)|_{\mathfrak{a}_P^{\tilde{P}}} = 0$ and so $V \preccurlyeq_0 \tilde{V}$. Finally $(\xi_V + \rho)|_{\mathfrak{a}_{\tilde{P}}} = \xi_{\tilde{V}} + \rho_{\tilde{P}}$ implies that $\tilde{Q} = Q_{\tilde{V}}^W$ and thus (iii) holds. Conversely (iii) implies $H(i_P^* i_{\tilde{P}*} i_{\tilde{P}}^* \hat{\imath}_{Q_{\tilde{V}}^W}^! \mathcal{M})_V \neq 0$. We have $Q_{\tilde{V}}^W = Q_V^W \vee \tilde{P}$ and $Q_{\tilde{V}}'^W = Q_V'^W$ so $Q_{\tilde{V}}^W \in [Q_V^W, Q_V'^W]$ and hence (ii) holds.

(iii)$\Longleftrightarrow$(iv): Assume that (iii) holds and consider the long exact sequence

$$\cdots \longrightarrow H^{j-1}(i_{\tilde{P}}^* \hat{\jmath}_{Q_{\tilde{V}}^W *} \hat{\jmath}_{Q_{\tilde{V}}^W}^* \hat{\imath}_{Q_{\tilde{V}}'^W}^! \mathcal{M})_{\tilde{V}} \longrightarrow H^j(i_{\tilde{P}}^* \hat{\imath}_{Q_{\tilde{V}}^W}^! \mathcal{M})_{\tilde{V}} \xrightarrow{\beta} H^j(i_{\tilde{P}}^* \hat{\imath}_{Q_{\tilde{V}}'^W}^! \mathcal{M})_{\tilde{V}} \longrightarrow \cdots \ .$$

Either β is nonzero, in which case (iv) holds, or the first term of the sequence is nonzero. By Lemma 3.7 and equation (3.6.6) this implies that (ii) and hence (iii) holds for a larger $\tilde{P}$ and we may repeat the argument. Obviously (iv) $\Longrightarrow$ (iii). $\square$

9.3. **Corollary.** *If* $V \in \operatorname{SS}(\mathcal{M})$ *there exists* $\tilde{V} \in \operatorname{SS}_{\mathrm{ess}}(\mathcal{M})$ *with* $V \preccurlyeq_0 \tilde{V}$. *If* $\tilde{V} \in \operatorname{SS}(\mathcal{M})$ *and* $V \in \mathfrak{Irr}(\mathcal{L}_W)$ *is an irreducible* L_P*-module for which* $V|_{M_P}$ *is conjugate self-contragredient and* $V \preccurlyeq_0 \tilde{V}$, *then* $V \in \operatorname{SS}(\mathcal{M})$.

Thus $\operatorname{SS}_{\mathrm{ess}}(\mathcal{M})$ determines $\operatorname{SS}(\mathcal{M})$; the converse is not true however—this is because the implication (ii) $\Longrightarrow$ (i) does not determine Q.

Part II. A Global Vanishing Theorem for $\mathcal{L}$-modules

In this part we present a vanishing theorem for the global cohomology of an $\mathcal{L}$-module $\mathcal{M}$. The well-known local-global principle implies that if the local cohomology of a sheaf vanishes, then the global hypercohomology vanishes. Our result implies that if the micro-support of an $\mathcal{L}$-module is empty, then the global cohomology vanishes. More generally we give a bound based on the micro-support on the degrees in which cohomology can be nonzero.

10. The Vanishing Theorem

10.1. Let $P \in \mathcal{P}$ and let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{l}_P$. Given $\mu \in \mathfrak{h}_{\mathbb{C}}^*$ view μ as an element of $\mathfrak{l}_{P\mathbb{C}}^*$ by extending μ to be zero on all root spaces. Let $L_P(\mu) \subseteq L_P$ be the stabilizer of μ under the coadjoint action. This is a reductive subgroup (defined over $\mathbb{C}$) whose Lie algebra $\mathfrak{l}_P(\mu)$ is generated by $\mathfrak{h}$ and the root spaces for

$$\{\, \gamma \in \Phi(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) \mid \langle \mu, \gamma^\vee \rangle = 0 \,\} \; . \tag{10.1.1}$$

As extreme cases we have $L_P(0) = L_P$ and $L_P(\rho) = H$, the Cartan subgroup.

10.2. Let V be an irreducible regular L_P-module for which $(V|_{M_P})^* \cong \overline{V|_{M_P}}$. Choose $\mathfrak{h}$ and Φ^+ so that $\tau_P = \theta$ for some Cartan involution and let $\mu \in \mathfrak{h}_{\mathbb{C}}^*$ be the corresponding highest weight of V. In this case $\overline{\mu|_{\mathfrak{b}_P}} = -\mu|_{\mathfrak{b}_P}$ and so $L_P(\mu)$ is defined over $\mathbb{R}$. Let

$$D_P(\mu) = L_P(\mu)(\mathbb{R})/(K_P \cap L_P(\mu))A_P$$

denote the corresponding symmetric space (compare [7]). The dimension of $D_P(\mu)$ can vary depending upon the θ-stable positive system Φ^+; we let $D_P(V)$ denote any one of the $D_P(\mu)$ with maximal dimension. Then $\dim D_P(V)$ is well-defined and independent of the choice of $\mathfrak{h}$ and θ. It will be convenient beginning in §25 to allow V in $D_P(V)$ to be merely isotypical as opposed to irreducible.

10.3. Let $\mathcal{M}$ be an $\mathcal{L}$-module on $\widehat{X}$. For $V \in \operatorname{SS}(\mathcal{M})$ let $c(V;\mathcal{M}) \leq d(V;\mathcal{M})$ be the least and greatest degrees in which $H^i(i_P^* \hat{\imath}_Q^! \mathcal{M})_V \neq 0$ (for any $Q \in [Q_V, Q'_V]$). Define

$$c(\mathcal{M}) = \inf_{V \in \operatorname{SS}(\mathcal{M})} \tfrac{1}{2}(\dim D_P - \dim D_P(V)) + c(V;\mathcal{M}) \; , \text{ and} \tag{10.3.1}$$

$$d(\mathcal{M}) = \sup_{V \in \operatorname{SS}(\mathcal{M})} \tfrac{1}{2}(\dim D_P + \dim D_P(V)) + d(V;\mathcal{M}) \; . \tag{10.3.2}$$

Note that the shift $\frac{1}{2}(\dim D_P \pm \dim D_P(V))$ corresponds to the range of degrees in which $H_{(2)}(X_P;\mathbb{V})$ can be nonzero (see Theorem 14.1).

10.4. **Theorem.** *Let $\mathcal{M}$ be an $\mathcal{L}$-module on $\widehat{X}$. Then $H^i(\widehat{X};\mathcal{M}) = 0$ for $i \notin [c(\mathcal{M}), d(\mathcal{M})]$.*

The proof will appear in §15 after a number of analytic preliminaries. In the remainder of this section we will indicate how $c(\mathcal{M})$ and $d(\mathcal{M})$ may be computed in terms of the essential micro-support.

10.5. **Lemma.** *Let V be an irreducible regular L_P-module for $P \in \mathcal{P}$ and assume $V|_{M_P}$ is conjugate self-contragredient. If $V \preccurlyeq_0 \tilde{V}$ then*

$$\dim D_P(V) \le \dim D_{\tilde{P}}(\tilde{V}) - \dim \mathfrak{a}_P^{\tilde{P}} \ . \tag{10.5.1}$$

Consequently

$$\tfrac12\bigl(\dim D_P - \dim D_P(V)\bigr) \ge \tfrac12\bigl(\dim D_{\tilde{P}} - \dim D_{\tilde{P}}(\tilde{V})\bigr) - \tfrac12 \dim \mathfrak{n}_P^{\tilde{P}} \ , \tag{10.5.2}$$

$$\begin{aligned}\tfrac12\bigl(\dim D_P + \dim D_P(V)\bigr) \le \tfrac12\bigl(\dim D_{\tilde{P}} + \dim D_{\tilde{P}}(\tilde{V})\bigr) - \tfrac12 \dim \mathfrak{n}_P^{\tilde{P}}\\ - \dim \mathfrak{a}_P^{\tilde{P}} \ .\end{aligned} \tag{10.5.3}$$

Proof. Let $\mathfrak{h}$ be a θ-stable Cartan subalgebra for $\mathfrak{l}_P$ and hence for $\mathfrak{l}_{\tilde{P}}$. Choose $\mathfrak{h}$ and Φ^+ so that $\tau_P = \theta$ and let μ be the corresponding highest weight of V. Then we can define $D_P(\mu)$ to be the symmetric space of $L_P(\mu)(\mathbb{R})$ as in §10.2. Now by hypothesis and Lemma 8.8 we know that $V = H^{\ell(w)}(\mathfrak{n}_P^{\tilde{P}}; \tilde{V})_w$ for $w \in W_P^{\tilde{P}}$ such that $w\tau_{\tilde{P}}w^{-1} = \tau_P$. Then $\widetilde{\tau}_{\tilde{P}} = \theta$, where $\widetilde{\tau}_{\tilde{P}} = w\tau_{\tilde{P}}w^{-1}$ is the operator of §8.1 defined with respect to the positive system $w\Phi^+$. So if $\tilde{\mu}$ is the highest weight of $\tilde{V}$ with respect to Φ^+, we can define $D_{\tilde{P}}(w\tilde{\mu})$ to be the symmetric space of $L_{\tilde{P}}(w\tilde{\mu})(\mathbb{R})$ as in §10.2.

Now $\mu = w(\tilde{\mu} + \rho) - \rho$ by Kostant's theorem §0.10.21. So if γ is a root of $\mathfrak{l}_{P\mathbb{C}}$ and $\langle \mu, \gamma^\vee\rangle = 0$ then $\langle w\tilde{\mu}, \gamma^\vee\rangle = 0$ as well, since both $w\tilde{\mu}$ and $w\rho - \rho$ are dominant for $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. Thus $L_P(\mu) \subseteq L_{\tilde{P}}(w\tilde{\mu})$ and hence $\dim D_P(\mu) \le \dim D_{\tilde{P}}(w\tilde{\mu}) - \dim \mathfrak{a}_P^{\tilde{P}}$; the subtraction of $\dim \mathfrak{a}_P^{\tilde{P}}$ adjusts for taking quotient by A_P on the left and $A_{\tilde{P}}$ on the right. The lemma follows by taking the maximum over possible Φ^+ and by applying $\dim D_P = \dim D_{\tilde{P}} - \dim \mathfrak{a}_P^{\tilde{P}} - \dim \mathfrak{n}_P^{\tilde{P}}$. $\square$

10.6. Let $\mathcal{M}$ be an $\mathcal{L}$-module on $\widehat{X}$. For $V \in \operatorname{SS}(\mathcal{M})$ let $[c_i(V;\mathcal{M}), d_i(V;\mathcal{M})]$ ($i = 1, \dots, 4$) be as in Proposition 9.2. Define $c_i(\mathcal{M})$ and $d_i(\mathcal{M})$ analogously to (10.3.1) and (10.3.2) with the exception that we set

$$d_2(\mathcal{M}) = \sup_{V \in \operatorname{SS}(\mathcal{M}), \tilde{P}} \tfrac12(\dim D_P + \dim D_P(V)) + \dim \mathfrak{a}_P^{\tilde{P}} + d_2(V, \tilde{P}; \mathcal{M}) \ .$$

Of course $[c(\mathcal{M}), d(\mathcal{M})] = [c_1(\mathcal{M}), d_1(\mathcal{M})]$. In fact we have

Proposition. *The intervals $[c_i(\mathcal{M}), d_i(\mathcal{M})]$ are equal for $i = 1, \dots, 4$.*

Proof. We will prove all $d_i(\mathcal{M})$ are equal; the proof of the other equalities is similar. For $V \in \operatorname{SS}(\mathcal{M})$ let $V \preccurlyeq_0 \tilde{V} \preccurlyeq_0 \tilde{\tilde{V}}$ be chosen as in Proposition 9.2 such that $d_2(V;\mathcal{M}) = d_2(V, \tilde{P}; \mathcal{M})$. Then

$$\begin{aligned} d_1(V;\mathcal{M}) \le d_2(V, \tilde{P}; \mathcal{M}) &= d_3(\tilde{V};\mathcal{M}) + \tfrac12 \dim \mathfrak{n}_P^{\tilde{P}} \\ &\le d_4(\tilde{\tilde{V}};\mathcal{M}) + \dim \mathfrak{a}_{\tilde{P}}^{\tilde{\tilde{P}}} + \tfrac12 \dim \mathfrak{n}_P^{\tilde{\tilde{P}}} \end{aligned}$$

by (9.2.1). These inequalities together with (10.5.3) (applied both to $V \preccurlyeq_0 \tilde{V}$ and $\tilde{V} \preccurlyeq_0 \tilde{\tilde{V}}$) yield $d_1(\mathcal{M}) \le d_2(\mathcal{M}) \le d_3(\mathcal{M}) \le d_4(\mathcal{M})$. On the other hand, for

$V \in \mathrm{SS}_{\mathrm{ess}}(\mathcal{M})$ the obvious inequality $d_4(V;\mathcal{M}) \le d_1(V;\mathcal{M})$ implies that $d_4(\mathcal{M}) \le d_1(\mathcal{M})$. ☐

Corollary. *For* $V \in \mathrm{SS}_{\mathrm{ess}}(\mathcal{M})$ *let* $c_{\mathrm{ess}}(V;\mathcal{M}) \le d_{\mathrm{ess}}(V;\mathcal{M})$ *be the least and greatest degrees in which*

$$\operatorname{Im}\bigl(H(i_P^*\hat{i}^!_{Q_V^W}\mathcal{M})_V \longrightarrow H(i_P^*\hat{i}^!_{Q_V'^W}\mathcal{M})_V\bigr) \ne 0 \ .$$

Then

$$\begin{aligned} c(\mathcal{M}) &= \inf_{V\in \mathrm{SS}_{\mathrm{ess}}(\mathcal{M})} \tfrac{1}{2}(\dim D_P - \dim D_P(V)) + c_{\mathrm{ess}}(V;\mathcal{M}) \ , \quad \textit{and} \\ d(\mathcal{M}) &= \sup_{V\in \mathrm{SS}_{\mathrm{ess}}(\mathcal{M})} \tfrac{1}{2}(\dim D_P + \dim D_P(V)) + d_{\mathrm{ess}}(V;\mathcal{M}) \ . \end{aligned}$$

11. L^2-COHOMOLOGY WITH BOUNDARY VALUES

In this section we introduce a variant of L^2-cohomology whose elements admit restrictions to boundary faces.

11.1. L^2-cohomology.

We first recall ordinary L^2-cohomology; references are [18] and [56]. Let Y be a Riemannian manifold (possibly with boundary or corners) and let $\mathbb{E}$ be a metrized locally constant sheaf. For a measurable $\mathbb{E}$-valued form ω on Y let

$$\|\omega\| = \|\omega\|_Y = \left(\int_Y |\omega|^2\, dV\right)^{\frac{1}{2}}$$

denote the usual L^2 norm (we will omit the subscript Y if this would not cause confusion). Let $L_2(Y;\mathbb{E})$ be the Hilbert space of those forms ω for which $\|\omega\| < \infty$; equivalently this is the completion of $A_c(Y;\mathbb{E})$ with respect to the L^2 norm. Let $d = d_Y = d_{Y,\mathbb{E}}$ denote the unbounded densely defined operator given by exterior differentiation on the domain

$$\operatorname{Dom} d = \{\, \omega \in A(Y;\mathbb{E}) \mid \|\omega\|, \|d\omega\| < \infty \,\}$$

and let $\overline{d}$ be its closure in the graph norm. That is, $\omega \in L_2(Y;\mathbb{E})$ is in $\operatorname{Dom}\overline{d}$ if and only if there exists a sequence $\omega_i \in \operatorname{Dom} d$ such ω_i and $d\omega_i$ are Cauchy sequences and $\omega_i \to \omega$ in L^2 norm. The *L^2-cohomology* $H_{(2)}(Y;\mathbb{E})$ is defined to be the cohomology of the complex $\operatorname{Dom}\overline{d}$. A smoothing argument [19, §15], [18, §8] shows that the inclusion $\operatorname{Dom} d \subseteq \operatorname{Dom}\overline{d}$ induces an isomorphism $H_{(2)}(Y;\mathbb{E}) \cong H(\operatorname{Dom} d)$. Consequently

$$H_{(2)}(Y;\mathbb{E}) \cong H(Y;\mathbb{E}) \qquad \text{if } Y \text{ is compact.} \tag{11.1.1}$$

We give $H_{(2)}(Y;\mathbb{E})$ the quotient topology; this is Hausdorff if and only if $\operatorname{Range}\overline{d}$ is closed. If it is Hausdorff, then $H_{(2)}(Y;\mathbb{E})$ is representable by harmonic forms, namely $\operatorname{Ker}\overline{d} \cap \operatorname{Ker}\overline{d}^*$ where $\overline{d}^*$ is the adjoint of $\overline{d}$ in the sense of unbounded operators on the Hilbert space $L_2(Y;\mathbb{E})$; conversely if it is not Hausdorff the L^2-cohomology is infinite dimensional.

If $g > 0$ is a positive function on Y one may also consider the weighted norm $\|\omega\|_g^2 = \int_Y |\omega|^2\, g dV$, the corresponding Hilbert space $L_2(Y;\mathbb{E},g)$, and the L^2-cohomology $H_{(2)}(Y;\mathbb{E},g)$. Since this case may be absorbed into the previous by modifying the metric on $\mathbb{E}$, we do not mention it again in this section.

11.2. **Spectral vanishing criterion for $H_{(2)}(Y;\mathbb{E})$.** If Y is complete as a metric space, a certain estimate implies the vanishing of L^2-cohomology. More precisely, define the space of *ϵ-approximate harmonic forms*

$$\mathcal{H}_\epsilon(Y;\mathbb{E}) = \{\, \omega \in \operatorname{Dom} \overline{d}^* \cap A_c(Y;\mathbb{E}) \mid \|d\omega\|^2 + \|d^*\omega\|^2 \le \epsilon \|\omega\|^2 \,\} \ .$$

(Note that the intersection with $\operatorname{Dom} \overline{d}^*$, which forces ω to satisfy Neumann boundary conditions, may be omitted if $\partial Y = \emptyset$.)

Proposition. ([47, Prop. 1.2]) *Assume that Y is complete as a metric space. Then $\mathcal{H}^i_\epsilon(Y;\mathbb{E}) = 0$ for some $\epsilon > 0$ if and only if $H^i_{(2)}(Y;\mathbb{E}) = 0$ and $H^{i+1}_{(2)}(Y;\mathbb{E})$ is Hausdorff.*

11.3. **L^2-cohomology with boundary values.** For $\omega \in A(Y;\mathbb{E})$ define the norm

$$\|\omega\|_b^2 = \|\omega\|_Y^2 + \sum_F \|\omega|_F\|_F^2,$$

where F ranges over the closed proper boundary faces of Y. We will write, for example, "$\|\cdot\|_b$-bounded" if we mean to use this norm. Let $d_b = d_{Y,b} = d_{Y,b,\mathbb{E}}$ denote the unbounded operator on $L_2(Y;\mathbb{E})$ given by exterior differentiation on the domain

$$\operatorname{Dom} d_b = \{\, \omega \in A(Y;\mathbb{E}) \mid \|\omega\|_b, \|d\omega\|_b < \infty \,\}$$

and let $\overline{d}_b$ be its $\|\cdot\|_b$-closure in the graph norm. That is, $\omega \in L_2(Y;\mathbb{E})$ is in $\operatorname{Dom} \overline{d}_b$ if and only if there exists a sequence $\omega_i \in \operatorname{Dom} d_b$ such ω_i and $d\omega_i$ are Cauchy sequences in the norm $\|\cdot\|_b$ and $\|\omega_i - \omega\| \to 0$ (usual L^2 norm). Since $\|\cdot\| \le \|\cdot\|_b$, we have an inclusion of complexes $\operatorname{Dom} \overline{d}_b \subseteq \operatorname{Dom} \overline{d}$, but it is important to note that $\overline{d}_b$ is not a closed operator on $L_2(Y;\mathbb{E})$ in general. (In §11.7 below we will consider a larger Hilbert space in which $\overline{d}_b$ is closed.) The *L^2-cohomology with boundary values* is defined to be

$$H_{(2),b}(Y;\mathbb{E}) = H(\operatorname{Dom} \overline{d}_b) \ .$$

(Actually in general one would want to consider various weight functions on the boundary faces in defining $\|\cdot\|_b$, but this is not necessary here; see Remark 11.6 below.)

11.4. L^2-cohomology with boundary values is convenient since $\omega \in \operatorname{Dom} \overline{d}_b$ admits a strong L_2 trace or restriction $\omega|_F \in L_2(F;\mathbb{E}|_F)$ for each closed boundary face F. Specifically

Lemma. *Restriction to a closed bounded face F induces a well-defined $\|\cdot\|_b$-bounded map*

$$\operatorname{Dom} \overline{d}_{Y,b} \to \operatorname{Dom} \overline{d}_{F,b}$$

and consequently a map

$$H_{(2),b}(Y;\mathbb{E}) \longrightarrow H_{(2),b}(F;\mathbb{E}|_F) \ .$$

Proof. Given $\omega \in \operatorname{Dom} \overline{d}_b$ let ω_i be a sequence of smooth forms with $\|\omega_i - \omega\|_b \to 0$ and $d\omega_i$ a Cauchy sequence in the norm $\|\cdot\|_b$. Then $\omega_i|_F$ is a Cauchy sequence in $L_2(F;\mathbb{E}|_F)$ and we set $\omega|_F$ to be its limit. Clearly $\omega|_F \in \operatorname{Dom} \overline{d}_{F,b}$.

To see that $\omega|_F$ is well-defined, it suffices to consider $\omega = 0$ and F a codimension 1 boundary face. (For if $\omega|_F$ is well-defined and $F' \subset F$, then $\omega|_F \in \operatorname{Dom} \overline{d}_{F,b}$ and $\omega|_{F'} = (\omega|_F)|_{F'}$.) Let $\operatorname{int} F$ denote the interior of F and ν the outward unit normal vector. For any form $\alpha \in A_c(\operatorname{int} F;\mathbb{E}|_F)$, there clearly exists a form $\tilde{\alpha} \in A_c(Y;\mathbb{E})$

of one degree higher with Neumann boundary data $(\iota_\nu\tilde{\alpha})|_F = \alpha$. Then Stokes's theorem implies that

$$(\omega_i|_F, \alpha)_F = (d\omega_i, \widetilde{\alpha}) - (\omega_i, \delta\widetilde{\alpha})$$

where δ is the formal adjoint to d. Now if $\omega_i \to 0$ with $d\omega_i$ Cauchy, we would have $d\omega_i \to 0$ as usual and hence by the above formula, $(\omega_i|_F, \alpha)_F \to 0$ for all α. Since such α are dense in $L_2(F; \mathbb{E}|_F)$, we see that $\omega|_F = \lim \omega_i|_F = 0$. $\square$

11.5. On the other hand, we have an inclusion of complexes $\operatorname{Dom} \overline{d}_b \hookrightarrow \operatorname{Dom} \overline{d}$ which induces a homomorphism

$$H_{(2),b}(Y; \mathbb{E}) \longrightarrow H_{(2)}(Y; \mathbb{E}).$$

We will give a simple criterion for this to be an isomorphism which will suffice for our purposes.

Consider a system of tubular neighborhoods

$$(N_F, p_F, t_F)$$

of the closed boundary faces $F \subset Y$, where N_F is an open neighborhood of F, $p_F \colon N_F \to F$ is a smooth retraction, $t_F \colon N_F \to [0,1)^{\operatorname{codim} F}$ is a normal variable with $F = t_F^{-1}(0)$, and

$$(t_F, p_F) \colon N_F \xrightarrow{\sim} [0,1)^{\operatorname{codim} F} \times F$$

is a smooth diffeomorphism. We require that this data is compatible in the sense that if F and F' intersect, then

$$\begin{aligned} &N_{F\cap F'} = N_F \cap N_{F'}, \\ &p_{F\cap F'} = p_F p_{F'} = p_{F'} p_F \qquad \text{on } N_{F\cap F'}, \\ &t_{F\cap F'} = (t_F, t_{F'}) \qquad \text{on } N_{F\cap F'}, \text{ up to permutation of coordinates.} \end{aligned} \tag{11.5.1}$$

It is not difficult to see that for any manifold with corners (always assumed paracompact) such a system of tubular neighborhoods always exists. Note also that such data induces canonical isomorphisms

$$\mathbb{E}|_{N_F} \cong p_F^* \mathbb{E}|_F.$$

Lemma. *Let Y be a Riemannian manifold with corners and $\mathbb{E}$ a metrized locally constant sheaf on Y. Assume that a system of tubular neighborhoods satisfying* (11.5.1) *can be chosen so that the Riemannian metric ds^2 on Y and the metric h on $\mathbb{E}$ satisfy the quasi-isometry*[6] *relations*

$$ds^2|_{N_F} \sim dt_F^2 + p_F^* ds_F^2 \tag{11.5.2}$$

and

$$h|_{N_F} \sim p_F^*(h|_F)$$

uniformly for all codimension 1 *closed boundary faces F.* (*This holds in particular if Y is compact.*) *Then the natural map induces an isomorphism*

$$H_{(2),b}(Y; \mathbb{E}) \xrightarrow{\sim} H_{(2)}(Y; \mathbb{E}).$$

---

[6] Two metrics g, g' are said to be quasi-isometric if there exist constants C, $C' > 0$ such that $Cg' \leq g \leq Cg'$; we denote this by $g \sim g'$.

Proof. Let $\eta(t)$ be a smooth function on $[0,1]$ which is identically 1 on $[0,1/4]$ and has support in $[0,1/2)$. Assume first that Y has a smooth boundary F and set $t=t_F$ and $p=p_F$. As in [18], the homotopy operator

$$H\omega = 2\eta(t)\int_0^{1/2}\left(\int_a^t \iota_{\partial/\partial t}\omega\right) da$$

yields a homotopy formula

$$dH + Hd = I - P \tag{11.5.3}$$

on smooth forms, where

$$P\omega = (1-\eta(t))\omega + 2\eta(t)\, p^* \int_0^{1/2} (\omega|_{a\times F})\, da - 2\eta'(t)\, dt \wedge \int_0^{1/2}\left(\int_a^t \iota_{\partial/\partial t}\omega\right) da$$

Given our hypotheses on the metrics, an easy estimate using the Cauchy-Schwarz inequality demonstrates that H and P are bounded in the usual L^2 norm. Thus (11.5.3) holds on $\operatorname{Dom}\overline{d}$. On the other hand,

$$\|H\omega|_F\| = 2\|\int_0^{1/2}\left(\int_a^0 \iota_{\partial/\partial t}\omega\right) da\| \le C\|\omega\|$$

and

$$\|P\omega|_F\| = 2\|\int_0^{1/2}(\omega|_{a\times F})\, da\| \le C\|\omega\|.$$

Thus P defines a bounded map $\operatorname{Dom}\overline{d} \to \operatorname{Dom}\overline{d}_b$ and (11.5.3) holds on $\operatorname{Dom}\overline{d}_b$; this proves the lemma for the case of a single boundary face. In general one performs this construction for each codimension 1 closed boundary face and composes the homotopies and projections in the usual way [19, §15]; the hypotheses (11.5.1) ensure that the t_F-invariance produced by P_F is not disturbed by $P_{F'}$ or $H_{F'}$. $\square$

11.6. *Remark.* If Y is not compact then analogous results may be proven even if (11.5.2) fails. For example, if instead of (11.5.2) we have

$$ds^2|_{N_F} \sim (p_F^* w_F)^2 dt_F^2 + p_F^* ds_F^2 \ ,$$

where $w_F > 0$ is a bounded function on F, then the lemma continues to hold provided that $\|\cdot\|_b$ is defined using an L^2-norm for F which is weighted by w_F^2 (and a product weight for faces of higher codimension). Even weight functions not independent of t_F may be handled—for a nontrivial example and application see [42, §§2–4]. (In particular Lemma 3.2 of [42] is the analogue of the lemma above.)

11.7. Since $\overline{d}_b$ is not closed, we need to be careful about the meaning of the adjoint operator $\overline{d}_b^*$ before extending the spectral vanishing criterion of Proposition 11.2 to $H^i_{(2),b}(Y;\mathbb{E})$. Let $L_{2,b}(Y;\mathbb{E})$ denote the Hilbert space completion of $A_c(Y;\mathbb{E})$ with respect to $\|\cdot\|_b$; by associating to a form $\omega \in A_c(Y;\mathbb{E})$ the collection $(\omega|_F)_F$ of restrictions to faces, we obtain an isometry $L_{2,b}(Y;\mathbb{E}) \cong \bigoplus_F L_2(F;\mathbb{E}|_F)$. In this way $\operatorname{Dom}\overline{d}_b$ may be viewed as a subset of $L_{2,b}(Y;\mathbb{E})$ and the map $(\omega|_F)_F \mapsto (d\omega|_F)_F$ is a closed unbounded operator. Define $\overline{d}_b^*$ to be the adjoint of this operator in the sense of unbounded operators on $L_{2,b}(Y;\mathbb{E})$.

To understand $\overline{d}_b^*$ better, assume that $\psi = (\psi_F)_F \in \operatorname{Dom}\overline{d}_b^*$ with each ψ_F a smooth form on F. Then from the relation $(\overline{d}_b^*\psi, \omega)_b = (\psi, \overline{d}_b\omega)_b$ which is valid for all $\omega \in \operatorname{Dom}\overline{d}_b$, but in particular for all $\omega \in A_c(Y;\mathbb{E})$, it is easy to see that

$$(\overline{d}_b^*\psi)_F = \delta_F\psi_F + \sum_{\widetilde{F}\supset F}(\iota_{\nu_F^{\widetilde{F}}}\psi_{\widetilde{F}})|_F.$$

Here δ_F is the formal adjoint to d on F and the sum is over boundary faces $\widetilde{F}$ for which F is a codimension 1 face with unit normal vector $\nu_F^{\widetilde{F}}$.

Note that ψ belonging to $\operatorname{Dom}\overline{d}_b^*$ does not necessarily imply that the various ψ_F satisfy Neumann boundary conditions. Furthermore even if ψ is induced from a smooth form on Y, the same is not necessarily true for $\overline{d}_b^*\psi$.

11.8. **Spectral vanishing criterion for $H_{(2),b}(Y;\mathbb{E})$.** To state the vanishing criterion, define

$$\mathcal{H}_{\epsilon,b}(Y;\mathbb{E}) = \{\,\omega \in \operatorname{Dom}\overline{d}_b \cap \operatorname{Dom}\overline{d}_b^*,\ \operatorname{supp}\omega \text{ compact} \mid \\ \|\overline{d}_b\omega\|_b^2 + \|\overline{d}_b^*\omega\|_b^2 \le \epsilon\|\omega\|_b^2\,\}.$$

Note that from now on we will resume viewing $\operatorname{Dom}\overline{d}_b$ (and hence ω above) as being contained in $L_2(Y;\mathbb{E})$. Even so, $\overline{d}_b^*\omega$ must still be viewed as an element of $L_{2,b}(Y;\mathbb{E})$ as in the previous paragraph. We give $\operatorname{Ker}\overline{d}_b$ and $\operatorname{Range}\overline{d}_b$ the topology coming from the $\|\cdot\|_b$-norm and let $H_{(2),b}(Y;\mathbb{E})$ have the quotient topology.

Proposition. *Assume that Y is complete as a metric space. Then $\mathcal{H}^i_{\epsilon,b}(Y;\mathbb{E}) = 0$ for some $\epsilon > 0$ if and only if $H^i_{(2),b}(Y;\mathbb{E}) = 0$ and $H^{i+1}_{(2),b}(Y;\mathbb{E}) = 0$ is Hausdorff.*

Proof. The desired equivalence would follow from Banach's closed range theorem [26, §1.1] if it were not for the restriction that $\operatorname{supp}\omega$ be compact in the definition of $\mathcal{H}^i_{\epsilon,b}(Y;\mathbb{E})$. To see that this doesn't matter, first note that by Cohn-Vossen's generalization [3, 2.4] of the Hopf-Rinow theorem, the completeness of Y implies that the function $r(x)$ (distance in Y from a fixed point) has compact sublevel sets. Then (following [21] except we don't bother regularizing) use $r(x)$ to create a sequence of compactly supported Lipschitz functions η_k such that $0 \le \eta_k \le 1$, $|d\eta_k| \le C$, and every compact $K \subseteq Y$ is contained in $\eta_k^{-1}(1)$ for k sufficiently large. For $\omega \in \operatorname{Dom}\overline{d}_b \cap \operatorname{Dom}\overline{d}_b^*$, it follows that $\eta_k\omega \to \omega$, $\overline{d}_b(\eta_k\omega) \to \overline{d}_b\omega$, and $\overline{d}_b^*(\eta_k\omega) \to \overline{d}_b^*\omega$ in the $\|\cdot\|_b$ norm. Thus the existence of a nonzero $\omega \in \operatorname{Dom}\overline{d}_b \cap \operatorname{Dom}\overline{d}_b^*$ satisfying $\|\overline{d}_b\omega\|_b^2 + \|\overline{d}_b^*\omega\|_b^2 \le \epsilon\|\omega\|_b^2$ would imply that $\mathcal{H}^i_{2\epsilon,b}(Y;\mathbb{E}) \ne 0$. ☐

Remark. The proposition would remain true if in the definition of $\mathcal{H}_{\epsilon,b}(Y;\mathbb{E})$ we restrict to ω smooth (as for ordinary L^2-cohomology). The proof requires a generalization of Friedrich's regularization techniques to handle boundary values; since we will not need this stronger result, we will not go into details.

11.9. **L^2-cohomology with boundary values of $\widehat{X}$.** In the next section we will define a Riemannian metric on $\overline{X}$ (actually on a space $\overline{X}_t$ diffeomorphic to $\overline{X}$) whose restriction to $Y_P \subseteq \partial\overline{X}$ is N_P-invariant. Thus there is an induced Riemannian metric on every strata X_P of $\widehat{X}$. Even though $\widehat{X}$ is not a manifold with corners, we can define the operators $\overline{d}_{\widehat{X}}$ and $\overline{d}_{\widehat{X},b}$ as well as the L^2-cohomology groups $H_{(2)}(\widehat{X};\mathbb{E})$ and $H_{(2),b}(\widehat{X};\mathbb{E})$ just as for $\overline{X}$ but with the following differences:

- Forms in the domain of $\overline{d}_{\widehat{X}}$ and $\overline{d}_{\widehat{X},b}$ belong to the complex of special differential forms $A_{\mathrm{sp}}(\widehat{X};\mathbb{E})$ as opposed to $A(\overline{X};\mathbb{E})$.
- The $\|\cdot\|_b$-norm is defined using the restrictions $\omega \mapsto \omega|_{X_P}$ from (4.4.1) as opposed to $\omega \mapsto \omega|_{Y_P}$.

All the results in the preceding subsections extend to this context (and likewise if $\widehat{X}$ is replaced by any domain $U \subset \widehat{X}$).

12. L^2-COHOMOLOGY OF $\mathcal{L}$-MODULES

In this section we will represent the cohomology $H(\widehat{X};\mathcal{M})$ of an $\mathcal{L}$-module $\mathcal{M}$ by a variant of L^2-cohomology. To do this, we need to consider $\overline{X}_t \subset X$, a diffeomorphic image of $\overline{X}$ which was constructed in [43], and replace $\widehat{X}$ by the corresponding $\widehat{X}_t$. We state a vanishing criterion for this L^2-cohomology analogous to those in Propositions 11.2 and 11.8.

We also introduce a variant of $\mathcal{L}$-modules on $\bar{A}_P \times X_P$ and their L^2-cohomology.

12.1. **Locally symmetric metrics.** We will work with a fixed basepoint $x \in D$. Let $KA_G \subseteq G(\mathbb{R})$ be the stabilizer of the basepoint x and let θ be the Cartan involution associated to K (see for example [8, V.24.6] or [11, 1.6] for the nonconnected and nonsemisimple case) with Cartan decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{p} + \mathfrak{a}_G$. Fix an $\operatorname{Ad} G(\mathbb{R})$-invariant, θ-invariant nondegenerate bilinear form B on $\mathfrak{g}$ such that $(X,Y) \mapsto -B(X,\theta Y)$ is a positive definite inner product on $\mathfrak{g}$. (On the derived algebra one may take B to be the Killing form.) The restriction of B to $\mathfrak{p} + \mathfrak{a}_G$ yields an $\operatorname{Ad} KA_G$-invariant inner product on $T_x D$ which may be extended to a $G(\mathbb{R})$-invariant metric on D; this descends to a locally symmetric Riemannian metric on $X = \Gamma\backslash D$. In addition one obtains from B left-invariant metrics on all Lie subgroups of $G(\mathbb{R})$.

If (E,σ) is a regular representation of G let $\mathbb{E} = D \times_\Gamma E$ be the corresponding flat bundle on X. An *admissible* inner product on E is a positive definite Hermitian bilinear form on E which is K-invariant and for which $\mathfrak{p} + \mathfrak{a}_G$ acts via self-adjoint operators; equivalently it is a form for which $\sigma(g)\sigma(\theta g)^* = \mathrm{id}_E$ for $g \in G(\mathbb{R})$. Admissible metrics always exist (see [34, p. 375] in the case where $G(\mathbb{R})$ is connected and semisimple; the generalization to our situation is clear). Such an admissible inner product on E induces a metric on $\mathbb{E}$; in the case that E is isotypical it is given by the formula $|(gKA_G, v)| = |\xi_E^k(g)|^{\frac{1}{k}} \cdot |\sigma(g^{-1})v|$, where ξ_E is the character by which S_G acts on E and $k \in \mathbb{N}$ is such that ξ_E^k extends to a character on G.

The above construction may be applied to all L_P (using the basepoint $A_P \mathbf{o} N_P x \in D_P$, corresponding maximal compact subgroup $K_P \subseteq L_P(\mathbb{R})$ and the bilinear form on $\mathfrak{l}_P$ induced from B) in order to define locally symmetric metrics on the X_P and metrics on flat bundles $\mathbb{E}_P$ associated to regular representations E_P of L_P.

For a parabolic P there is (analogously to §1.4) a flat nilmanifold fibration $\Gamma_P\backslash D \to Z_P \equiv \Gamma_{L_P}\backslash(N_P(\mathbb{R})\backslash D)$. The space $N_P(\mathbb{R})\backslash D$ is naturally a homogeneous space for $L_P(\mathbb{R})$ with the stabilizer of a point being a maximal compact subgroup times A_G. The metric on D induces a $L_P(\mathbb{R})$-invariant metric on $N_P(\mathbb{R})\backslash D$ and hence a metric on Z_P. There is a trivial principal A_P^G-bundle $Z_P \to X_P$ given by geodesic retraction; canonical trivializing sections are given by quotients of $L_P(\mathbb{R})$-orbits. The metric on Z_P decomposes under a canonical trivialization into the sum of an invariant metric on A_P^G and the locally symmetric metric on X_P. Given a regular representations E_P of L_P, let $\widetilde{\mathbb{E}}_P = (N_P(\mathbb{R})\backslash D) \times_{\Gamma_{L_P}} E_P$ be the corresponding flat bundle on Z_P; alternatively $\widetilde{\mathbb{E}}_P$ is the pullback of $\mathbb{E}_P$ on X_P. The admissible inner product on E_P induces a metric on $\widetilde{\mathbb{E}}_P$ by the formula $|(rK_PA_P, v)| = |\sigma(r^{-1})v|$ for all $r \in M_P(\mathbb{R})A_P^G$; note that this is *not* the pullback of the metric on $\mathbb{E}_P$ unless A_P^G acts trivially.

12.2. **Diffeomorphisms of $\overline{X}$ into X.** By §1.3 we may identify D with $A_P^G \times e_P$ such that the ${}^0P(\mathbb{R})$-orbit of the fixed basepoint x is identified with $\{1\} \times e_P$. This induces identifications $\Gamma_P\backslash D \cong A_P^G \times Y_P$ and $Z_P \cong A_P^G \times X_P$.

By [43, Theorem 6.1][7] there exists a family (depending on $t \in A^G$ sufficiently dominant) of Γ-equivariant piecewise-analytic diffeomorphisms

$$s_t \colon \overline{D} \xrightarrow{\sim} \overline{D}_t \subseteq D \ ,$$

such that $\overline{X}_t = \Gamma\backslash\overline{D}_t \subseteq D$ is a compact subdomain with corners. For all[8] $P \in \mathcal{P}$ each diffeomorphism s_t satisfies:

$$e_P \xrightarrow{s_t|_{e_P}} e_{P,t} \equiv s_t(e_P) \text{ is } N_P(\mathbb{R})\text{-equivariant} \tag{12.2.1}$$

and

$$e_{P,t} \subseteq \{t_P\} \times e_P \tag{12.2.2}$$

(where $t_P \in A_P^G$ is the image of t under the canonical projection $A^G \cong A_P^G \times A^P \to A_P^G$). We obtain an induced diffeomorphism

$$\overline{X} \xrightarrow{\sim} \overline{X}_t = \Gamma\backslash\overline{D}_t \subseteq X \ .$$

12.3. We can describe $\overline{X}_t \subseteq X$ more precisely by using a cylindrical cover of X. Given $s_P \in A_P^G$ and $t \in A^G$, set

$$\begin{aligned} A_P^G(s_P) &= \{\, a \in A_P^G \mid a^\alpha > s_P^\alpha \text{ for all } \alpha \in \Delta_P \,\}, \text{ and} \\ \langle A_P^G\rangle_t &= \{\, a \in A_P^G \mid a^\beta \le t^\beta \text{ for all } \beta \in \widehat{\Delta}_P \,\}. \end{aligned}$$

For $\Omega_P \subseteq Y_P$ open, relatively compact, and $N_P(\mathbb{R})$-invariant, let $W_P = A_P^G(s_P) \times \Omega_P \subseteq A_P^G \times Y_P \cong \Gamma_P\backslash D$ be the associated *cylindrical set.* We always assume that s_P is sufficiently dominant (depending on Ω_P) so that W_P may be identified with its image in $X = \Gamma\backslash D$. For appropriate s_P and Ω_P we obtain a *cylindrical cover* $\{W_P\}_{P\in\mathcal{P}}$ of X. Given any cylindrical cover $\{W_P\}$, the intersections $\overline{X}_t \cap W_P$ form a cover of $\overline{X}_t$; for t sufficiently dominant one can prove [43, Theorem 5.7] that $\overline{X}_t \cap W_P \neq \emptyset$ and that

$$\overline{X}_t \cap W_P = (\langle A_P^G\rangle_t \times Y_P) \cap W_P \ . \tag{12.3.1}$$

---

[7] In fact [43] constructs a disjoint decomposition of D (indexed by parabolic $\mathbb{Q}$-subgroups) called a *tiling* of which $\overline{D}_t$ is the central tile corresponding to G; see in particular Definition 2.1, Theorem 5.7, and Theorem 6.1. Note the differences in notation: in [43], X refers to the symmetric space itself rather than its arithmetic quotient. More significantly, in [43] X_P denotes the piece of the tiling corresponding to P, as opposed to a stratum of $\widehat{X}$ as in the current paper.

[8] Actually equation (12.2.2) is only valid for P belonging to a fixed set of representatives of Γ-conjugacy classes of parabolic $\mathbb{Q}$-subgroups; we will assume such representatives have been chosen. For other P one has $e_{P,t} \subseteq \{t_Pb_P\} \times e_P$ where $b_P \in A_P^G$ is a certain parameter needed to make the construction Γ-invariant; this is because $\gamma \in \Gamma$ may move the fixed basepoint x. See [43, §2], particularly §§2.6, 2.7.

We set $\overline{W}_{P,t} \equiv \overline{X}_t \cap W_P$.

12.4. The diffeomorphism $\overline{X} \xrightarrow{\sim} \overline{X}_t$ induces diffeomorphisms $Y_P \xrightarrow{\sim} Y_{P,t} = \Gamma_P \backslash e_{P,t} \subseteq \Gamma_P \backslash D$ of the boundary faces. For each P this diffeomorphism descends by (12.2.1) to

$$X_P \xrightarrow{\sim} X_{P,t} \subseteq Z_P \ , \tag{12.4.1}$$

where $X_{P,t}$ is formed from $Y_{P,t}$ by collapsing the $N_P(\mathbb{R})'$-fibers of the nilmanifold fibration. We set

$$\widehat{X}_t = \coprod_P X_{P,t}$$

and equip it with the topology so that (12.4.1) induces a homeomorphism $\widehat{X} \xrightarrow{\sim} \widehat{X}_t$. The closure of $X_{P,t}$ will be denoted $\widehat{X}_{P,t}$.

The locally symmetric metric on $Z_P = A_P^G \times X_P$ induces on each stratum $X_{P,t}$ of $\widehat{X}_t$ a metric which extends smoothly to its Borel-Serre compactification $\overline{X}_{P,t}$. Given a regular representation E_P of L_P there is a natural isomorphism of flat bundles

$$\mathbb{E}_P \xrightarrow{\sim} \mathbb{E}_{P,t} \equiv \widetilde{\mathbb{E}}_P|_{X_{P,t}}$$

over the diffeomorphism (12.4.1). As in §12.1, the choice of an admissible inner product on E_P induces a metric on $\widetilde{\mathbb{E}}_P$, and hence a metric on $\mathbb{E}_{P,t}$ that also extends smoothly over $\overline{X}_{P,t}$.

12.5. Let $\mathcal{M} = (E_\cdot, f_{\cdot\cdot})$ be an $\mathfrak{L}$-module on $\widehat{X}$. In §4 we constructed an incarnation $\mathcal{A}_{\widehat{X}}(\mathcal{M})$ of the realization using special differential forms; this was a complex of fine sheaves on $\widehat{X}$. The cohomology $H(\widehat{X};\mathcal{M})$ is the cohomology of the global sections of $\mathcal{A}_{\widehat{X}}(\mathcal{M})$.

We now consider an alternate incarnation and realization which is a complex of sheaves on $\widehat{X}_t$ instead of $\widehat{X}$. Namely we set $\mathcal{A}_P(E_P) = \mathcal{A}_{\mathrm{sp}}(\widehat{X}_{P,t};\mathbb{E}_{P,t})$ instead of $\mathcal{A}_{\mathrm{sp}}(\widehat{X}_P;\mathbb{E}_P)$ and let $\mathcal{A}_{\widehat{X}_t}$ be the corresponding realization functor provided by (a generalization of) Theorem 4.1. If $U \subseteq \widehat{X}_t$ is a domain define $H(U;\mathcal{M})$ to be the cohomology of global sections of $\mathcal{A}_{\widehat{X}_t}(\mathcal{M})|_U$. For $U = \widehat{X}_t$, clearly

$$H(\widehat{X};\mathcal{M}) \cong H(\widehat{X}_t;\mathcal{M}) \ .$$

Explicitly for $U \subseteq \widehat{X}_t$ the cohomology $H(U;\mathcal{M})$ is the cohomology of the complex

$$\begin{cases} A_{\mathrm{sp}}(U;\mathcal{M}) = \bigoplus\limits_P A_{\mathrm{sp}}(U \cap \widehat{X}_{P,t};\mathbb{E}_{P,t}), \\ \qquad d = \sum\limits_P d_P + \sum\limits_{P \le Q} d_{PQ}, \end{cases} \tag{12.5.1}$$

where d_P is the differential on $A_{\mathrm{sp}}(U \cap \widehat{X}_{P,t};\mathbb{E}_{P,t})$ and d_{PQ} is defined by

$$d_{PQ}(\omega_Q) = A_{\mathrm{sp}}(U \cap \widehat{X}_{P,t}; f_{PQ} \circ h_{PQ})(\omega_Q|_{U\cap\widehat{X}_{P,t}}).$$

For $\omega = (\omega_Q)_Q \in A_{\mathrm{sp}}(U;\mathcal{M})$ define the weighted norm

$$\|\omega\|_t^2 = \sum_Q \|t^{-\rho_Q}\omega_Q\|_{Q,b}^2 \tag{12.5.2}$$

where the norms on the right-hand side are as in §§11.3 and 11.9,

$$\|\omega_Q\|^2_{Q,b} = \sum_{P \leq Q} \|\omega_Q|_{U\cap X_{P,t}}\|^2_{U\cap X_{P,t}} \ . \tag{12.5.3}$$

We let $t^{-\rho}$ denote the family of weights $(t^{-\rho_Q})_Q$ and let

$$L_2(U;\mathcal{M},t^{-\rho}) = \bigoplus_P L_{2,b}(U\cap \widehat{X}_{P,t};\mathbb{E}_{P,t},t^{-\rho_Q})$$

denote the Hilbert space completions of $A_{\mathrm{sp}}(U;\mathcal{M})$ and $A_{\mathrm{sp}}(U\cap\widehat{X}_{P,t};\mathbb{E}_{P,t})$ under the norms (12.5.2) and (12.5.3).

For each P the closed unbounded operator $\overline{d}_{U\cap\widehat{X}_{P,t},b,t^{-\rho_P}}$ from §§11.3 and 11.9 is defined on $L_{2,b}(U\cap\widehat{X}_{P,t};\mathbb{E}_{P,t},t^{-\rho_Q})$ and has cohomology $H_{(2),b}(U\cap\widehat{X}_{P,t};\mathbb{E}_{P,t},t^{-\rho_Q})$. On the other hand, let $d_{U;\mathcal{M},t^{-\rho}}$ be the unbounded operator on $L_2(U;\mathcal{M},t^{-\rho})$ corresponding to the subcomplex of (12.5.1) consisting of ω such that ω, $d\omega \in L_2(U;\mathcal{M},t^{-\rho})$ and denote its graph norm closure by $\overline{d} = \overline{d}_{\mathcal{M}} = \overline{d}_{U;\mathcal{M},t^{-\rho}}$. The cohomology of $\operatorname{Dom}\overline{d}_{U;\mathcal{M},t^{-\rho}}$ is by definition the *L^2-cohomology of $\mathcal{M}$ over U with weights $t^{-\rho}$* and is denoted $H_{(2)}(U;\mathcal{M},t^{-\rho})$. In order to relate these L^2-cohomology groups we need the

12.6**. Lemma.** *The operators d_{PQ} are bounded with respect to* (12.5.2) *independently of t.*

Proof. There are bounds independent of t for both the restriction map and f_{PQ}: the first since $\|\cdot\|_{Q,b}$ incorporates the L^2-norm of boundary values, and the second since it is induced by a map of L_P-modules with admissible inner product. As for h_{PQ}, recall from the proof of Lemma 4.7 that this acts on the coefficients via the composition

$$\mathbb{A}_{\mathrm{inv}}(\mathcal{N}^Q_P(\mathbb{R})';\mathbb{E}_{Q,t}) \xrightarrow{h_x} \mathbb{C}(\mathfrak{n}^Q_P;E_Q)_x \longrightarrow \mathbb{H}(\mathfrak{n}^Q_P;E_Q) \ .$$

The second map here is induced from a map of L_P-modules so it remains to consider h_x at a point $z \in U\cap X_{P,t}$. Let $Y^Q_{P,t}$ denote the Borel-Serre boundary stratum of $\overline{X}_{Q,t}$ associated to P and let $p\colon Y^Q_{P,t}\to X_{P,t}$ denote the nilmanifold fibration with typical fiber $\mathcal{N}^Q_P(\mathbb{R})'$. The map h_x is induced by the Γ_{L_P}-morphism which transfers an invariant form on $\mathcal{N}^Q_P(\mathbb{R})'$ to one on $N^Q_P(\mathbb{R})'$ via n'_x (see §1.4) and then takes its value at the identity. If $\psi \in \mathbb{A}_{\mathrm{inv}}(\mathcal{N}^Q_P(\mathbb{R})';\mathbb{E}_{Q,t})_z$ is an invariant form on $p^{-1}(z)$, then the pointwise norm $|\psi_y|$ is independent of $y\in p^{-1}(z)$ and thus

$$|\psi|^2 = \int_{p^{-1}(z)} |\psi_y|^2 dy = \operatorname{Vol}(p^{-1}(z))|\psi_{y_0}|^2 = \operatorname{Vol}(p^{-1}(z))|h_x(\psi)|^2.$$

This implies $|h_x|^2 = \operatorname{Vol}(p^{-1}(z))^{-1} = t^{2\rho^Q_P}\operatorname{Vol}(N^Q_P(\mathbb{R})')^{-1}$ and hence $\|h_{PQ}\|^2_t$ is bounded independently of t. □

12.7**. Corollary.** *There is a decomposition* (*as vector spaces, not as complexes*)

$$\operatorname{Dom}\overline{d}_{U;\mathcal{M},t^{-\rho}} = \bigoplus_P \operatorname{Dom}\overline{d}_{U\cap\widehat{X}_{P,t},b,t^{-\rho_P}}$$

and a strict operator equality

$$\overline{d}_{U;\mathcal{M},t^{-\rho}} = \sum_P \overline{d}_{U\cap\widehat{X}_{P,t},b,t^{-\rho_P}} + \sum_{P\leq Q} d_{PQ} \ .$$

For $U = \widehat{X}$ we have the

12.8. **Corollary.** $H(\widehat{X}_t;\mathcal{M}) \cong H_{(2)}(\widehat{X}_t;\mathcal{M},t^{-\rho})$.

Proof. Filter both (12.5.1) and $\operatorname{Dom}\overline{d}_{U;\mathcal{M},t^{-\rho}}$ by $\#\Delta_P \le p$ and compare the resulting spectral sequences; they are isomorphic at E_1 by (11.1.1) and Lemma 11.5. □

12.9. Define

$$\mathcal{H}^i_\epsilon(U;\mathcal{M},t^{-\rho}) = \{\, \omega \in \operatorname{Dom}\overline{d}_{U;\mathcal{M},t^{-\rho}} \cap \operatorname{Dom}\overline{d}^{*}_{U;\mathcal{M},t^{-\rho}},\ \operatorname{supp}\omega \text{ compact} \mid \\ \|\overline{d}\omega\|_t^2 + \|\overline{d}^{*}\omega\|_t^2 \le \epsilon\|\omega\|_t^2 \,\}\ .$$

For $U = \widehat{X}$ we have as in Proposition 11.8

Proposition. *$\mathcal{H}^i_\epsilon(\widehat{X};\mathcal{M},t^{-\rho}) = 0$ for some $\epsilon > 0$ if and only if $H^i_{(2)}(\widehat{X};\mathcal{M},t^{-\rho}) = 0$ and $H^{i+1}_{(2)}(\widehat{X};\mathcal{M},t^{-\rho})$ is Hausdorff.*

Actually $H_{(2)}(\widehat{X};\mathcal{M},t^{-\rho})$ is finite-dimensional so the Hausdorff condition above is vacuous. However the proposition also applies in an different context which we will now introduce and where finite-dimensionality is not assured.

12.10. **$\mathcal{L}$-modules on $\langle A_P^G\rangle_t \times X_P$.** The space $\langle A_P^G\rangle_t \times X_P \subseteq Z_P$ is a manifold with corners with strata indexed by the parabolic $\mathbb{Q}$-subgroups $Q \ge P$. Explicitly the Q-stratum is $\langle A_P^G\rangle_t^Q \times X_P$ where

$$\langle A_P^G\rangle_t^Q = \{\, a \in A_P^G \mid a^\beta \le t^\beta \text{ for all } \beta \in \widehat{\Delta}_P \setminus \widehat{\Delta}_Q,\ a^\beta = t^\beta \text{ for all } \beta \in \widehat{\Delta}_Q \,\}\ .$$

We may extend the theory of $\mathcal{L}$-modules and their realizations to such spaces. In fact the theory becomes much simpler—an $\mathcal{L}$-module $\mathcal{M}' = (E'_\cdot, f'_{\cdot\cdot})$ on $\langle A_P^G\rangle_t \times X_P$ consists of a family $(E'_Q)_{Q\ge P}$ of graded regular representations of the *same* group L_P and degree 1 morphisms $f'_{QR}\colon E'_R \xrightarrow{[1]} E'_Q$. (Since the space has homotopically trivial links the functor $H(\mathfrak{n}_R^Q;\cdot)$ is replaced throughout by the identity functor.) Furthermore special differential forms are no longer needed for the realization.

The strata and the locally constant sheaves associated to E'_Q are metrized as in §12.4 by restricting the locally symmetric metrics on Z_P and $\widetilde{\mathbb{E}}'_Q$. We will be considering the L^2-space $L_2(\langle A_P^G\rangle_t \times X_P;\mathcal{M}')$ (without weights) and the corresponding L^2-cohomology. Since $\langle A_P^G\rangle_t \times X_P$ is a complete metric space the analogue of Proposition 12.9 holds in this context.

13. Analytic Lemmas

The goal of this section is Proposition 13.5, a vanishing criterion for $H^i(\widehat{X};\mathcal{M})$ in terms of certain L^2-cohomology groups of $\langle A_P^G\rangle_1 \times X_P$ for all $P \in \mathcal{P}$. In view of Proposition 12.9 this is accomplished by presenting a sequence of lemmas that reduce the vanishing of $\mathcal{H}^i_\epsilon(\widehat{X};\mathcal{M},t^{-\rho})$. As we proceed, the parameter t will have to be made successively more dominant.

13.1. Given a cylindrical cover $\{W_P\}$ assume t is sufficiently dominant so that (12.3.1) holds. Each boundary stratum $Y_{Q,t} \cap \overline{W}_{P,t}$ of $\overline{W}_{P,t}$ for $Q \ge P$ is part of a nilmanifold fibration with fibers $N_Q(\mathbb{R})'$; let $\widehat{W}_{P,t} \subseteq \widehat{X}_t$ denote the result of collapsing these fibers. We have the following spectral analogue of the Mayer-Vietoris sequence:

Lemma. *Given* $\epsilon > 0$ *there exists* $\epsilon' > 0$ *and a cylindrical cover* $\{W_P\}$ *of* X *such that for all* t *sufficiently dominant,* $\mathcal{H}^i_\epsilon(\widehat{W}_{P,t}; \mathcal{M}, t^{-\rho}) = 0$ *for all* P *implies* $\mathcal{H}^i_{\epsilon'}(\widehat{X}_t; \mathcal{M}, t^{-\rho}) = 0$.

Proof. For every $c > 0$ (to be chosen below) there exists [47, Prop. 2.1] a cylindrical cover $\{W_P\}$ possessing a partition of unity $\{\eta_P\}$ such that $|d\eta_P| < c$. (It is easy to eliminate the neatness hypothesis.) Examination of the construction shows that for $P \leq Q$ the function η_P is $N_Q(\mathbb{R})$-invariant on $W_P \setminus \bigcup_{P' \not\leq Q} W_{P'}$. But since $W_P \cap Y_{Q,t}$ is empty unless for $P \leq Q$ by (12.2.2) and (12.3.1), this implies that $\eta_P|_{Y_{Q,t}}$ is $N_Q(\mathbb{R})$-invariant for *any* P and Q. Thus we may pushdown to obtain a partition of unity $\{\hat\eta_P\}$ for the covering $\{\widehat{W}_{P,t}\}$ of $\widehat{X}_t$. Now for $\omega \in \mathcal{H}^i_{\epsilon'}(\widehat{X}_t; \mathcal{M}, t^{-\rho})$ (with $\epsilon' > 0$ to be chosen below) one may estimate (compare [47, Prop. 1.4]):

$$\begin{aligned}
\|\overline{d}(\hat\eta_P\omega)\|_t^2 + \|\overline{d}^*(\hat\eta_P\omega)\|_t^2 &\leq 2\big(\|\hat\eta_P\overline{d}\omega\|_t^2 + \|\hat\eta_P\overline{d}^*\omega\|_t^2 + \|\overline{d}\hat\eta_P \wedge \omega\|_t^2 + \|\overline{d}\hat\eta_P \lrcorner \omega\|_t^2\big) \\
&\leq 2\big(\|\overline{d}\omega\|_t^2 + \|\overline{d}^*\omega\|_t^2 + 2c^2\|\omega\|_t^2\big) \\
&< 2(\epsilon' + 2c^2)\|\omega\|_t^2 \\
&\leq 2(\#\mathcal{P})^2(\epsilon' + 2c^2)\max_R \|\hat\eta_R\omega\|_t^2 \leq \epsilon \max_R \|\hat\eta_R\omega\|_t^2,
\end{aligned}$$

where for the last line we choose $c > 0$ and $\epsilon' > 0$ small. Thus if P realizes the maximum on the last line we have $\hat\eta_P\omega \in \mathcal{H}^i_\epsilon(\widehat{W}_{P,t}; \mathcal{M}, t^{-\rho})$. Since $\omega \neq 0$ implies $\hat\eta_P\omega \neq 0$ for this P, the lemma is proved. □

13.2. Now assume that W_P is a cylindrical set for a fixed P and that t is sufficiently dominant so that (12.3.1) holds. Let O_P be the image of Ω_P under the nilmanifold fibration $Y_P \to X_P$. Each boundary stratum $X_{Q,t} \cap \widehat{W}_{P,t}$ of $\widehat{W}_{P,t}$ for $Q \geq P$ is part of a nilmanifold fibration with fibers $N_P^Q(\mathbb{R})'$; let

$$\widehat{W}_{P,t} \xrightarrow{p} (\langle A_P^G\rangle_t \cap A_P^G(s_P)) \times O_P \subseteq \langle A_P^G\rangle_t \times X_P$$

be the projection obtained by collapsing these fibers.

Define a pushforward $\mathcal{L}$-module $p_*\mathcal{M} = (E'_\cdot, f'_{\cdot\cdot})$ on $\langle A_P^G\rangle_t \times X_P \subseteq Z_P$ in the sense of §12.10 by $E'_Q = H(\mathfrak{n}_P^Q; E_Q)$ and $f'_{QR} = H(\mathfrak{n}_P^Q; f_{QR})$ and set $p_*\mathcal{M} \otimes \mathbb{C}_{\rho_P} = (E'_\cdot \otimes \mathbb{C}_{\rho_P}, f'_{\cdot\cdot} \otimes \mathrm{id}_{\mathbb{C}_{\rho_P}})$.

Lemma. *Given* $\epsilon > 0$ *and a cylindrical set* W_P *there exists* $\epsilon' > 0$ *and an injective map*

$$\mathcal{H}^i_{\epsilon'}(\widehat{W}_{P,t}; \mathcal{M}, t^{-\rho}) \longrightarrow \mathcal{H}^i_\epsilon(\langle A_P^G\rangle_t \times X_P; p_*\mathcal{M} \otimes \mathbb{C}_{\rho_P})$$

for all t *sufficiently dominant.*

Proof. For $Q \geq P$ let p_{Q*} be the operator on $L_2(X_{Q,t} \cap \widehat{W}_{P,t}; \mathbb{E}_{Q,t}, t^{-\rho_Q})$ which orthogonally projects onto $N_P^Q(\mathbb{R})$-invariant and $\mathfrak{n}_P^Q$-harmonic forms. Zucker [56, (4.24)] shows there is a homotopy formula

$$\overline{d}_Q H_Q + H_Q \overline{d}_Q = I - p_{Q*} \tag{13.2.1}$$

with H_Q and p_{Q*} bounded in the L^2 norm; both $\|H_Q\|_Q$ and $\|p_{Q*}\|_Q$ are independent of t (though dependent on W_P). In fact since the boundary faces of $\overline{X}_{Q,t} \cap \widehat{W}_{P,t}$ contain full $N_P^Q(\mathbb{R})$-fibers, the result holds for the $\|\cdot\|_{Q,b}$ norm and we obtain a homotopy formula for $\overline{d}_{Q,b,t^{-\rho_Q}}$.

We need the compatibility conditions

$$p_{Q*}d_{QR} = d_{QR}p_{R*} \qquad \text{and} \qquad H_Q d_{QR} = -d_{QR}H_R \tag{13.2.2}$$

(the sign is due to the fact that d_{QR} is degree 1 in the coefficients). To see that this can be arranged, note that Zucker's construction for H_Q depends on the choice of a central series defined over $\mathbb{Q}$,

$$1 = N_0^Q \subset N_1^Q \subset \cdots \subset N_\nu^Q = N_P^Q(\mathbb{R}),$$

and unit $\mathbb{Q}$-root vectors $Z_j^Q \in \mathfrak{n}_j^Q$ which satisfy $\mathfrak{n}_j^Q = \mathfrak{n}_{j-1}^Q + \mathbb{R}\cdot Z_j^Q$. Choose such a central series and root vectors for $N_P(\mathbb{R})$ (the case $Q = G$) and use the induced data for each $N_P^Q(\mathbb{R})$. (Some of the vectors Z_j^Q will now be zero but those stages in the series may be omitted.) One may now verify from the description in [56, §4(c)] that (13.2.2) holds.

By summing (13.2.1) over $Q \geq P$ we obtain

$$\overline{d}_{\mathcal{M}}H + H\overline{d}_{\mathcal{M}} = I - p_*, \tag{13.2.3}$$

where $H = \sum_{Q\geq P} H_Q$, $p_* = \sum_{Q\geq P} p_{Q*}$, and the d_{QR} terms vanish due to (13.2.2). Now for $\omega \in \mathcal{H}^i_{\epsilon'}(\widehat{W}_{P,t}; \mathcal{M}, t^{-\rho})$ calculate:

$$\begin{aligned}\|\overline{d}\omega\|_t^2 + \|\overline{d}^*\omega\|_t^2 &\leq \epsilon'\|\omega\|_t^2 \\ &\leq \epsilon'(\|p_*\omega\|_t^2 + \|(I-p_*)\omega\|_t^2) \\ &\leq \epsilon'\|p_*\omega\|_t^2 + 2\epsilon'\|H\|_t^2\cdot(\|\overline{d}\omega\|_t^2 + \|\overline{d}^*\omega\|_t^2);\end{aligned}$$

the final line uses (13.2.3)—see [47, Prop. 1.6]. This estimate shows that p_* is injective on $\mathcal{H}^i_{\epsilon'}$ provided we choose $\epsilon' > 0$ so that (say) $2\epsilon'\|H\|_t^2 \leq 1/2$. Furthermore in this case the last term may be absorbed into the left-hand side and we obtain

$$\|\overline{d}p_*\omega\|_t^2 + \|\overline{d}^*p_*\omega\|_t^2 \leq \|p_*\|_t^2\cdot(\|\overline{d}\omega\|_t^2 + \|\overline{d}^*\omega\|_t^2) \leq 2\epsilon'\|p_*\|_t^2\cdot\|p_*\omega\|_t^2.$$

This shows that p_* takes $\mathcal{H}^i_{\epsilon'}$ into $\mathcal{H}^i_\epsilon$ provided we choose $\epsilon' > 0$ so that $2\epsilon'\|p_*\|_t^2 \leq \epsilon$. Since $\|H\|_t$ and $\|p_*\|_t$ are bounded uniformly in t, $\epsilon' > 0$ may be chosen independently of t.

To complete the proof, we embed $p_*(\operatorname{Dom}\overline{d}_{\mathcal{M},t^{-\rho}})$ into $\operatorname{Dom}\overline{d}_{p_*\mathcal{M}\otimes\mathbb{C}_{\rho_P}}$ by extending an invariant compactly supported form by zero and applying Lemma 4.7 to the coefficients; we also need to choose a flat trivializing section of the bundle associated to $\mathbb{C}_{\rho_P}$. To see that this is an isometry, note that the L^2 norm integrand for the Q summand of $p_*(\operatorname{Dom}\overline{d}_{\mathcal{M},t^{-\rho}})$ involves $t^{-2\rho_Q}$ whereas for $\operatorname{Dom}\overline{d}_{p_*\mathcal{M}\otimes\mathbb{C}_{\rho_P}}$ the coefficient system $\mathbb{C}_{\rho_P}$ contributes $a^{-2\rho_P} = a^{-2\rho_Q}\cdot a^{-2\rho_P^Q} = t^{-2\rho_Q}\cdot a^{-2\rho_P^Q}$. The discrepancy is canceled by the map $\mathbb{A}_{\mathrm{inv}}(\mathcal{N}_P^Q(\mathbb{R})'; \mathbb{E}_{Q,t}) \xrightarrow{h_x} \mathbb{C}(\mathfrak{n}_P^Q; E_Q)_x$ which multiplies norm squared by $a^{2\rho_P^Q}$ (see the proof of Lemma 12.6). ☐

13.3. The isotypical decomposition of a regular L_P-module induces a decomposition of $\mathcal{L}$-modules

$$p_*\mathcal{M} = \bigoplus_V (p_*\mathcal{M})_V$$

where V ranges over irreducible regular L_P-modules.

Lemma. *For any $\epsilon > 0$ and any t, the vanishing $\mathcal{H}^i_\epsilon(\langle A_P^G\rangle_t \times X_P; p_*\mathcal{M}\otimes\mathbb{C}_{\rho_P}) = 0$ is equivalent to the vanishing conditions*

$$\mathcal{H}^i_\epsilon(\langle A_P^G\rangle_t \times X_P; (p_*\mathcal{M})_V\otimes\mathbb{C}_{\rho_P}) = 0, \qquad \textit{for all } V.$$

Proof. If $\omega = \sum_V \omega_V \in \mathcal{H}^i_\epsilon(\langle A^G_P\rangle_t \times X_P; p_*\mathcal{M} \otimes \mathbb{C}_{\rho_P})$ is nonzero, then for at least one V, $\omega_V \neq 0$ and $\omega_V \in \mathcal{H}^i_\epsilon(\langle A^G_P\rangle_t \times X_P; (p_*\mathcal{M})_V \otimes \mathbb{C}_{\rho_P})$. For otherwise we could sum the estimates $\|\overline{d}\omega_V\|^2 + \|\overline{d}^*\omega_V\|^2 > \epsilon\|\omega_V\|^2$ and obtain a contradiction. The converse is obvious. □

13.4. **Lemma.** *The vanishing condition $\mathcal{H}^i_\epsilon(\langle A^G_P\rangle_t \times X_P; (p_*\mathcal{M})_V \otimes \mathbb{C}_{\rho_P}) = 0$ is independent of t.*

Proof. Let $\Psi_t \colon \langle A^G_P\rangle_1 \times X_P \xrightarrow{\sim} \langle A^G_P\rangle_t \times X_P$ be the diffeomorphism given by the left action of t_P. Although Ψ_t is an isometry, the natural isomorphism of coefficient systems $(\mathbb{E}'_{Q,1})_V \otimes \mathbb{C}_{\rho_P} \xrightarrow{\sim} \Psi^*_t((\mathbb{E}'_{Q,t})_V \otimes \mathbb{C}_{\rho_P})$ multiplies norm by $t^{-(\xi_V+\rho_P)}$, where ξ_V is the character by which S_P acts on V. Thus the norm of a form depends on t only via a scaling factor independent of Q. This implies that $\overline{d}^*\Psi^*_t = \Psi^*_t\overline{d}^*$ and that the scaling factor may be canceled from all terms of the estimate $\|\overline{d}\omega\|^2 + \|\overline{d}^*\omega\|^2 \leq \epsilon\|\omega\|^2$. □

Remark. This lemma would not hold if we had not included the weight factors in (12.5.2).

13.5. We finally arrive at the following vanishing criterion for the cohomology of an $\mathcal{L}$-module.

Proposition. *Let $\mathcal{M}$ be an $\mathcal{L}$-module on $\widehat{X}$. Assume*

$$H^i_{(2)}(\langle A^G_P\rangle_1 \times X_P; (p_*\mathcal{M})_V \otimes \mathbb{C}_{\rho_P}) = 0 \text{ , and} \qquad (13.5.1)$$
$$H^{i+1}_{(2)}(\langle A^G_P\rangle_1 \times X_P; (p_*\mathcal{M})_V \otimes \mathbb{C}_{\rho_P}) \text{ is Hausdorff.}$$

for all parabolic $\mathbb{Q}$-subgroups P and $V \in \mathfrak{Irr}(L_P)$. Then $H^i(\widehat{X};\mathcal{M}) = 0$.

Proof. By Corollary 12.8 and Proposition 12.9 it suffices to show $\mathcal{H}^i_\epsilon(\widehat{X};\mathcal{M},t^{-\rho}) = 0$ for some $\epsilon > 0$. For this vanishing it suffices by Lemmas 13.1–13.4 to show there exists $\epsilon > 0$ such that

$$\mathcal{H}^i_\epsilon(\langle A^G_P\rangle_1 \times X_P; (p_*\mathcal{M})_V \otimes \mathbb{C}_{\rho_P}) = 0$$

for all P and V. However by Proposition 12.9 again (see §12.10) this is implied by (13.5.1). □

14. Vanishing Theorem for L^2-cohomology

In the course of verifying (13.5.1) we will need to apply the following vanishing theorem to the ordinary global L^2-cohomology $H_{(2)}(X_P;\mathbb{V})$. The heart of the theorem is a calculation originally due to Raghunathan [37], [38]. The theorem was proven in [47] but since it was not stated explicitly in this form we sketch the proof in this section.

14.1. **Theorem.** *Let $X = \Gamma\backslash G(\mathbb{R})/KA_G$ be the locally symmetric space associated to an arithmetic subgroup Γ of a reductive algebraic group G defined over $\mathbb{Q}$. Let E be an irreducible regular representation of G and let $\mathbb{E}$ be the associated locally constant sheaf on X. Endow X with a locally symmetric Riemannian metric and let $\mathbb{E}$ have a metric induced from an admissible inner product on E.*

(i) *If $(E|_{{}^0G})^* \ncong \overline{E|_{{}^0G}}$, then $H_{(2)}(X;\mathbb{E}) = 0$.*

(ii) *If* $(E|_{{}^0G})^* \cong \overline{E|_{{}^0G}}$, *then* $H^i_{(2)}(X;\mathbb{E}) = 0$ *for*

$$i \notin [(\dim D - \dim D(E))/2, (\dim D + \dim D(E))/2] ,$$

where $\dim D(E)$ *is defined in* §10.2. *Furthermore,* $H^i_{(2)}(X;\mathbb{E})$ *is Hausdorff (and hence representable by harmonic forms) for* $i = (\dim D - \dim D(E))/2$.

Sketch of Proof (compare [47, §§8–10]). We will determine a sufficient condition on a degree i so that the estimate

$$\|d\omega\|^2 + \|d^*\omega\|^2 > \epsilon\|\omega\|^2 \tag{14.1.1}$$

holds for all $\omega \in A^i_c(X;\mathbb{E})$ and some $\epsilon > 0$. Since X is complete and without boundary, Proposition 11.2 shows that this estimate for a given i implies $H^i_{(2)}(X;\mathbb{E}) = 0$ and $H^{i+1}_{(2)}(X;\mathbb{E})$ is Hausdorff.

Let $K \subseteq G(\mathbb{R})$ be a maximal compact subgroup with associated Cartan involution θ and Cartan decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{p} + \mathfrak{a}_G$. Let σ_1 denote the coadjoint action of $\mathfrak{k}_{\mathbb{C}}$ on $\bigwedge \mathfrak{p}^*_{\mathbb{C}}$ and let σ_2 denote the action of $\mathfrak{g}_{\mathbb{C}}$ on E. Since $X \cong \Gamma\backslash({}^0G(\mathbb{R}))^0/K^0$, the complex $A_c(X;\mathbb{E})$ may as usual be isometrically identified with the complex [13, I, §§1.2, 5.1, VII, §2.7], [34, §4]

$$\begin{cases} C({}^0\mathfrak{g}, \mathfrak{k}; E \otimes A^0_c(\Gamma\backslash({}^0G(\mathbb{R}))^0)) \cong \operatorname{Hom}_{\mathfrak{k}}(\bigwedge \mathfrak{p}, E \otimes A^0_c(\Gamma\backslash({}^0G(\mathbb{R}))^0)) , \\ \qquad d = \sum_k \epsilon(X_k)(\sigma_2(X_k) + X_k) , \end{cases}$$

where $\{X_k\}$ denotes an orthonormal basis of $\mathfrak{p}$, acting on $A^0_c(\Gamma\backslash({}^0G(\mathbb{R}))^0)$ via differentiation by the corresponding left-invariant vector field, and $\epsilon(X_k)$ denotes exterior multiplication by the corresponding element of the dual basis. Note that the operator d extends to the larger space $\bigwedge \mathfrak{p}^*_{\mathbb{C}} \otimes E \otimes A^0_c(\Gamma\backslash({}^0G(\mathbb{R}))^0)$ (without the $\mathfrak{k}$-invariance condition) though it is no longer a differential here.

Integration by parts and some multi-linear algebra allow one to extract 0-order terms and estimate that

$$\|d\omega\|^2 + \|d^*\omega\|^2 \geq (\Delta_0\omega, \omega) \geq 0 \tag{14.1.2}$$

[47, Props. 9.2 and 9.4] for a certain operator Δ_0. Explicitly

$$\Delta_0 = \sigma_2(C) - \frac{1}{2}\sigma_2(C_{\mathfrak{k}}) + \frac{1}{2}\sigma_1(C_{\mathfrak{k}}) - \frac{1}{2}(\sigma_1 \otimes \sigma_2)(C_{\mathfrak{k}}) , \tag{14.1.3}$$

where C (resp. $C_{\mathfrak{k}}$) denotes the Casimir operator of ${}^0\mathfrak{g}$ (resp. $\mathfrak{k}$). In fact Δ_0 is induced from the algebraic Laplacian on $C({}^0\mathfrak{g}, \mathfrak{k}; E) = \operatorname{Hom}_{\mathfrak{k}}(\bigwedge \mathfrak{p}, E) \subseteq \bigwedge \mathfrak{p}^*_{\mathbb{C}} \otimes E$.

We now need some more notation. Let $\mathfrak{h} = \mathfrak{b}_G + \mathfrak{a}_G = \mathfrak{b}_{G,\mathfrak{k}} + \mathfrak{b}_{G,\mathfrak{p}} + \mathfrak{a}_G$ be a fundamental θ-stable Cartan subalgebra of $\mathfrak{g}$. Fix a positive system $\Phi^+_{\mathfrak{k}} = \Phi^+(\mathfrak{k}_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}})$ of roots for $\mathfrak{k}_{\mathbb{C}}$ and consider any θ-stable positive system $\Phi^+ = \Phi^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ compatible with $\Phi^+_{\mathfrak{k}}$. (A specific choice will be made below.) Let $\Phi^+(\mathfrak{p}_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}})$ denote the positive nonzero weights of $\mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}$ in $\mathfrak{p}_{\mathbb{C}}$, counted with multiplicity. (A weight is "positive" here if it is the restriction to $\mathfrak{b}_{G,\mathfrak{k}}$ of a positive root [47, 10.1].) Clearly

$$\dim D = 2 \cdot \#\Phi^+(\mathfrak{p}_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) + \dim \mathfrak{b}_{G,\mathfrak{p}} . \tag{14.1.4}$$

Let λ be the highest weight of E and recall the reductive Lie subgroup $G(\lambda) \subseteq G$ defined in §10.1 with positive roots $\Phi^+(\mathfrak{g}(\lambda)_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) = \{\, \gamma \in \Phi^+ \mid \gamma \perp \lambda \,\}$; define $\Phi^+(\mathfrak{p}(\lambda)_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) = \{\, \alpha \in \Phi^+(\mathfrak{p}_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) \mid \alpha \perp \lambda \,\}$ similarly. Note that if $\lambda|_{\mathfrak{b}_{G,\mathfrak{p}}} = 0$ (so that $G(\lambda)$ is defined over $\mathbb{R}$ as in §10.2) then $\Phi^+(\mathfrak{p}(\lambda)_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}})$ is indeed the set of positive $\mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}$-weights of the split part in the Cartan decomposition of $\mathfrak{g}(\lambda)$. In this case

$$\dim D(\lambda) = 2 \cdot \#\Phi^+(\mathfrak{p}(\lambda)_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) + \dim \mathfrak{b}_{G,\mathfrak{p}} \ . \tag{14.1.5}$$

Equation (14.1.3) shows that for any fixed irreducible submodule R of $\sigma_1 \otimes \sigma_2|_{\mathfrak{k}}$, Δ_0 acts as a certain scalar on all forms in $R \otimes A_c^0(\Gamma\backslash({}^0G(\mathbb{R}))^0))$. We need to determine when this scalar can be zero. It suffices to assume $R \subseteq S_1 \otimes S_2$, where S_1 is an irreducible submodule of σ_1 with pure degree i and highest weight ν_1, and S_2 is an irreducible submodule of $\sigma_2|_{\mathfrak{k}}$ with highest weight ν_2. From the usual basis of weight vectors for $\bigwedge^i \mathfrak{p}_{\mathbb{C}}^*$ we see that ν_1 may be expressed as $\sum_{\alpha\in A^+} \alpha - \sum_{\alpha \in A^-} \alpha$, where A^+, $A^- \subseteq \Phi^+(\mathfrak{p}_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}})$ and

$$\#A^+ + \#A^- \le i \le \#A^+ + \#A^- + \dim \mathfrak{b}_{G,\mathfrak{p}} \ . \tag{14.1.6}$$

Choose Φ^+ now so that ν_2 (extended by zero on $\mathfrak{b}_{G,\mathfrak{p}}$) is dominant. A calculation [47, Prop. 10.2] using the well-known expression in terms of highest weights for the value of the Casimir operator acting on an irreducible representation shows that Δ_0 is zero on R precisely when

1. the highest weight of R is $\nu_1 + \nu_2$,
2. $\lambda|_{\mathfrak{b}_G} = \nu_2$ (where ν_2 has been extended to be zero on $\mathfrak{b}_{G,\mathfrak{p}}$), and
3. $(\Phi^+(\mathfrak{p}_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) \setminus A^+)$, $A^- \subseteq \Phi^+(\mathfrak{p}(\lambda)_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}})$.

If Δ_0 acts as a nonzero scalar for all R with a fixed i then (14.1.2) will yield (14.1.1) and so $H^i_{(2)}(X;\mathbb{E}) = 0$. So if $H^i_{(2)}(X;\mathbb{E}) \neq 0$ conditions (1)–(3) must hold for some R. Condition (2) implies $\tau(\lambda|_{\mathfrak{b}_G}) = \lambda|_{\mathfrak{b}_G}$ (since $\tau = \theta$ for a fundamental Cartan subalgebra and θ-stable positive system) and thus part (i) of the theorem follows from §8.2. Condition (3) allows one to estimate the possible maximum and minimums values of $\#A^\pm$; inserting these into (14.1.6) yields that

$$\begin{aligned}\#\Phi^+(\mathfrak{p}_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) - \#\Phi^+(\mathfrak{p}(\lambda)_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) \le i \le \\ \#\Phi^+(\mathfrak{p}_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) + \#\Phi^+(\mathfrak{p}(\lambda)_{\mathbb{C}}, \mathfrak{b}_{G,\mathfrak{k}\mathbb{C}}) + \dim \mathfrak{b}_{G,\mathfrak{p}} \ .\end{aligned} \tag{14.1.7}$$

This is equivalent to the degree range in part (ii) by applying (14.1.4) and (14.1.5) and taking the maximum over Φ^+. ☐

15. PROOF OF THEOREM 10.4

By Proposition 13.5 we need to show for all P and V that

$$H^i_{(2)}(\langle A_P^G\rangle_1 \times X_P; (p_*\mathcal{M})_V \otimes \mathbb{C}_{\rho_P}) = 0 \qquad \text{for } i \notin [c(\mathcal{M}), d(\mathcal{M})] \tag{15.0.8}$$

and in addition

$$H^{c(\mathcal{M})}_{(2)}(\langle A_P^G\rangle_1 \times X_P; (p_*\mathcal{M})_V \otimes \mathbb{C}_{\rho_P}) \text{ is Hausdorff.} \tag{15.0.9}$$

We may write the complex C which computes (15.0.8) as
(15.0.10)

$$\begin{cases} C = \bigoplus\limits_{R\ge P} \operatorname{Dom} \overline{d}_{R,b} \otimes \operatorname{Hom}_{L_P}(V, H(\mathfrak{n}_P^R; E_R)) \ , \\ d_C = \sum\limits_{R \ge P} \overline{d}_{R,b} \otimes (-1)^{\text{degree}} + \sum\limits_{R'\ge R\ge P} i_R^{R'*} \otimes \operatorname{Hom}_{L_P}(V, H(\mathfrak{n}_P^{R'}; f_{RR'})) \ , \end{cases}$$

where $\overline{d}_{R,b} = \overline{d}_{\langle A_P^R\rangle_1 \times X_P, b, \mathbb{V}\otimes\mathbb{C}_{\rho_P}}$ and $i_R^{R'*}\colon \operatorname{Dom}\overline{d}_{R',b} \to \operatorname{Dom}\overline{d}_{R,b}$ restricts a form to the smaller stratum.

Let $P' \geq P$ be the parabolic $\mathbb{Q}$-subgroup with type $\Delta_P^{P'} = \{\, \alpha \in \Delta_P \mid \langle \xi_V + \rho, \alpha^\vee \rangle = 0 \,\}$ and set $S = (P, P')$. Then any R with $P \leq R$ belongs to $[\tilde P, \tilde S]$ for a unique $\tilde P \in [P, P']$ and $\tilde S = \tilde P \vee S$:



.

(Compare (3.7.3).) For each such $\tilde P$ let $C(\tilde P)$ be the subquotient complex obtained by restricting the sum in (15.0.10) to be over those $R \in [\tilde P, \tilde S]$. It suffices to prove the vanishing and Hausdorff assertions of (15.0.8) and (15.0.9) for each complex $C(\tilde P)$. (Compare the Fary spectral sequence from Lemma 3.7.)

Filter the double complex $C(\tilde P)$ by the degree of the second factor in (15.0.10); the associated spectral sequence has

$$E_1^{i-k,k} = \bigoplus_{R \in [\tilde P, \tilde S]} H^k_{(2)}(\langle A_P^R \rangle_1 \times X_P; \mathbb{V} \otimes \mathbb{C}_{\rho_P}) \otimes \operatorname{Hom}^{i-k}_{L_P}(V, H(\mathfrak{n}_P^R; E_R)). \tag{15.0.11}$$

Note that we may use ordinary L^2-cohomology here as opposed to L^2-cohomology with boundary values in view of Lemma 11.5.

For $\alpha \in \Delta_P$, let A_P^α denote the one-parameter subgroup corresponding to $\alpha^\vee$. We have a factorization $\langle A_P^R \rangle_1 = \prod_{\alpha \in \Delta_P^R} \langle A_P^\alpha \rangle_1$; for each α, Zucker's L^2-Künneth theorem [56, (2.34)] may be applied to correspondingly factor out the L^2-cohomology of $\langle A_P^\alpha \rangle_1$ in (15.0.11) provided it is Hausdorff. To calculate the result write $\xi_V + \rho = \sum_{\alpha \in \Delta_P} \langle \xi_V + \rho, \alpha^\vee \rangle \beta_\alpha$ and recall that

$$H^{\cdot}_{(2)}(\langle A_P^\alpha \rangle_1; \mathbb{C}_{\langle \xi_V + \rho, \alpha^\vee \rangle \beta_\alpha}) \cong \begin{cases} 0 & \text{if } \langle \xi_V + \rho, \alpha^\vee \rangle > 0, \\ M[-1] & \text{if } \langle \xi_V + \rho, \alpha^\vee \rangle = 0, \\ \mathbb{C} & \text{if } \langle \xi_V + \rho, \alpha^\vee \rangle < 0, \end{cases}$$

where M is an infinite dimensional non-Hausdorff space [41, Prop. 3.2], [56, (2.37)–(2.40)]. Thus let $Q_V \leq S$ have type $\Delta_P^{Q_V} = \{\, \alpha \in \Delta_P \mid \langle \xi_V + \rho, \alpha^\vee \rangle < 0 \,\}$ as in §7.2 and set $\tilde Q = \tilde P \vee Q_V$. We find that

$$E_1^{i-k,k} = H^k_{(2)}(\langle A_P^{\tilde P} \rangle_1 \times X_P; \mathbb{C}_{\xi_V^{-1}} \otimes \mathbb{V}) \otimes \Biggl(\bigoplus_{R \in [\tilde P, \tilde Q]} \operatorname{Hom}^{i-k}_{L_P}(V, H(\mathfrak{n}_P^R; E_R)) \Biggr).$$

The advantage of this expression is that the differentials in the spectral sequence act only on the final factor. These differentials are induced by $H(\mathfrak{n}_P^R; f_{RR'})$ for $\tilde P \leq R \lneq R' \leq \tilde Q$ and so from the definitions in §3.4 we see that $H^i(C(\tilde P))$ equals

$$\bigoplus_k H^k_{(2)}(\langle A_P^{\tilde P} \rangle_1 \times X_P; \mathbb{C}_{\xi_V^{-1}} \otimes \mathbb{V}) \otimes \operatorname{Hom}_{L_P}(V, H^{i-k}(i_P^* i_{\tilde P *} i_{\tilde P}^* \hat\imath_{\tilde Q}^! \mathcal{M})) \;. \tag{15.0.12}$$

By the L^2-Künneth theorem this would be isomorphic to

$$\bigoplus_{k_1, k_2} H^{k_1}_{(2)}(\langle A_P^{\tilde P} \rangle_1) \otimes H^{k_2}_{(2)}(X_P; \mathbb{V}) \otimes \operatorname{Hom}_{L_P}(V, H^{i-k_1-k_2}(i_P^* i_{\tilde P *} i_{\tilde P}^* \hat\imath_{\tilde Q}^! \mathcal{M})) \;; \tag{15.0.13}$$

if at least one of the first two L^2-cohomology factors were finite-dimensional in all degrees, however we cannot assume this. Instead note that by the technique in [56, (2.29)–(2.34)] one can show that if each factor of (15.0.13) vanishes outside a certain interval of degrees, then (15.0.12) vanishes outside the sum of the intervals. The first factor is 0 outside $[0, \dim \mathfrak{a}_P^{\tilde{P}}]$, the vanishing interval of the second factor is given by Theorem 14.1, and the third factor vanishes outside $[c_2(V, \tilde{P}; \mathcal{M}), d_2(V, \tilde{P}; \mathcal{M})]$ by definition (see Proposition 9.2). Thus by Proposition 10.6 we have vanishing for $i \notin [c_2(\mathcal{M}), d_2(\mathcal{M})] = [c(\mathcal{M}), d(\mathcal{M})]$ which proves (15.0.8).

Finally consider (15.0.9). If $P = \tilde{P}$ this follows from the final assertion of Theorem 14.1. If $P \neq \tilde{P}$ then $H^{k_1}_{(2)}(\langle A_P^{\tilde{P}} \rangle_1)$ vanishes in degree 0 (and is non-Hausdorff for $1 \leq k_1 \leq \dim \mathfrak{a}_P^{\tilde{P}}$ [56, (4.51)]) and thus (15.0.12) vanishes in degree $c(\mathcal{M})$ as well. □

Part III. Micro-support Calculations

16. Micro-support of Weighted Cohomology

16.1. Fix a weight profile η. For V an irreducible L_P-module, let $T_V^\eta \geq P$ have type[9]

$$\{\, \alpha \in \Delta_P \mid \langle \xi_V - \eta_P, \beta_\alpha^\vee \rangle < 0 \,\} \,.$$

relative to P. Note that $T_V^\eta = P$ if and only if $\xi_V \in \eta_P + {}^+\mathfrak{a}_P^{G*}$ and $T_V^\eta = G$ if and only if $\xi_V \in \eta_P - \operatorname{int}({}^+\mathfrak{a}_P^{G*})$.

Lemma. *Let V be an irreducible L_P-module occurring in $H(i_P^* \mathcal{W}^\eta \mathcal{C}(E))$. Then V has the form $H^{\ell(w)}(\mathfrak{n}_P; E)_w$ for some $w \in W_P$. Furthermore, the natural morphism $\mathcal{W}^\eta \mathcal{C}(E) \to i_{G*} E$ induces*

$$H(i_P^* \mathcal{W}^\eta \mathcal{C}(E))_V \cong \begin{cases} V[-\ell(w)] & \text{if } T_V^\eta = P, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Apply Proposition 6.4. □

16.2. **Proposition.** *Let V be an irreducible L_P-module occurring in $H(i_P^* \hat{i}_Q^! \mathcal{W}^\eta \mathcal{C}(E))$ for $P \leq Q$. Then V has the form $H^{\ell(w)}(\mathfrak{n}_P; E)_w$ for some $w \in W_P$. Furthermore,*

$$H(i_P^* \hat{i}_Q^! \mathcal{W}^\eta \mathcal{C}(E))_V \cong \begin{cases} V[-\ell(w) - \#\Delta_Q] & \text{if } Q = (P, T_V^\eta), \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Assume first that $T_V^\eta = P$. Then we claim that $H(i_P^* \hat{i}_Q^! \mathcal{W}^\eta \mathcal{C}(E))_V \cong H(i_P^* \hat{i}_Q^! i_{G*} E)_V$ from which proposition follows immediately. To prove the claim, consider the commutative diagram

$$\begin{tikzcd}
\cdots \arrow[r] & H^i(i_P^* \hat{i}_Q^! \mathcal{W}^\eta \mathcal{C}(E))_V \arrow[r] \arrow[d] & H^i(i_P^* \mathcal{W}^\eta \mathcal{C}(E))_V \arrow[r] \arrow[d] & H^i(i_P^* \hat{j}_{Q*} \hat{j}_Q^* \mathcal{W}^\eta \mathcal{C}(E))_V \arrow[r] \arrow[d] & \cdots \\
\cdots \arrow[r] & H^i(i_P^* \hat{i}_Q^! i_{G*} E)_V \arrow[r] & H^i(i_P^* i_{G*} E)_V \arrow[r] & H^i(i_P^* \hat{j}_{Q*} \hat{j}_Q^* i_{G*} E)_V \arrow[r] & \cdots
\end{tikzcd}$$

in which both rows are long exact sequences (obtained from (3.6.4) by setting $Q' = G$ and then taking V-isotypical components). The middle vertical arrow is an isomorphism by Lemma 16.1. To show the right-hand vertical arrow is an isomorphism, consider the corresponding map on the Mayer-Vietoris spectral sequences from Lemma 3.7. For $H(i_P^* \hat{\jmath}_{Q*} \hat{\jmath}_Q^* \mathcal{W}^\eta \mathcal{C}(E))_V$ we have

---

[9] This type corresponds to what in [25] would be denoted $I_\eta(\xi_V)$.

$$E_1^{p,\cdot} = \bigoplus_{\substack{P < \tilde{P} \leq (P,Q) \\ \#\Delta_P^{\tilde{P}} = p+1}} H(\mathfrak{n}_P^{\tilde{P}}; H(i_{\tilde{P}}^* \mathcal{W}^\eta \mathcal{C}(E)))_V \ . \tag{16.2.1}$$

The $\tilde{P}$-term can be rewritten $H(\mathfrak{n}_P^{\tilde{P}}; H(i_{\tilde{P}}^* \mathcal{W}^\eta \mathcal{C}(E))_{\tilde{V}})_V$ for an irreducible $L_{\tilde{P}}$-module $\tilde{V}$ satisfying $\xi_{\tilde{V}} = \xi_V|_{\mathfrak{a}_{\tilde{P}}}$ (see §0.10.22). By Lemma 16.1 this is isomorphic to the corresponding term in E_1 for $H(i_P^* \hat{\jmath}_{Q*} \hat{\jmath}_Q^* i_{G*} E)_V$ provided that $T_{\tilde{V}}^\eta = \tilde{P}$. But this holds since $T_{\tilde{V}}^\eta = T_V^\eta \vee \tilde{P}$. The claim then follows from the 5-lemma.

In the case that $T_V^\eta > P$, Lemma 16.1 shows that $H(i_P^* \mathcal{W}^\eta \mathcal{C}(E))_V = 0$ and hence $H(i_P^* \hat{\imath}_Q^! \mathcal{W}^\eta \mathcal{C}(E))_V \cong H(i_P^* \hat{\jmath}_{Q*} \hat{\jmath}_Q^* \mathcal{W}^\eta \mathcal{C}(E))_V[-1]$. It also shows that the sum in (16.2.1) yields one copy of $V[-\ell(w)]$ for each $\tilde{P} \in [T_V^\eta, (P,Q)]$. The differential between two such copies of $V[-\ell(w)]$ for adjacent $\tilde{P}$ will be an isomorphism by comparison with the spectral sequence for $H(i_P^* \hat{\jmath}_{Q*} \hat{\jmath}_Q^* i_{G*} E)_V$. Consequently $H(i_P^* \hat{\jmath}_{Q*} \hat{\jmath}_Q^* \mathcal{W}^\eta \mathcal{C}(E))_V = 0$ unless $T_V^\eta = (P,Q)$, in which case we obtain $V[-\ell(w) - \#\Delta_Q + 1]$. This proves the proposition. □

16.3. Let η be one of the two *middle weight profiles*:

$$\textit{upper middle}\colon\ \mu = -\rho + \epsilon\rho \ ,$$
$$\textit{lower middle}\colon\ \nu = -\rho \ .$$

Here $\epsilon > 0$ is chosen sufficiently small such that $\epsilon\rho_R < \chi_R$ for all $R \in \mathcal{P}$ maximal, where χ_R is a positive generator for $X(S_R^G)$.

Theorem. *Let E be an irreducible regular G-module and let η be a middle weight profile. The weak micro-support* $\mathrm{SS}_w(\mathcal{W}^\eta \mathcal{C}(E))$ *consists of those irreducible L_P-modules V such that*

(i) $V = H^{\ell(w)}(\mathfrak{n}_P; E)_w$ *with* $w \in W_P$, *and*
(ii) $(\xi_V + \rho)|_{\mathfrak{a}_P^G} = 0$.

For such V we have

$$c(V;\mathcal{M}) = d(V;\mathcal{M}) = \begin{cases} \ell(w) + \Delta_P & \textit{if } \eta = \mu, \\ \ell(w) & \textit{if } \eta = \nu. \end{cases}$$

The micro-support $\mathrm{SS}(\mathcal{W}^\eta \mathcal{C}(E))$ *consists of those* $V \in \mathrm{SS}_w(\mathcal{W}^\eta \mathcal{C}(E))$ *such that*

(iii) $V|_{M_P}$ *is conjugate self-contragredient*;

in this case w from (i) *is fundamental. The micro-support is nonempty if and only if* $E|_{{}^0G}$ *is conjugate self-contragredient*; *in this case* $\mathrm{SS}_{\mathrm{ess}}(\mathcal{W}^\eta \mathcal{C}(E)) = \{E\}$.

Proof of Theorem 16.3. By Proposition 16.2, an irreducible L_P-module V is in $\mathrm{SS}(\mathcal{W}^\eta \mathcal{C}(E))$ if and only if (i) and

(ii)′ $(P, T_V^\eta) \in [Q_V, Q_V']$

hold. Assume that $\eta = \nu = -\rho$. Then Condition (ii)′ is equivalent to

(ii)″ $\langle \xi_V + \rho, \alpha^\vee \rangle < 0 \quad \Rightarrow \quad \langle \xi_V + \rho, \beta_\alpha^\vee \rangle \geq 0 \quad \Rightarrow \quad \langle \xi_V + \rho, \alpha^\vee \rangle \leq 0$

for all $\alpha \in \Delta_P$. For $R \geq P$ define

$$\begin{aligned}\langle \mathfrak{a}_P^* \rangle_R &= \operatorname{int} \mathfrak{a}_R^{*+} - {}^+\mathfrak{a}_P^{R*} \\ &= \{\, \zeta \in \mathfrak{a}_P^* \mid \langle \zeta, \gamma^\vee \rangle > 0 \text{ for } \gamma \in \Delta_R, \text{ and} \\ &\qquad\qquad \langle \zeta, \beta_\alpha^{R\vee} \rangle \leq 0 \text{ for } \alpha \in \Delta_P^R \,\}.\end{aligned}$$

By Langlands's "geometric lemmas" [13, IV,§6.11] these sets form a disjoint decomposition of $\mathfrak{a}_P^*$, so

$$\xi_V + \rho_P \in -\langle \mathfrak{a}_P^* \rangle_R \tag{16.3.1}$$

for a unique $R \geq P$. Now for any $\alpha \in \Delta_P \setminus \Delta_P^R$, write $\alpha^\vee = \alpha_R^\vee + \alpha^{\vee R} = \gamma^\vee + \alpha^{\vee R}$, where $\gamma = \alpha_R \in \Delta_R$. Since $\alpha^{\vee R} \in -\mathfrak{a}_P^{R+}$, we see from (16.3.1) that $\langle \xi_V + \rho, \alpha^\vee \rangle < 0$. Thus by the first implication of (ii)″, $(\xi_V + \rho)_R \in {}^+\mathfrak{a}_R^*$ which contradicts (16.3.1). Consequently we have $R = G$, $\xi_V + \rho \in {}^+\mathfrak{a}_P^*$, and $G = (P, T_V^\eta)$. But then the second implication of (ii)″ implies $\xi_V + \rho \in -\mathfrak{a}_P^{*+}$ and thus (ii) must hold. Conversely if (ii) holds, then (ii)″ is automatic. The fact that $c(V; \mathcal{M}) = d(V; \mathcal{M}) = \ell(w)$ comes from Proposition 16.2.

The case of $\eta = \mu$ is similar except that (ii)″ is replaced by

(ii)‴ $\langle \xi_V + \rho, \alpha^\vee \rangle < 0 \quad \Rightarrow \quad \langle \xi_V + \rho, \beta_\alpha^\vee \rangle > 0 \quad \Rightarrow \quad \langle \xi_V + \rho, \alpha^\vee \rangle \leq 0$

and we consider the disjoint decomposition of $\mathfrak{a}_P^*$ given by

$$\begin{aligned}\langle\langle \mathfrak{a}_P^* \rangle\rangle_R &= \mathfrak{a}_R^{*+} - \operatorname{int} {}^+\mathfrak{a}_P^{R*} \\ &= \{\, \zeta \in \mathfrak{a}_P^* \mid \langle \zeta, \gamma^\vee \rangle \geq 0 \text{ for } \gamma \in \Delta_R, \text{ and} \\ &\qquad\qquad \langle \zeta, \beta_\alpha^{R\vee} \rangle < 0 \text{ for } \alpha \in \Delta_P^R \,\}\end{aligned}$$

for all $R \geq P$. We find that (ii)‴ is equivalent to (ii) and that in this case $\xi_V + \rho_P \in -\langle\langle \mathfrak{a}_P^* \rangle\rangle_R$ with $R = P = T_V^\eta$.

The assertion that w is fundamental for $V \in \operatorname{SS}(\mathcal{W}^\eta\mathcal{C}(E))$ follows from (ii) and (iii) by Lemma 8.8. For the final assertion, note that we can rephrase our result as $V \in \operatorname{SS}(\mathcal{W}^\eta\mathcal{C}(E))$ if and only if $V \preccurlyeq_0 E$ and $V|_{M_P}$ is conjugate self-contragredient. By Lemma 9.1 the existence of such a V implies $E|_{{}^0G}$ is conjugate self-contragredient. Also note that if $V \in \operatorname{SS}_{\mathrm{ess}}(\mathcal{W}^\eta\mathcal{C}(E))$ then $Q_V = Q'_V$ by Proposition 16.2 and therefore $\langle \xi_V + \rho, \alpha^\vee \rangle \neq 0$ for all $\alpha \in \Delta_P$. In view of (ii) we must have $P = G$ and thus $V = E$. □

17. MICRO-SUPPORT OF INTERSECTION COHOMOLOGY

In §§17, 18, and 20, we consider a *middle perversity* p, either $m(k) = \lfloor \frac{k-2}{2} \rfloor$ or $n(k) = \lfloor \frac{k-1}{2} \rfloor$.

17.1. **Theorem.** *Assume the irreducible components of the $\mathbb{Q}$-root system of G are of type A_n, B_n, C_n, BC_n, or G_2. Let E be an irreducible regular G-module and let p be a middle perversity. The micro-support $\operatorname{SS}(\mathcal{I}_p\mathcal{C}(E))$ consists of those irreducible L_P-modules V such that*

(i) $V = H^{\ell(w)}(\mathfrak{n}_P; E)_w$ *with $w \in W_P$ a fundamental Weyl element for P in G;*

(ii) $$\langle \xi_V + \rho, \alpha^\vee \rangle \begin{cases} \geq 0 & \textit{for all } \alpha \in \Delta_P \ (\textit{if } p = m), \\ \leq 0 & \textit{for all } \alpha \in \Delta_P \ (\textit{if } p = n); \end{cases}$$

(iii) $V|_{M_P}$ *is conjugate self-contragredient.*

For such a L_P-module V we have

$$c(V;\mathcal{M}) = d(V;\mathcal{M}) = \begin{cases} \ell(w) + \Delta_P & \text{if } p = m, \\ \ell(w) & \text{if } p = n. \end{cases}$$

The essential micro-support $\mathrm{SS}_{\mathrm{ess}}(\mathcal{I}_p\mathcal{C}(E))$ *consists of the maximal elements of* $\mathrm{SS}(\mathcal{I}_p\mathcal{C}(E))$ *under* $\preccurlyeq_0$ (*see* §9.1); *this is equivalent to requiring strict inequalities in* (ii).

Unlike the case of weighted cohomology, the weak micro-support is not characterized by simply omitting (iii).

17.2**. Corollary.** *In the setting of Theorem* 17.1 *assume that* $E|_{{}^0G}$ *is conjugate self-contragredient. Then* $\mathrm{SS}_{\mathrm{ess}}(\mathcal{I}_p\mathcal{C}(E)) = \{E\}$ *with* $c(E;\mathcal{I}_p\mathcal{C}(E)) = d(E;\mathcal{I}_p\mathcal{C}(E)) = 0$ *and* $(\xi_V+\rho)|_{\mathfrak{a}_P^G} = 0$ *for all* $V \in \mathrm{SS}(\mathcal{I}_p\mathcal{C}(E))$.

17.3. In the remainder of this section we will reduce the theorem to a combinatorial vanishing result which will be proven in the following sections.

Recall the "combinatorial" intersection cohomology $I_{p_w}H(U)$ (for U an open constructible subset of $|\Delta_P|$ or $c(|\Delta_P|)$) defined in §5.5 which depends on $w \in W$ as well as p. We also consider the version with supports $I_{p_w}H_Z(U)$ defined for Z constructible and relatively closed in U. This may be interpreted in terms of relative intersection cohomology:

$$I_{p_w}H_Z(U) = I_{p_w}H(U, U \setminus Z)\ .$$

17.4**. Proposition.** *Let* $P \le Q \in \mathcal{P}$. *Then*

$$H(i_P^*\hat{i}_Q^!\mathcal{I}_p\mathcal{C}(E)) \cong \bigoplus_{w\in W_P} H(\mathfrak{n}_P;E)_w \otimes I_{p_w}H_{c(|\Delta_P^Q|)}(c(|\Delta_P|))\ .$$

Proof. There is a long exact sequence (set $Q' = G$ in (3.6.4))

$$\ldots \longrightarrow H^j(i_P^*\hat{i}_Q^!\mathcal{I}_p\mathcal{C}(E)) \longrightarrow H^j(i_P^*\mathcal{I}_p\mathcal{C}(E)) \longrightarrow H^j(i_P^*\hat{j}_{Q*}\hat{j}_Q^*\mathcal{I}_p\mathcal{C}(E)) \longrightarrow \cdots\ .$$

On the other hand there is a topological long exact sequence

$$\cdots \to I_{p_w}H^i_{c(|\Delta_P^Q|)}(c(|\Delta_P|)) \to I_{p_w}H^i(c(|\Delta_P|)) \to I_{p_w}H^i(c(|\Delta_P|)\setminus c(|\Delta_P^Q|)) \to \cdots\ .$$

We need to show that if the first term of the second sequence is tensored with $H(\mathfrak{n}_P;E)_w$ and then direct summed over $w \in W_P$, then the result is isomorphic to the first term of the first sequence. This is true for the middle term by Proposition 5.5. It is also true for the last term: compare the Mayer-Vietoris sequences abutting to each and apply Proposition 5.5 again. One may check that these isomorphisms are compatible with the maps in the long exact sequences. Application of the 5-Lemma concludes the proof. □

17.5. From the proposition it follows that if an irreducible L_P-module V occurs in $\mathrm{SS}(\mathcal{I}_p\mathcal{C}(E))$, then $V = H^{\ell(w)}(\mathfrak{n}_P;E)_w$ for some $w \in W_P$, and this V will actually occur only if $V|_{M_P}$ is conjugate self-contragredient and $I_{p_w}H_{c(|\Delta_P^Q|)}(c(|\Delta_P|) \neq 0$ for some $Q \in [Q_V, Q'_V]$ (see §7.1). This condition on Q is equivalent to

$$\begin{aligned} \langle \xi_V + \rho, \alpha^\vee\rangle &\le 0 \qquad \text{for } \alpha \in \Phi(\mathfrak{n}_P^Q;\mathfrak{a}_P), \\ \langle \xi_V + \rho, \alpha^\vee\rangle &\ge 0 \qquad \text{for } \alpha \in \Phi(\mathfrak{n}_P^{P,Q};\mathfrak{a}_P). \end{aligned} \tag{17.5.1}$$

17.6. Thus we need to show $I_{p_w} H_{c(|\Delta_P^Q|)}(c(|\Delta_P|)$ vanishes except for the cases indicated in Theorem 17.1; this is a combinatorial statement involving the data $\ell(w_R)$ for $R \geq P$. However our hypotheses give us information on V, or equivalently, its highest weight. Following an argument from [47], the simple basic lemma below will translate the weight information into combinatorial information; it is a generalization of a result of Casselman [16, Prop. 2.6] in the real rank one case. A stronger result will be presented in §24.

Let $\mathfrak{h} = \mathfrak{b}_P + \mathfrak{a}_P$ be a Cartan subalgebra of $\mathfrak{l}_P$ and fix a positive system Φ^+ of $\Phi(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ containing $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. For $\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)$, let $\mathfrak{n}_\alpha \subseteq \mathfrak{n}_P$ be the corresponding root space and write $\Phi(\mathfrak{n}_{\alpha\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) = \{\, \gamma \in \Phi^+ \mid \gamma|_{\mathfrak{a}_P} = \alpha \,\}$. For $w \in W$, the contribution to $\ell(w)$ attributable to α is $\ell_\alpha(w) = \#(\Phi(\mathfrak{n}_\alpha, \mathfrak{h}_{\mathbb{C}}) \cap \Phi_w)$. Note that $\ell_\alpha(w) = \ell_\alpha(w_P)$ for $\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)$. Also if $w \in W_P$ then $\ell(w) = \sum_{\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)} \ell_\alpha(w)$.

Lemma. *Let P be a parabolic $\mathbb{Q}$-subgroup and let $w \in W_P$. Let $V = H^{\ell(w)}(\mathfrak{n}_P; E)_w$ for an irreducible G-module E and assume $V|_{M_P}$ is conjugate self-contragredient. For any $\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)$ we have*:

(i) $\langle \xi_V + \rho, \alpha^\vee \rangle \leq 0 \implies \ell_\alpha(w) \geq \frac{1}{2} \dim \mathfrak{n}_\alpha$.
(ii) $\langle \xi_V + \rho, \alpha^\vee \rangle = 0 \implies \ell_\alpha(w) = \frac{1}{2} \dim \mathfrak{n}_\alpha$.
(iii) $\langle \xi_V + \rho, \alpha^\vee \rangle \geq 0 \implies \ell_\alpha(w) \leq \frac{1}{2} \dim \mathfrak{n}_\alpha$.

Furthermore if $\gamma \in \Phi(\mathfrak{n}_\alpha, \mathfrak{h}_{\mathbb{C}})$, then Φ_w contains respectively (i) *at least one,* (ii) *exactly one,* (iii) *at most one of γ and $\tau'_P\gamma$.*

Proof. Let $\lambda \in \mathfrak{h}_{\mathbb{C}}$ be the highest weight of E; by Kostant's theorem the highest weight of V is then $w(\lambda + \rho) - \rho$ and $\xi_V + \rho = w(\lambda + \rho)|_{\mathfrak{a}_P}$.

Let $\gamma \in \Phi(\mathfrak{n}_\alpha, \mathfrak{h}_{\mathbb{C}})$. Since $w(\lambda+\rho)$ is regular and λ is dominant, $\gamma \in \Phi_w$ if and only if $\langle w(\lambda + \rho), \gamma^\vee \rangle \leq 0$. Decompose $\mathfrak{b}_P = \mathfrak{b}_{P,1} + \mathfrak{b}_{P,-1}$ according to the eigenvalues of τ_P; by §§8.2, 8.3 the hypothesis that $V|_{M_P}$ is conjugate self-contragredient is equivalent to $w(\lambda + \rho)|_{\mathfrak{b}_{P,-1}} = 0$. On the other hand, $\gamma^\vee|_{\mathfrak{a}_P} = c\alpha$ where $c > 0$. Thus

$$\langle w(\lambda + \rho), \gamma^\vee \rangle = \langle w(\lambda + \rho)|_{\mathfrak{b}_{P,1}}, \gamma^\vee \rangle + c\langle w(\lambda + \rho), \alpha^\vee \rangle \ .$$

Replacing γ by $\tau'_P\gamma$ in this equation negates the first term and leaves the second term unchanged, so $\langle w(\lambda + \rho), \alpha^\vee \rangle \leq 0$ implies $\langle w(\lambda + \rho), \gamma^\vee \rangle \leq 0$ or $\langle w(\lambda + \rho), \tau'_P\gamma^\vee \rangle \leq 0$. Thus either γ or $\tau'_P\gamma$ is in Φ_w. This proves (i) and the proof of (iii) is similar; (ii) follows from (i) and (iii). □

By (17.5.1) and the lemma we have

$$\begin{aligned} &\ell_\alpha(w) \geq \tfrac{1}{2} \dim \mathfrak{n}_\alpha \qquad \text{for } \alpha \in \Phi(\mathfrak{n}_P^Q; \mathfrak{a}_P), \\ &\ell_\alpha(w) \leq \tfrac{1}{2} \dim \mathfrak{n}_\alpha \qquad \text{for } \alpha \in \Phi(\mathfrak{n}_P^{P,Q}; \mathfrak{a}_P). \end{aligned} \tag{17.6.1}$$

17.7. We begin with a special case of the vanishing theorem. The proof will appear in §18; the key point is that in the case $Q = P$ or G the hypotheses (17.6.1) are preserved under passing by induction from P to $R > P$.

Proposition. *Let p be a middle perversity and let $w \in W$. Let $P \leq Q$ be parabolic subgroups so that* (17.6.1) *is satisfied. Also assume that $Q = P$ or G. Then*

$$I_{p_w} H_{c(|\Delta_P^Q|)}(c(|\Delta_P|)) = \begin{cases} \mathbb{Z}[-\#\Delta_P] & \textit{if } Q = P,\ p = m, \textit{ and } \ell(w_P) = \frac{1}{2} \dim \mathfrak{n}_P, \\ \mathbb{Z} & \textit{if } Q = G,\ p = n, \textit{ and } \ell(w_P) = \frac{1}{2} \dim \mathfrak{n}_P, \\ 0 & \textit{otherwise.} \end{cases}$$

17.8. For $P < Q < G$ the analogue of (17.6.1) will no longer remain true when we pass by induction to $R > P$. Nor will $H(\mathfrak{n}_R; E)_{w_R}$ necessarily be conjugate self-contragredient (for any E) so Lemma 17.6 will no longer apply directly. The following lemma yields an alternate hypothesis that is suitable for induction and is sufficient.

Lemma. *Assume the irreducible components of the $\mathbb{Q}$-root system of G are of type A_n, B_n, C_n, BC_n, or G_2. Let P be a proper parabolic subgroup and let $w \in W$. Let $V = H^{\ell(w)}(\mathfrak{n}_P; E)_w$ for an irreducible G-module E and assume $V|_{M_P}$ is conjugate self-contragredient. Assume $Q \geq P$ is a parabolic $\mathbb{Q}$-subgroup such that* (17.5.1) *is satisfied. Then there exists $T \geq P$ so that*

$$(17.8.1) \qquad 1 \leq \#\Delta_T \leq 2 \qquad \text{and} \qquad 0 \leq \#\Delta_T^{Q\vee T},\ \#(\Delta_T \setminus \Delta_T^{Q\vee T}) \leq 1$$

and that

(17.8.2)

$$\ell_\alpha(w) \geq \tfrac{1}{2}\dim \mathfrak{n}_\alpha \qquad \text{for } \alpha \in \Phi(\mathfrak{n}_P; \mathfrak{a}_P) \text{ such that } \alpha|_{\mathfrak{a}_T} \in \Phi(\mathfrak{n}_T^{Q\vee T}; \mathfrak{a}_T),$$

$$\ell_\alpha(w) \leq \tfrac{1}{2}\dim \mathfrak{n}_\alpha \qquad \text{for } \alpha \in \Phi(\mathfrak{n}_P; \mathfrak{a}_P) \text{ such that } \alpha|_{\mathfrak{a}_T} \in \Phi(\mathfrak{n}_T^{T,Q\vee T}; \mathfrak{a}_T).$$

Furthermore, if $P < Q < G$ we can assume that

$$(17.8.3) \qquad \ell(w_T) \neq \tfrac{1}{2}\dim \mathfrak{n}_T \text{ when } \#\Delta_T = 1.$$

The lemma will be proved in §19. When $P < Q < G$ this lemma provides the hypotheses for the following theorem:

17.9**. Theorem.** *Let p be a middle perversity and let $w \in W$. Let $P \leq Q$ be parabolic $\mathbb{Q}$-subgroups and assume there exists $T \geq P$ satisfying* (17.8.1), (17.8.2), *and* (17.8.3). *Then*

$$I_{p_w} H_{c(|\Delta_P^Q|)}(c(|\Delta_P|)) = 0.$$

The proof will appear in §20.

17.10. **Proof of Theorem 17.1.** Note that the nonvanishing cases of Proposition 17.7 are relevant only when $Q_V = P$ (for $p = m$) and $Q'_V = G$ (for $p = n$). Together with Theorem 17.9, this implies that an irreducible L_P-module V belongs to $\mathrm{SS}(\mathcal{I}_p\mathcal{C}(E))$ if and only if

(i′) $V = H^{\ell(w)}(\mathfrak{n}_P; E)_w$ with $w \in W_P$ satisfying $\ell(w) = \frac{1}{2}\dim \mathfrak{n}_P$;

together with (iii) and (ii) of Theorem 17.1 hold. The last assertion of Lemma 17.6 implies in this case that τ'_P interchanges Φ_w and $\Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C}) \setminus \Phi_w$; in other words, w is a fundamental Weyl element for P in G by Lemma 8.6. Thus (i)$\iff$(i′) given (iii) and (ii). Such a V will lie in $\mathrm{SS}_{\mathrm{ess}}(\mathcal{I}_p\mathcal{C}(E))$ furthermore if and only if $Q_V = Q'_V$, that is, if and only if the inequalities in (ii) are all strict. This is equivalent to V being maximal in $\mathrm{SS}(\mathcal{I}_p\mathcal{C}(E))$ by Lemma 8.8. □

18. PROOF OF PROPOSITION 17.7

First assume that $\ell(w_P) = \frac{1}{2}\dim \mathfrak{n}_P$. This implies $\ell(w_R) = \frac{1}{2}\dim \mathfrak{n}_R$ for all $R \geq P$ by (17.6.1) and so

$$p_w(R) = \begin{cases} \left\lfloor \frac{\#\Delta_R - 2}{2} \right\rfloor & \text{if } p = m, \\ \left\lfloor \frac{\#\Delta_R - 1}{2} \right\rfloor & \text{if } p = n. \end{cases}$$

For $p = m$ we see that $p_w(R) < \#\Delta_R - 1$ for $R < G$ and thus the link cohomology for the R-stratum (which inductively is $\mathbb{Z}[-\#\Delta_R + 1]$) will always be truncated. For $p = n$ we see that $p_w(R) \geq 0$ and thus the link cohomology (which inductively is $\mathbb{Z}$) will never be truncated. This proves the proposition in this case.

Now assume that $\ell(w_P) \neq \frac{1}{2} \dim \mathfrak{n}_P$ and consider the case $Q = P$. Let $S > P$ be the unique parabolic $\mathbb{Q}$-subgroup which is minimal with the property that $\ell(w_S) = \frac{1}{2} \dim \mathfrak{n}_S$. Consider the Fary spectral sequence

$$E_1^{-p,i+p} = \bigoplus_{\#\Delta_P^R = p} I_{p_w} H^i_{|\Delta_P^R|^\circ}(|\Delta_P|) \cong \bigoplus_{\#\Delta_P^R = p} I_{p_w} H^i_{c(\emptyset)}(c(|\Delta_R|)) \Rightarrow I_{p_w} H^i(|\Delta_P|).$$

The hypothesis (17.6.1) remains valid if we replace P by R (with $Q = R$) so by induction all the E_1 terms vanish except in the case that $p = m$. In this case, only the terms for $R \in [S, G]$ are nonvanishing and the spectral sequence is isomorphic to the Fary spectral sequence abutting to $I_{p_w} H^i(c(|\Delta_S|))$. This vanishes by the argument in the previous paragraph unless $S = G$. Thus

$$I_{p_w} H(|\Delta_P|) = \begin{cases} 0 & \text{if } S < G \text{ or } p = n, \\ \mathbb{Z} & \text{if } S = G \text{ and } p = m. \end{cases}$$

However in this second case, $p_w(P) \geq \left\lfloor \frac{1 + \#\Delta_P - 2}{2} \right\rfloor \geq 0$ and so a degree 0 link cohomology class will not be truncated at the P-stratum. Thus $I_{p_w} H_{c(\emptyset)}(c(|\Delta_P|)) = 0$.

The argument for the case $Q = G$ is similar except that we use the Mayer-Vietoris spectral sequence converging to $I_{p_w} H^i(|\Delta_P|)$ from (5.5.2):

$$E_1^{p,i-p} = \bigoplus_{\#\Delta_P^R = p+1} I_{p_w} H^{i-p}(U_R) \cong \bigoplus_{\#\Delta_P^R = p+1} I_{p_w} H^{i-p}_{c(|\Delta_R|)}(c(|\Delta_R|)) \,. \qquad \square$$

19. PROOF OF LEMMA 17.8

We begin with some generalities on chambers in $\mathfrak{a}_P$ and an analogue of the lemma for the $\mathbb{Q}$-Weyl group ${}_{\mathbb{Q}}W$ of G (Lemma 19.4). This is a purely combinatorial result regarding the $\mathbb{Q}$-root system and its Weyl group. We then reduce Lemma 17.8 to this result. Thus w in this section will refer to an element of ${}_{\mathbb{Q}}W$ until §19.5.

19.1. Let ${}_{\mathbb{Q}}\Phi \subset \mathfrak{a}^*$ denote the $\mathbb{Q}$-root system of G with simple roots Δ. Let P be a parabolic $\mathbb{Q}$-subgroup and let ${}_{\mathbb{Q}}\Phi^P$ denote the subroot system with basis Δ^P in $\mathfrak{a}^{P*}$. The Weyl group of ${}_{\mathbb{Q}}\Phi^P$ is the parabolic subgroup ${}_{\mathbb{Q}}W^P \subseteq {}_{\mathbb{Q}}W$ generated by the simple reflections $\{s_\alpha\}_{\alpha \in \Delta^P}$. Let ${}_{\mathbb{Q}}\Phi_P$ denote the restriction of the elements${}_{\mathbb{Q}}\Phi \setminus {}_{\mathbb{Q}}\Phi^P$ to $\mathfrak{a}_P$. An element of ${}_{\mathbb{Q}}\Phi_P$ may be expressed as a $\mathbb{Z}$-linear combination of the elements of Δ_P with coefficients all ≥ 0 or ≤ 0, however in general ${}_{\mathbb{Q}}\Phi_P$ is not a root system.

19.2. **Chambers in $\mathfrak{a}_P$.** The "chambers" of $\mathfrak{a}_P$ are the connected components of $\mathfrak{a}_P \setminus \bigcup_{\alpha \in {}_{\mathbb{Q}}\Phi_P} \operatorname{Ker} \alpha$. Even though ${}_{\mathbb{Q}}\Phi_P$ is not a root system there is a combinatorial object that parametrizes these chambers. Let ${}_{\mathbb{Q}}W_P$ denote the minimal length representatives of the cosets in ${}_{\mathbb{Q}}W^P \backslash {}_{\mathbb{Q}}W$ and define

(19.2.1) $\qquad {}_{\mathbb{Q}}W(P) = \{\, w \in {}_{\mathbb{Q}}W \mid w^{-1}\Delta^P \subseteq \Delta \,\} \subseteq {}_{\mathbb{Q}}W_P$

(cf. [35, I.1.7][10]). This is not a subgroup in general.

There is a nice geometric way to think of ${}_{\mathbb{Q}}W_P$ and ${}_{\mathbb{Q}}W(P)$. Namely let C denote the strictly dominant cone of ${}_{\mathbb{Q}}\Phi$ in $\mathfrak{a}$ and let C^P denote the strictly dominant cone of ${}_{\mathbb{Q}}\Phi^P$ in $\mathfrak{a}^P$. Then $w \in {}_{\mathbb{Q}}W_P$ if and only if the chamber wC projects into C^P under the projection $\mathfrak{a} \to \mathfrak{a}^P$. Furthermore such an element belongs to ${}_{\mathbb{Q}}W(P)$ if and only if the closure of wC is adjacent to $\mathfrak{a}_P$ in the sense that $w\overline{C} \cap \mathfrak{a}_P$ is a boundary stratum of $w\overline{C}$ with codimension $\#\Delta^P$. In this case we let $w \cdot C_P \equiv \operatorname{int}_{\mathfrak{a}_P}(w\overline{C} \cap \mathfrak{a}_P)$ denote the corresponding chamber of $\mathfrak{a}_P$. (Here $\operatorname{int}_{\mathfrak{a}_P}$ denotes the interior of a subset of $\mathfrak{a}_P$.)

The correspondence

$$w \quad \longleftrightarrow \quad w \cdot C_P \tag{19.2.2}$$

is a bijection of ${}_{\mathbb{Q}}W(P)$ with the chambers of $\mathfrak{a}_P$ [35, I.1.10].

19.3. **An Alternative Characterization of ${}_{\mathbb{Q}}W(P)$.** Let pr_P denote the map ${}_{\mathbb{Q}}\Phi \setminus {}_{\mathbb{Q}}\Phi^P \to {}_{\mathbb{Q}}\Phi_P$ given by restriction to $\mathfrak{a}_P$. For $w \in {}_{\mathbb{Q}}W$ let ${}_{\mathbb{Q}}\Phi_w = \{\gamma \in {}_{\mathbb{Q}}\Phi^+ | w^{-1}\gamma < 0\}$.

Lemma. ${}_{\mathbb{Q}}W(P) = \{\, w \in {}_{\mathbb{Q}}W_P \mid {}_{\mathbb{Q}}\Phi_w \cap \operatorname{pr}_P^{-1}(\alpha) = \operatorname{pr}_P^{-1}(\alpha)$ *or* $\emptyset$ *for all* $\alpha \in {}_{\mathbb{Q}}\Phi_P^+ \}$. (*The cases* $\operatorname{pr}_P^{-1}(\alpha)$ *and* $\emptyset$ *corresponding respectively to* $\langle \alpha, w \cdot C_P\rangle < 0$ *and* $\langle \alpha, w \cdot C_P\rangle > 0$.)

Proof. If $w \in {}_{\mathbb{Q}}W(P)$ and $\gamma \in \operatorname{pr}_P^{-1}(\alpha)$ we have $\gamma \in {}_{\mathbb{Q}}\Phi_w \Leftrightarrow \langle \gamma, wC\rangle < 0 \Leftrightarrow \langle \gamma, w\overline{C}\rangle \leq 0 \Leftrightarrow \langle \alpha, w \cdot C_P\rangle < 0$. Conversely, if $w \in {}_{\mathbb{Q}}W_P$ and ${}_{\mathbb{Q}}\Phi_w \cap \operatorname{pr}_P^{-1}(\alpha) = \operatorname{pr}_P^{-1}(\alpha)$ or $\emptyset$ for all $\alpha \in {}_{\mathbb{Q}}\Phi_P^+$, consider $\gamma \in \Delta^P$. If $w^{-1}\gamma = \delta_1 + \delta_2$ (δ_1 and δ_2 positive roots) is not simple, then $w\delta_1 > 0$ (say) and $w\delta_2 < 0$. Thus $-w\delta_2 \in {}_{\mathbb{Q}}\Phi_w$ and $-w\delta_2 + \gamma \notin {}_{\mathbb{Q}}\Phi_w$, while $\operatorname{pr}_P(-w\delta_2) = \operatorname{pr}_P(-w\delta_2 + \gamma)$. This contradiction implies $w^{-1}\Delta^P \subset \Delta$ and thus $w \in {}_{\mathbb{Q}}W(P)$. $\square$

19.4. **Lemma.** *Assume the irreducible components of the $\mathbb{Q}$-root system of G are of type A_n, B_n, C_n, BC_n, or G_2. Let P be a proper parabolic $\mathbb{Q}$-subgroup and let $w \in {}_{\mathbb{Q}}W(P)$. Then there exists $T \geq P$ such that one of the following hold*:

Case 1: $\#\Delta_T = 1$. *In this case* ${}_{\mathbb{Q}}\Phi_w \cap \operatorname{pr}_T^{-1}({}_{\mathbb{Q}}\Phi_T^+) = \operatorname{pr}_P^{-1}({}_{\mathbb{Q}}\Phi_T^+)$ *or* $\emptyset$.

Case 2: $\#\Delta_T = 2$. *In this case* $\Delta_T = \{\beta_-, \beta_+\}$ *and*

$$\begin{aligned} {}_{\mathbb{Q}}\Phi_w \cap \operatorname{pr}_T^{-1}(\mathbb{N}\beta_-) &= \operatorname{pr}_T^{-1}(\mathbb{N}\beta_-), \\ {}_{\mathbb{Q}}\Phi_w \cap \operatorname{pr}_T^{-1}(\mathbb{N}\beta_+) &= \emptyset. \end{aligned}$$

Remark. The lemma does not quite imply that $w_T \in {}_{\mathbb{Q}}W(T)$ since in Case 2 nothing is asserted regarding ${}_{\mathbb{Q}}\Phi_w \cap \operatorname{pr}_T^{-1}(\beta)$ when $\beta \in {}_{\mathbb{Q}}\Phi_T^+$ is not a multiple of a simple root.

Proof. It suffices to assume that ${}_{\mathbb{Q}}\Phi$ is irreducible and even reduced (the case BC_n follows from the case B_n). Let $\widetilde{G}$ be the simply connected semisimple $\mathbb{Q}$-split group with root system ${}_{\mathbb{Q}}\Phi$. There always exists an irreducible representation V of $\widetilde{G}$ whose nonzero weights form a single Weyl orbit [32, §2], [28]. We partially order the weights $\Phi(V)$ as usual: $\mu \prec \nu$ if $\nu - \mu$ is a sum of nonnegative integral multiples of elements of Δ. Then the root systems A_n, B_n, C_n, and G_2 are precisely those for which we can arrange that $\Phi(V)$ is totally ordered. (In the notation of the appendices of [14] we choose V to have highest weight ω_1.) Write $\Phi(V) = \{\varepsilon_1 \succ e_2 \succ \cdots \succ \varepsilon_N\}$. Since the representation is faithful every root $\gamma \in {}_{\mathbb{Q}}\Phi^+$ may be expressed (nonuniquely) as $\varepsilon_i - \varepsilon_j$, where $1 \leq i < j \leq N$. Moreover, if $\alpha \in \Delta$ is in the support of γ and we represent α as $\varepsilon_k - \varepsilon_{k+1}$ for a fixed k, we may choose i and j so that $i \leq k < j$. This can be proven in general but is easily checked for the cases at hand. Note that to simplify the exposition that follows we will treat $\varepsilon_i - \varepsilon_j$ as if it were always a root; any assertion regarding $\varepsilon_i - \varepsilon_j$ should be ignored if it is not a root. For $w \in {}_{\mathbb{Q}}W$ we will also denote by w the corresponding permutation of $\{1, 2, \dots, N\}$; thus $w(\varepsilon_i - \varepsilon_j) = \varepsilon_{w(i)} - \varepsilon_{w(j)}$.

---

[10] Our notation differs in several ways from [35]. For one thing, the meaning of ${}_{\mathbb{Q}}W_P$ and ${}_{\mathbb{Q}}W^P$ are reversed. Furthermore, since we consider right cosets of ${}_{\mathbb{Q}}W^P$ rather than left cosets, we have w^{-1} where [35] would have w.

First assume that $w(1) > 1$. Then

$$\begin{aligned} \varepsilon_i - \varepsilon_{w(1)} &\in {}_{\mathbb{Q}}\Phi_w \quad \text{for } i < w(1), \\ \varepsilon_{w(1)} - \varepsilon_k &\notin {}_{\mathbb{Q}}\Phi_w \quad \text{for } w(1) < k. \end{aligned} \tag{19.4.1}$$

Since $w \in {}_{\mathbb{Q}}W_P$ it follows that $\varepsilon_{w(1)-1} - \varepsilon_{w(1)} \notin \Delta^P$. If $\varepsilon_{w(1)} - \varepsilon_{w(1)+1}, \dots, \varepsilon_{N-1} - \varepsilon_N \in \Delta^P$, then $\mathrm{pr}_P(\varepsilon_i - \varepsilon_j) = \mathrm{pr}_P(\varepsilon_i - \varepsilon_{w(1)})$ for all $i < w(1) \leq j$ and so by Lemma 19.3 and (19.4.1) such $\varepsilon_i - \varepsilon_j$ must belong to ${}_{\mathbb{Q}}\Phi_w$. We can thus let T have type $\Delta \setminus \{\varepsilon_{w(1)-1} - \varepsilon_{w(1)}\}$ and Case 1 is satisfied. Otherwise let $b \geq w(1)$ be the least index such that $\varepsilon_b - \varepsilon_{b+1} \notin \Delta^P$. Then Lemma 19.3 and (19.4.1) imply that

$$\begin{aligned} \varepsilon_i - \varepsilon_j &\in {}_{\mathbb{Q}}\Phi_w \quad \text{for } i < w(1) \leq j \leq b, \\ \varepsilon_j - \varepsilon_k &\notin {}_{\mathbb{Q}}\Phi_w \quad \text{for } w(1) \leq j \leq b < k. \end{aligned}$$

Thus we can let T have type $\Delta \setminus \{\varepsilon_{w(1)-1} - \varepsilon_{w(1)}, \varepsilon_b - \varepsilon_{b+1}\}$ and Case 2 is satisfied.

Now assume that $w(1) = 1$. If $w(i) = i$ for all i then ${}_{\mathbb{Q}}\Phi_w = \emptyset$ and letting T have type $\Delta \setminus \{\varepsilon_i - \varepsilon_{i+1}\}$ for any $\varepsilon_i - \varepsilon_{i+1} \notin \Delta^P$ will do. Otherwise let $a > 1$ be the least integer such that $w(a) > a$. Since $\varepsilon_a - \varepsilon_{w(a)} \in {}_{\mathbb{Q}}\Phi_w$ whereas $\varepsilon_{a-1} - \varepsilon_{w(a)} \notin {}_{\mathbb{Q}}\Phi_w$ we must have $\varepsilon_{a-1} - \varepsilon_a \notin \Delta^P$ by Lemma 19.3. Furthermore $\varepsilon_i - \varepsilon_j \notin {}_{\mathbb{Q}}\Phi_w$ for $i < a \leq j$ so we can let T have type $\Delta \setminus \{\varepsilon_{a-1} - \varepsilon_a\}$. □

19.5. **Proof of Lemma 17.8.** We return to the setting of Lemma 17.8; thus w is now an element of W. Let $\lambda \in \mathfrak{h}_{\mathbb{C}}$ be the highest weight of E; as in the proof of Lemma 17.6 the hypotheses on V and Q imply that λ satisfies

$$\begin{aligned} \langle w(\lambda + \rho), \alpha^\vee \rangle &\leq 0 \qquad \text{for } \alpha \in \Phi(\mathfrak{n}_P^Q; \mathfrak{a}_P), \\ \langle w(\lambda + \rho), \alpha^\vee \rangle &\geq 0 \qquad \text{for } \alpha \in \Phi(\mathfrak{n}_P^{P,Q}; \mathfrak{a}_P). \end{aligned} \tag{19.5.1}$$

and

$$\lambda \in B = \{\, \lambda \text{ dominant} \mid w(\lambda + \rho)|_{\mathfrak{b}_{P,-1}} = 0 \,\}$$

(see (8.2.3) and (8.3.1)). The transformation $\lambda \mapsto c\lambda + (c-1)\rho$ for $c \geq 1$ preserves these conditions (since it corresponds to rescaling $w(\lambda + \rho)$ by c) so we may assume that λ is strictly dominant and hence in the interior of B. By perturbing λ within B slightly we can arrange that all the inequalities in (19.5.1) are strict inequalities and that in addition that $\langle w(\lambda + \rho), \alpha^\vee \rangle \neq 0$ for all $\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)$.

Since we have arranged that $w(\lambda+\rho)|_{\mathfrak{a}_P}$ lies in a chamber of $\mathfrak{a}_P$, the discussion of §19.2 and in particular (19.2.2) shows that it determines an element ${}_{\mathbb{Q}}w \in {}_{\mathbb{Q}}W(P)$.

By (19.5.1) and Lemma 19.3 we have
(19.5.2)

$$\begin{aligned} \Phi_{\mathbb{Q}w} \cap \operatorname{pr}_P^{-1}(\alpha) = \operatorname{pr}_P^{-1}(\alpha) &\iff \langle w(\lambda+\rho), \alpha^\vee \rangle < 0 &&\Longleftarrow \quad \alpha \in \Phi(\mathfrak{n}_P^Q; \mathfrak{a}_P), \\ \Phi_{\mathbb{Q}w} \cap \operatorname{pr}_P^{-1}(\alpha) = \emptyset &\iff \langle w(\lambda+\rho), \alpha^\vee \rangle > 0 &&\Longleftarrow \quad \alpha \in \Phi(\mathfrak{n}_P^{P,Q}; \mathfrak{a}_P). \end{aligned}$$

for all $\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)$. Apply Lemma 19.4. If $\#\Delta_T = 1$, denote the unique element of Δ_T by β_- or β_+ depending on whether $\Phi_w \cap \operatorname{pr}_T^{-1}(\Phi_T^+) = \operatorname{pr}_P^{-1}(\Phi_T^+)$ or $\emptyset$. In general let $\alpha_\pm \in \Delta_P$ restrict to $\beta_\pm$. The lemma and (19.5.2) imply that $\alpha_- \in \Delta_P^Q$ and $\alpha_+ \in \Delta_P \setminus \Delta_P^Q$ (which establishes (17.8.1)) and furthermore that for any $\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)$,

$$\begin{aligned} &\langle w(\lambda+\rho), \alpha^\vee \rangle < 0 \qquad \text{if } \alpha|_{\mathfrak{a}_T} \in \mathbb{N}\beta_-, \\ &\langle w(\lambda+\rho), \alpha^\vee \rangle > 0 \qquad \text{if } \alpha|_{\mathfrak{a}_T} \in \mathbb{N}\beta_+. \end{aligned} \tag{19.5.3}$$

Lemma 17.6 concludes the proof of (17.8.2).

The final assertion may be proved by induction. Suppose that $P < Q < G$, $\Delta_T = 1$ and $\ell(w_T) = \frac{1}{2}\dim \mathfrak{n}_T$. Then $\ell_\alpha(w) = \frac{1}{2}\dim \mathfrak{n}_\alpha$ for all $\alpha \in \Phi(\mathfrak{n}_T, \mathfrak{a}_P)$. If $P = Q \cap T$, then $\Delta_P^T = \Delta_P \setminus \Delta_P^Q$ and so $\ell_\alpha(w) \leq \frac{1}{2}\dim \mathfrak{n}_\alpha$ for all $\alpha \in \Phi(\mathfrak{n}_P^T, \mathfrak{a}_P)$ by Lemma 17.6. Then $\widetilde{T}$ with $\Delta_P^{\widetilde{T}} = \Delta_P^T \setminus \{\alpha\}$ for any $\alpha \in \Delta_P^T$ has the desired properties. An analogous argument applies if $Q \cap T = T$. Otherwise we can apply the lemma by induction to the parabolic $\mathbb{Q}$-subgroups $P/N_T < (Q \cap T)/N_T < T/N_T$ with w and λ replaced by w^T and $w_T(\lambda+\rho)-\rho$ respectively to obtain a parabolic $\mathbb{Q}$-subgroup $\widetilde{T}/N_T$. Then $\widetilde{T}$ has the desired properties. □

20. Proof of Theorem 17.9

If $T = P$ and $Q = P$ or G we can apply Proposition 17.7 to obtain the desired vanishing; the hypothesis (17.6.1) follows from (17.8.2) and the nonvanishing cases are excluded by (17.8.3). So in this section we can assume that if $T = P$, then $P < Q < G$; this is necessary for Lemma 20.6 below.

20.1. **Notation.** In order to simplify the notation, we will simply write $IH(U)$ for $I_{p_w}H(U)$. Furthermore, if $Z \subseteq |\Delta_P|$ is a locally closed constructible subset, we write

$$IH_Z = IH_Z(U)$$

where $U \subseteq |\Delta_P|$ is any open constructible subset containing Z as a relatively closed subset. This is well-defined by excision.

20.2. Consider the commutative diagram

$$\begin{tikzcd} \cdots \arrow[r] & IH^i_{c(|\Delta_P^Q|)}(c(|\Delta_P|)) \arrow[r] & IH^i(c(|\Delta_P|)) \arrow[r, "r"] \arrow[d, "s"] & IH^i(c(|\Delta_P|) \setminus c(|\Delta_P^Q|)) \arrow[r] \arrow[d, "\cong"] & \cdots \\ \cdots \arrow[r] & IH^i_{|\Delta_P^Q|} \arrow[r] & IH^i(|\Delta_P|) \arrow[r] & IH^i(|\Delta_P| \setminus |\Delta_P^Q|) \arrow[r] & \cdots \end{tikzcd}$$

in which both rows are long exact sequences. The vanishing assertion of Theorem 17.9 is equivalent to r being an isomorphism. However by the local characterization of intersection cohomology, the map s is an isomorphism in degrees

$i \le p_w(P)$ and the inclusion of 0 in degrees $i > p_w(P)$. Thus we need to show that

$$
\begin{aligned}
&\text{(20.2.1a)} \quad IH^{i-1}(|\Delta_P| \setminus |\Delta_P^Q|) = 0 && \text{for } i > p_w(P)+1,\\
&\text{(20.2.1b)} \quad \operatorname{Im}\Big(IH^{p_w(P)}(|\Delta_P| \setminus |\Delta_P^Q|) \longrightarrow IH^{p_w(P)+1}_{|\Delta_P^Q|} \Big) = 0 && \text{for } i = p_w(P)+1,\\
&\text{(20.2.1c)} \quad IH^i_{|\Delta_P^Q|} = 0 && \text{for } i < p_w(P)+1.
\end{aligned}
$$

To prove (20.2.1a) we will use the long exact sequence
(20.2.2)

$$
\dots \longrightarrow IH^{i-1}_{|\Delta_P^{Q\vee T}| \setminus |\Delta_P^Q|} \longrightarrow IH^{i-1}(|\Delta_P| \setminus |\Delta_P^Q|) \longrightarrow IH^{i-1}(|\Delta_P| \setminus |\Delta_P^{Q\vee T}|) \longrightarrow \dots
$$

and show the outside terms vanish in the appropriate degrees. Equation (20.2.1c) will follow similarly using

$$
\text{(20.2.3)} \qquad \dots \longrightarrow IH^i_{|\Delta_P^Q \setminus \Delta_P^{Q\cap T}|} \longrightarrow IH^i_{|\Delta_P^Q|} \longrightarrow IH^i_{|\Delta_P^Q| \setminus |\Delta_P^Q \setminus \Delta_P^{Q\cap T}|} \longrightarrow \dots \ .
$$

For (20.2.1b) we will use these arguments to show that either $IH^{p_w(P)}(|\Delta_P| \setminus |\Delta_P^Q|) = 0$ or $IH^{p_w(P)+1}_{|\Delta_P^Q|} = 0$.

20.3. Lemma. $IH^{i-1}_{|\Delta_P^{Q\vee T}| \setminus |\Delta_P^Q|} = 0$ *for all* i.

Proof. For $R > P$, let U_R be the star neighborhood of the open face $|\Delta_P^R|^\circ$. Decompose

$$
|\Delta_P^{Q\vee T}| \setminus |\Delta_P^Q| = \coprod_{\emptyset \ne \Delta_P^R \subseteq \Delta_P^T \setminus \Delta_P^{Q\cap T}} U_R \cap |\Delta_P^{Q\vee R}|.
$$

The corresponding Fary spectral sequence abutting to $IH^{i-1}_{|\Delta_P^{Q\vee T}| \setminus |\Delta_P^Q|}$ has (compare Lemma 3.7)

$$
E_1^{-p,i-1+p} = \bigoplus_{\substack{\emptyset \ne \Delta_P^R \subseteq \Delta_P^T \setminus \Delta_P^{Q\cap T} \\ \#\Delta_P^R = p}} IH^{i-1}_{U_R \cap |\Delta_P^{Q\vee R}|} \cong \bigoplus_{\substack{\emptyset \ne \Delta_P^R \subseteq \Delta_P^T \setminus \Delta_P^{Q\cap T} \\ \#\Delta_P^R = p}} IH^{i-1}_{c(|\Delta_R^{Q\vee R}|)}(c(|\Delta_R|)) \ .
$$

Theorem 17.9 may be applied by induction (with $P \le Q$ replaced by $R \le Q \vee R$ and T remaining the same) to prove that this vanishes. □

Similarly we have the

20.4. Lemma. $IH^i_{|\Delta_P^Q| \setminus |\Delta_P^Q \setminus \Delta_P^{Q\cap T}|} = 0$ *for all* i.

Proof. Cover $|\Delta_P^Q| \setminus |\Delta_P^Q \setminus \Delta_P^{Q\cap T}|$ by $\{U_\alpha\}_{\alpha \in \Delta_P^{Q\cap T}}$, where U_α is the open star of the vertex α. The corresponding Mayer-Vietoris spectral sequence abutting to $IH^i_{|\Delta_P^Q| \setminus |\Delta_P^Q \setminus \Delta_P^{Q\cap T}|}$ has (compare Lemma 3.7)

$$
E_1^{p,i-p} = \bigoplus_{\substack{\emptyset \ne \Delta_P^R \subseteq \Delta_P^{Q\cap T} \\ \#\Delta_P^R = p+1}} IH^{i-p}_{U_R \cap |\Delta_P^Q|} \cong \bigoplus_{\substack{\emptyset \ne \Delta_P^R \subseteq \Delta_P^{Q\cap T} \\ \#\Delta_P^R = p+1}} IH^{i-p}_{c(|\Delta_R^Q|)}(c(|\Delta_R|)) \ .
$$

Theorem 17.9 may be applied by induction (with $P \le Q$ replaced by $R \le Q$ and T remaining the same) to prove that this vanishes. □

20.5. In view of the preceding two lemmas, Lemma 20.6 below will conclude the proof of the theorem. First we need a useful formula. For a perversity p let

$$dp(k) = p(k+1) - p(k) \qquad \text{and} \qquad \delta(k) = \begin{cases} 1 & \text{if } k \text{ odd}, \\ 0 & \text{if } k \text{ even}, \end{cases}$$

Thus for the middle perversities we have $dm = \delta$ and $dn = 1 - \delta$.

Lemma. *For p a middle perversity, $w \in W$, and $P \leq R$,*

$$\begin{aligned} p_w(P) = p_w(R) &+ \left(\tfrac{1}{2}(\dim \mathfrak{n}_P^R + \#\Delta_P^R) - \ell(w_P^R)\right) \\ &+ \delta(\dim \mathfrak{n}_P^R + \#\Delta_P^R)\left(\tfrac{1}{2} - dp(\dim \mathfrak{n}_P + \#\Delta_P)\right). \end{aligned}$$

Proof. This is a simple verification using $\ell(w_P) = \ell(w_P^R) + \ell(w_R)$ and the definition of a middle perversity. $\square$

20.6**. Lemma.** *In addition to the hypotheses of Theorem* 17.9*, assume that if $T = P$, then $P < Q < G$. Then the following vanishing results hold*:

(a) $IH^{i-1}(|\Delta_P| \setminus |\Delta_P^{Q\vee T}|) = 0$ *for* $\begin{cases} i > p_w(P) + 1, \textit{ or} \\ i = p_w(P) + 1 \textit{ and} \\ \qquad dp(\dim \mathfrak{n}_P + \#\Delta_P) = 0. \end{cases}$

(c) $IH^i_{|\Delta_P^Q \setminus \Delta_P^{Q\cap T}|} = 0$ *for* $\begin{cases} i = p_w(P) + 1 \textit{ and } dp(\dim \mathfrak{n}_P + \#\Delta_P) = 1, \textit{ or} \\ i < p_w(P) + 1. \end{cases}$

Proof. Consider part (a). Either $\Delta_P \setminus \Delta_P^{Q\vee T}$ is empty (in which case the lemma is trivial) or by (17.8.1) it has just one element. Let $R = (P, Q \vee T)$ be the corresponding parabolic subgroup with $\#\Delta_P^R = 1$. We know that $R \neq G$ since otherwise $Q \vee T = P$ and hence $Q = T = P$. Thus $IH^{i-1}(|\Delta_P| \setminus |\Delta_P^{Q\vee T}|) \cong IH^{i-1}(c(|\Delta_R|))$ will be zero due to truncation at the cone point if $i - 1 > p_w(R)$. By Lemma 20.5 this can be re-expressed as

$$\begin{aligned} \left[i - (p_w(P) + 1)\right] &+ \left[\tfrac{1}{2}(\dim \mathfrak{n}_P^R + 1) - \ell(w_P^R)\right] \\ &+ \left[\delta(\dim \mathfrak{n}_P^R + 1)\left(\tfrac{1}{2} - dp(\dim \mathfrak{n}_P + \#\Delta_P)\right)\right] > 0. \end{aligned} \tag{20.6.1}$$

Since $\ell(w_P^R) \leq \frac{1}{2}\dim \mathfrak{n}_P^R$ by (17.8.2), the second bracketed term of (20.6.1) is at least $\frac{1}{2}$. We are considering $i \geq p_w(P) + 1$ so the first term is nonnegative (and at least 1 if $i > p_w(P) + 1$), while the third term is at least $-\frac{1}{2}$ (and nonnegative if $dp(\dim \mathfrak{n}_P + \#\Delta_P) = 0$). This proves part (a).

Part (c) is similar. Let R have type $\Delta_P^R = \Delta_P^Q \setminus \Delta_P^{Q\cap T} = \Delta_P^{Q\vee T} \setminus \Delta_P^T$. Again $R \neq G$ since otherwise $Q = G$ and $T = P$. Then $IH^i_{|\Delta_P^Q \setminus \Delta_P^{Q\cap T}|} \cong IH^i_{c(\emptyset)}(c(|\Delta_R|))$ will be zero if $i - 1 \leq p_w(R)$. The application of Lemma 20.5 and the inequality $\ell(w_P^R) \geq \frac{1}{2}\dim \mathfrak{n}_P^R$ from (17.8.2) concludes the proof. $\square$

Part IV. Satake Compactifications and Functoriality of Micro-support

This part of the paper begins with two independent sections. In §21 we introduce Satake compactifications X_σ^* of X and recall Zucker's result [57] that there is a natural quotient map $\pi\colon \widehat{X} \to X_\sigma^*$. In §22 we discuss in a fairly general context the functorial behavior of micro-support of an $\mathcal{L}_W$-module $\mathcal{M}$ under k^* and $k^!$, where $k\colon Z \hookrightarrow W$ is an inclusion of admissible spaces. Following this we proceed to our main result, Theorem 26.1, which is a significant strengthening of this functoriality in the case of restriction to the fibers of $\pi\colon \widehat{X} \to X_\sigma^*$.

21. Satake Compactifications

We will briefly outline the theory of Satake compactifications X_σ^* and give Zucker's realization [57] of X_σ^* as a quotient space of $\widehat{X}$. References are [49], [50], [5], [57], and [17].

21.1. **Notation.** Let $S \subseteq {}_{\mathbb{R}}S$ be maximal tori of G split over $\mathbb{Q}$ and $\mathbb{R}$ respectively and choose compatible orderings on the roots. For $k = \mathbb{Q}$ or $\mathbb{R}$ let ${}_k\Delta$ denote the simple restricted k-roots; as usual we omit the left subscript when $k = \mathbb{Q}$. We will be working with a regular representation $\sigma\colon G \to GL(U)$. Let μ denote highest weight of σ and let ${}_k\mu = \mu|_{{}_k\mathfrak{a}}$ be the restricted highest k-weight. A subset $\Theta \subseteq {}_k\Delta$ is said to be *σ-connected* if $\Theta \cup \{{}_k\mu\}$ is connected. Here a finite subset Θ of an inner product space is said to be *connected* if the associated graph (with vertices Θ and an edge between α and β whenever $(\alpha, \beta) \neq 0$) is connected. For any subset $\Theta \subseteq {}_k\Delta$, let $\kappa(\Theta)$ be the largest σ-connected subset of Θ. If Θ is a σ-connected subset of ${}_k\Delta$, let $\omega(\Theta)$ be the largest subset of ${}_k\Delta$ such that $\kappa(\omega(\Theta)) = \Theta$. As in [17] we extend this notation to arbitrary subsets $\Theta \subseteq {}_k\Delta$ by setting $\omega(\Theta) = \omega(\kappa(\Theta))$; it is easy to see that $\omega(\Theta) \supseteq \Theta$.

If P is a parabolic $\mathbb{R}$-subgroup, let $P^\dagger$ be the parabolic $\mathbb{R}$-subgroup containing P with $\mathbb{R}$-type $\omega({}_{\mathbb{R}}\Delta^P)$. This is the unique largest parabolic $\mathbb{R}$-subgroup containing P with $\kappa({}_{\mathbb{R}}\Delta^{P^\dagger}) = \kappa({}_{\mathbb{R}}\Delta^P)$. In general a parabolic $\mathbb{R}$-subgroup will be called *σ-saturated* if it has $\mathbb{R}$-type $\omega(\Theta)$ for some $\Theta \subseteq {}_{\mathbb{R}}\Delta$. Let $\rho\colon {}_{\mathbb{R}}\Delta \to \Delta \cup \{0\}$ denote[11] the restriction map on simple roots. For a subset $\Theta \subseteq {}_{\mathbb{R}}\Delta$ set $\Theta^{\wedge} = \rho(\Theta) \setminus \{0\}$ and for a subset $\Upsilon \subseteq \Delta$ set $\widetilde{\Upsilon} = \rho^{-1}(\Upsilon \cup \{0\})$. Subsets of ${}_{\mathbb{R}}\Delta$ of the form $\widetilde{\Upsilon}$ are called *$\mathbb{Q}$-rational* and are precisely the $\mathbb{R}$-types of the parabolic $\mathbb{Q}$-subgroups. Clearly $(\widetilde{\Upsilon})^{\wedge} = \Upsilon$ and $\widetilde{(\Theta^{\wedge})}$ is the smallest $\mathbb{Q}$-rational set containing Θ.

21.2. **Satake compactifications of D.** In this subsection and the next we depart from our usual context and assume that G is merely defined over $\mathbb{R}$ with $K \subseteq G(\mathbb{R})$ a maximal compact subgroup and that D denotes the associated symmetric space $G(\mathbb{R})/K_{\mathbb{R}}A_G$.

Let $\sigma\colon G \to \mathrm{GL}(U)$ be an irreducible regular representation of G which is nontrivial on every simple factor of $G(\mathbb{R})$ and fix an admissible metric h_0 on U with respect to K as in §12.1. Let T^* denote the adjoint with respect to h_0 of an endomorphism T of U and let $S(U)$ denote the real vector space of self-adjoint endomorphisms. The representation σ induces a natural representation $\tilde{\sigma}\colon G(\mathbb{R}) \to \mathrm{GL}(\mathbb{P}S(U))$ given by

$$\tilde{\sigma}(g)[T] = [\sigma(g) \circ T \circ \sigma(g)^*] \ .$$

The class of the identity endomorphism is fixed by $K_{\mathbb{R}}A_G$ under $\tilde{\sigma}$ and thus the map $g \mapsto [\sigma(g)\sigma(g)^*]$ descends a $G(\mathbb{R})$-equivariant map

$$D \longrightarrow \mathbb{P}S(U) \ .$$

---

[11] Our use in this section of this standard notation should not be confused with our equally standard use elsewhere of ρ to denote one-half the sum of the positive roots.

This map is an embedding; the closure of the image is denoted ${}_{\mathbb{R}}D^*_\sigma$, the *Satake compactification*[12] *of D associated to σ.* The Satake compactification is independent of the choice of K and h_0; usually σ will be fixed and we omit it from the notation. If D is Hermitian symmetric then the closure of the Harish-Chandra embedding of D as a bounded domain is called the *natural compactification* and it is topologically equivalent to the Satake compactification for a certain σ.

21.3. **Real boundary components.** A *real Satake boundary component* $D_{P,h} \subseteq {}_{\mathbb{R}}D^*$ is the set of points fixed under the action of $N_P(\mathbb{R})$ for some parabolic $\mathbb{R}$-subgroup P. The action of $P(\mathbb{R})$ descends to an action of $L_P(\mathbb{R})$ on $D_{P,h}$. The centralizer of $D_{P,h}$ under this action is the group of real points of a normal reductive $\mathbb{R}$-subgroup $L'_{P,\ell} \subseteq L_P$. The action then descends further to the real points of $L'_{P,h} = L_P/L'_{P,\ell}$, and thus $D_{P,h}$ is realized as $L'_{P,h}(\mathbb{R})/(K_P \cap L'_{P,h}(\mathbb{R}))$, the symmetric space associated to $L'_{P,h}(\mathbb{R})$. (We use the subscripts h and ℓ to mimic the notation frequently used in the Hermitian case.)

We can describe these groups concretely in terms of roots. Set $\mathit{\Theta} = \kappa({}_{\mathbb{R}}\Delta^P)$; then the orthogonal decomposition ${}_{\mathbb{R}}\Delta^P = \mathit{\Theta} \coprod ({}_{\mathbb{R}}\Delta^P \setminus \mathit{\Theta})$ yields a decomposition of the Levi factor as an almost direct product,

$$L_P = \widetilde{L'_{P,h}} L'_{P,\ell}. \tag{21.3.1}$$

Here $\widetilde{L'_{P,h}}$ is the minimal normal connected semisimple $\mathbb{R}$-subgroup with $\mathit{\Theta}$ as simple $\mathbb{R}$-roots; it is a lift of $L'_{P,h}$. The group $L'_{P,\ell}$ is the almost direct product of the minimal normal connected semisimple $\mathbb{R}$-subgroup with ${}_{\mathbb{R}}\Delta^P \setminus \mathit{\Theta}$ as simple $\mathbb{R}$-roots together with the center of L_P and all almost simple factors of L_P with compact groups of real points [12, 5.11].

The parabolic $\mathbb{R}$-subgroup P giving rise to a boundary component $D_{P,h}$ as above is not unique. To index $D_{P,h}$ in a unique fashion we use its normalizer $\{ g \in G(\mathbb{R}) \mid gD_{P,h} = D_{P,h} \}$; this is the group of real points of a σ-saturated parabolic $\mathbb{R}$-subgroup, namely $P^\dagger$. Recall that $P^\dagger$ is the unique parabolic $\mathbb{R}$-subgroup containing P with $\mathbb{R}$-type $\omega({}_{\mathbb{R}}\Delta^P)$. For later use we note that the centralizer $\{ g \in G(\mathbb{R}) \mid gy = y \text{ for all } y \in D_{P,h} \}$ of $D_{P,h}$ is the group of real points of the inverse image of $L'_{P^\dagger,\ell}$ under $P^\dagger \to L_{P^\dagger}$. On the other hand, if R is a σ-saturated parabolic $\mathbb{R}$-subgroup, $R(\mathbb{R})$ is the normalizer of a unique boundary component $D_{R,h}$, namely the points fixed by $N_R(\mathbb{R})$. We have

$${}_{\mathbb{R}}D^*_\sigma = \coprod_{\substack{R\ \sigma\text{-saturated} \\ \text{parabolic } \mathbb{R}\text{-subgroup}}} D_{R,h} \ .$$

21.4. **Rational boundary components.** Now consider again a connected reductive algebraic $\mathbb{Q}$-group G with $D = G(\mathbb{R})/KA_G$. We assume that ${}_{\mathbb{R}}S_G = S_G$. (This is not necessary but simplifies the exposition; the assumption is satisfied if

---

[12] Here we have followed most closely Zucker [57]; Borel [5] uses an action on the right for U and $S(U)$ and hence replaces μ by the lowest weight of U. Satake's construction [49] is identical except that he represents $S(U)$ by Hermitian matrices. Casselman [17] dispenses with U altogether and works directly with an irreducible representation V containing a K-fixed vector. In [44, §3] it is shown that given U there exists V (in fact contained in $S(U)_{\mathbb{C}}$) such that Casselman's construction agrees with Satake's construction. The highest weight of this V is $2_{\mathbb{R}}\mu$ and hence the notion of σ-connected defined here agrees with that in [17] for subsets of Δ or ${}_{\mathbb{R}}\Delta$.

D is Hermitian or ${}^0G$ is equal-rank.) Then the preceding two subsections may be applied to yield a Satake compactification ${}_{\mathbb{R}}D^*_\sigma$.

A real boundary component $D_{R,h}$ of ${}_{\mathbb{R}}D^*_\sigma$ is called *rational* [2, §§3.5, 3.6] if

(i) its normalizer R is defined over $\mathbb{Q}$, and
(ii) the group $L'_{R,\ell}$ contains a normal subgroup $L_{R,\ell}$ of L_R defined over $\mathbb{Q}$ such that $L'_{R,\ell}(\mathbb{R})/L_{R,\ell}(\mathbb{R})$ is compact.[13]

Here Condition (i) is equivalent to $\Gamma_{N_R}\backslash N_R(\mathbb{R})$ being compact while Condition (ii) ensures that $D_{R,h}$ may be realized as the symmetric space associated to the reductive algebraic $\mathbb{Q}$-group $L_{R,h} \equiv L_R/L_{R,\ell}$. For a rational boundary component $D_{R,h}$, equation (21.3.1) may be replaced by a factorization by $\mathbb{Q}$-subgroups,

$$L_R = \widetilde{L_{R,h}}L_{R,\ell}. \tag{21.4.1}$$

On the other hand, consider the real boundary components $D_{R,h}$ for which

$${}_{\mathbb{R}}\Delta^R = \omega(\widetilde{\varUpsilon}) \qquad \text{for some } \sigma\text{-connected } \varUpsilon \subseteq \Delta. \tag{21.4.2}$$

Real boundary components satisfying (21.4.2) are exactly those in the closure of a Siegel set [17, Lemma 8.1]. By [57, Proposition 3.3(i)] any real boundary component satisfying Condition (i) satisfies (21.4.2). The Satake compactification ${}_{\mathbb{R}}D^*_\sigma$ is called *geometrically rational* if every real boundary component satisfying (21.4.2) is rational.[14] Examples of geometrically rational Satake compactifications include those where σ is defined over $\mathbb{Q}$ [44, Theorem 8] or where D is a Hermitian symmetric space and ${}_{\mathbb{R}}D^*_\sigma$ is the natural compactification [2]. Casselman [17] gives a necessary and sufficient criterion for geometric rationality in terms of ${}_{\mathbb{R}}\mu$ and the Tits index of G.

For geometrically rational Satake compactifications, Condition (i) in the definition of rational boundary component implies Condition (ii). Thus the rational boundary components of a geometrically rational Satake compactification are indexed by the σ-saturated parabolic $\mathbb{Q}$-subgroups. In fact by the following lemma these are precisely the parabolic $\mathbb{Q}$-subgroups with $\mathbb{Q}$-type $\omega(\varUpsilon)$ for $\varUpsilon \subseteq \Delta$.

Lemma. *If ${}_{\mathbb{R}}D^*_\sigma$ is geometrically rational,*

$$\widetilde{\omega(\varUpsilon)} = \omega(\widetilde{\varUpsilon}) \qquad \textit{for } \varUpsilon \subseteq \Delta, \tag{21.4.3}$$

$$\omega(\varTheta)^{\wedge} = \omega(\varTheta^{\wedge}) \qquad \textit{for } \varTheta \subseteq {}_{\mathbb{R}}\Delta \textit{ with } \omega(\varTheta)\ \mathbb{Q}\textit{-rational.} \tag{21.4.4}$$

Proof. Note that a subset $\varUpsilon \subseteq \Delta$ is σ-connected if and only if there is a σ-connected subset $\varTheta \subseteq {}_{\mathbb{R}}\Delta$ with $\varTheta^{\wedge} = \varUpsilon$ [57, (2.4)] and that this implies that

$$\kappa(\varTheta^{\wedge}) = \kappa(\varTheta)^{\wedge} \qquad \text{for any } \varTheta \subseteq {}_{\mathbb{R}}\Delta\ . \tag{21.4.5}$$

To prove (21.4.3) we first claim that $\kappa(\widetilde{\omega(\varUpsilon)}) = \kappa(\widetilde{\varUpsilon})$. Since $\widetilde{\omega(\varUpsilon)}$ contains $\kappa(\widetilde{\varUpsilon})$ it follows that $\kappa(\widetilde{\omega(\varUpsilon)}) \supseteq \kappa(\widetilde{\varUpsilon})$. On the other hand, $\kappa(\widetilde{\omega(\varUpsilon)})^{\wedge}$ is σ-connected and contained in $\omega(\varUpsilon)$ and therefore $\kappa(\widetilde{\omega(\varUpsilon)})^{\wedge} \subseteq \kappa(\varUpsilon) \subseteq \varUpsilon$; it follows that $\kappa(\widetilde{\omega(\varUpsilon)}) \subseteq$

---

[13] In [5] it is assumed that $L'_{R,\ell}$ itself is defined over $\mathbb{Q}$, but it is noted in [2] that this is too restrictive.

[14] Our presentation here follows Casselman [17]. The assumption that Condition (i) holds for the real boundary components satisfying (21.4.2) is the assertion that $\omega(\widetilde{\varUpsilon})$ is $\mathbb{Q}$-rational for any σ-connected subset $\varUpsilon \subseteq \Delta$. A weaker assumption is made in [57, (3.3)] (Assumption 1) and the above assertion is deduced via [57, Proposition 3.3(ii)]. However as pointed out in [17, §9], this proposition is incorrect.

$\kappa(\widetilde{\Upsilon})$ which finishes the proof of the claim. The claim implies that $\widetilde{\omega(\Upsilon)} \subseteq \omega(\widetilde{\Upsilon})$. For the opposite inclusion it suffices (since by geometric rationality both sets are $\mathbb{Q}$-rational) to check that $\omega(\Upsilon) \supseteq \omega(\widetilde{\Upsilon})^{\wedge}$. This follows since $\kappa(\Upsilon) = \kappa(\widetilde{\Upsilon})^{\wedge} = \kappa(\omega(\widetilde{\Upsilon})^{\wedge})$ by (21.4.5). For (21.4.4) suppose $\omega(\Theta)$ is $\mathbb{Q}$-rational for $\Theta \subseteq {}_{\mathbb{R}}\Delta$ and set $\Upsilon = \kappa(\Theta)^{\wedge}$. By [57, Proposition 3.3(i)], $\kappa(\Theta) = \kappa(\widetilde{\Upsilon})$ and hence $\omega(\Theta) = \omega(\widetilde{\Upsilon}) = \widetilde{\omega(\Upsilon)}$ (use (21.4.3) for the last equality); we thus see that $\omega(\Theta)^{\wedge} = \omega(\Upsilon)$ which in view of (21.4.5) completes the proof. $\square$

21.5. **Satake compactifications of X.** Assume ${}_{\mathbb{R}}D^*_\sigma$ is a geometrically rational Satake compactification of D. Let D^*_σ be the union of D and the rational boundary components. By the preceding discussion we have

$$D^*_\sigma = \coprod_{\substack{R\ \sigma\text{-saturated} \\ \text{parabolic } \mathbb{Q}\text{-subgroup}}} D_{R,h}$$

and $G(\mathbb{Q})$ acts on D^*_σ. There is a topology on D^*_σ (the "Satake topology") for which $X^*_\sigma \equiv \Gamma\backslash D^*_\sigma$ is a compact Hausdorff space containing X as a dense open subset; X^*_σ is the *Satake compactification of X associated to σ* and as usual we will omit σ from the notation. For example, if D is a Hermitian symmetric space and ${}_{\mathbb{R}}D^*_\sigma$ is the natural compactification, then X^* is the Baily-Borel-Satake compactification [2].

Let $\mathcal{P}^*_\sigma$ denote the Γ-conjugacy classes of σ-saturated parabolic $\mathbb{Q}$-subgroups. For $R \in \mathcal{P}^*_\sigma$, set $F_R = \Gamma_{L_{R,h}}\backslash D_{R,h}$, where we view $D_{R,h}$ as the symmetric space associated to $L_{R,h}$ as above and $\Gamma_{L_{R,h}} = \Gamma_{L_R}/(\Gamma_{L_R} \cap L_{R,\ell})$. Then X^*_σ has a stratification

$$X^*_\sigma = \coprod_{R \in \mathcal{P}^*_\sigma} F_R \ .$$

We define a partial order on the rational boundary components (and similarly on the strata of X^*_σ) by setting $D_{R,h} \leq D_{R',h}$ if and only if $D_{R,h} \subseteq \operatorname{cl}(D_{R',h})$; this corresponds to the conditions that $R^\gamma \cap R'$ is a parabolic $\mathbb{Q}$-subgroup (for some $\gamma \in \Gamma$ and $\kappa(\Delta^R) \subseteq \kappa(\Delta^{R'})$.

21.6. **X^*_σ as a quotient of $\widehat{X}$.** Continue to assume that ${}_{\mathbb{R}}D^*_\sigma$ is a geometrically rational Satake compactification of D. Let P be a parabolic $\mathbb{Q}$-subgroup and let ${}_{\mathbb{R}}\Delta^P = \widetilde{\Upsilon}$ be its $\mathbb{R}$-type. By (21.4.3), $\omega(\widetilde{\Upsilon})$ is $\mathbb{Q}$-rational so $P^\dagger$ is a σ-saturated parabolic $\mathbb{Q}$-subgroup and thus defines a stratum $D_{P^\dagger,h}$ of D^*. Let $D_{P,\ell} = L_{Q,\ell}(\mathbb{R})/K_{Q,\ell}A_Q$ be the symmetric space of the second factor of (21.4.1) (where $K_{Q,\ell} = K_Q \cap L_{Q,\ell}(\mathbb{R})$). The factorization (21.4.1) induces a decomposition

$$D_P = L_P(\mathbb{R})/(K_P A_P) = D_{P,h} \times D_{P,\ell} \tag{21.6.1}$$

and we let

$$q_P \colon D_P \to D_{P^\dagger,h}$$

be the projection onto the first factor. (Note that $D_{P,h} = D_{P^\dagger,h}$ since $\kappa(\widetilde{\Upsilon}) = \kappa(\omega(\widetilde{\Upsilon}))$.) Together these maps yield a surjection

$$q \colon \widehat{D} \to D^*_\sigma \ .$$

Let $\pi \colon \widehat{X} \to X^*_\sigma$ be the map induced by q. Zucker's main result in [57] is that π is a quotient map. (Note however that the quotient topology on D^*_σ induced by q may be finer than the Satake topology on D_σ^* [57, (3.10)].) Note that in the notation of §3.5 we have $\pi^{-1}(F_R) = X_R(L_{R,\ell})$ for any σ-saturated parabolic $\mathbb{Q}$-subgroup R.

22. Functoriality under Inverse Images

Let $k\colon Z \hookrightarrow W$ be an inclusion map of admissible spaces possessing unique maximal strata and let $\mathcal{M}$ be an $\mathcal{L}_W$-module. Let X_S denote the unique maximal stratum of W. We wish to study the micro-support of the inverse images $k^*\mathcal{M}$ and $k^!\mathcal{M}$. In order to preserve the condition of conjugate self-contragredience, we shall sometimes assume certain boundary components are equal-rank: a symmetric space is called *equal-rank* if it can be expressed as $G(\mathbb{R})/K$, where G is a reductive $\mathbb{R}$-group, K is a maximal compact subgroup of $G(\mathbb{R})$, and G is equal-rank (see §8.5). Thus if G is defined over $\mathbb{Q}$ and D is defined as usual to be $G(\mathbb{R})/KA_G$, then D is equal-rank if and only if ${}^0G$ is equal-rank, which is the case if and only if $V|_{{}^0G}$ is conjugate self-contragredient for any regular G-module V.

22.1. **Notation.** Given an admissible space W with unique maximal stratum X_S and an irreducible L_P-module $V \in \mathfrak{Irr}(W)$, recall that Q_V^W (resp. $Q_V'^W$) is the parabolic $\mathbb{Q}$-subgroup containing P with type $\{\, \alpha \in \Delta_P^S \mid \langle \xi_V + \rho, \alpha^\vee \rangle < 0 \,\}$ (resp. $\{\, \alpha \in \Delta_P^S \mid \langle \xi_V + \rho, \alpha^\vee \rangle \leq 0 \,\}$) with respect to P.

22.2. **The case of an open embedding.** The following proposition is clear:

Proposition. *Let $k\colon Z \hookrightarrow W$ be the inclusion of a open admissible subspace and let $\mathcal{M}$ be an $\mathcal{L}_W$-module. Then* $\operatorname{SS}(k^*\mathcal{M}) = \operatorname{SS}(\mathcal{M}) \cap \mathfrak{Irr}(\mathcal{L}_Z)$. (*Note that $k^* = k^!$ in this case.*)

22.3. **More partial orderings on $\mathfrak{Irr}(\mathcal{L}_W)$.** Before treating the case of a closed embedding, we need to generalize the partial ordering of §9.1. Let V, $\tilde{V} \in \mathfrak{Irr}(\mathcal{L}_W)$ be irreducible L_P- and $L_{\tilde{P}}$-modules respectively. Let ${}^+\mathfrak{a}_P^{\tilde{P}*}$ denote the convex cone generated by all $\alpha \in \Delta_P^{\tilde{P}}$. Set $V \preccurlyeq_+ \tilde{V}$ if the following three conditions are fulfilled:

(i) $P \leq \tilde{P}$;
(ii) $V = H^{\ell(w)}(\mathfrak{n}_P^{\tilde{P}}; \tilde{V})_w$ for some $w \in W_P^{\tilde{P}}$;
(iii) $(\xi_V + \rho)|_{\mathfrak{a}_P^{\tilde{P}}} \in {}^+\mathfrak{a}_P^{\tilde{P}*}$.

Similarly set $V \preccurlyeq_- \tilde{V}$ if instead of (iii) the condition

$$(\xi_V + \rho)|_{\mathfrak{a}_P^{\tilde{P}}} \in -{}^+\mathfrak{a}_P^{\tilde{P}*}$$

is fulfilled. If both $V \preccurlyeq_+ \tilde{V}$ and $V \preccurlyeq_- \tilde{V}$ we recover $V \preccurlyeq_0 \tilde{V}$. If $V \preccurlyeq_\pm \tilde{V}$ we set $[\tilde{V} : V] = \ell(w)$ where w is as in (ii).

22.4. **Lemma.** *Both $\preccurlyeq_+$ and $\preccurlyeq_-$ are partial orderings on $\mathfrak{Irr}(\mathcal{L}_W)$. They are generated by the relations where $\#\Delta_P^{\tilde{P}} = 1$.*

Proof. The only issue in the first assertion is transitivity. This is clear for (i) and for (ii) it follows from (3.2.1). As for (iii), suppose $V \preccurlyeq_+ \tilde{V} \preccurlyeq_+ \tilde{\tilde{V}}$. Then

$$(\xi_V + \rho)|_{\mathfrak{a}_P^{\tilde{\tilde{P}}}} = (\xi_V + \rho)|_{\mathfrak{a}_P^{\tilde{P}}} + (\xi_{\tilde{V}} + \rho)|_{\mathfrak{a}_{\tilde{P}}^{\tilde{\tilde{P}}}}$$

so we need to check that the elements of both $\Delta_P^{\tilde{P}}$ and $\Delta_{\tilde{P}}^{\tilde{\tilde{P}}}$ are in ${}^+\mathfrak{a}_P^{\tilde{\tilde{P}}*}$ when viewed as linear functions on $\mathfrak{a}_P^{\tilde{\tilde{P}}} = \mathfrak{a}_P^{\tilde{P}} + \mathfrak{a}_{\tilde{P}}^{\tilde{\tilde{P}}}$. But this is one of Langlands's "geometric lemmas" [13, IV, §6.5(2)].

For the second assertion, suppose that $V \preccurlyeq_+ \tilde{\tilde{V}}$. If $\langle \xi_V + \rho, \alpha^\vee \rangle < 0$ for all $\alpha \in \Delta_P^{\tilde{\tilde{P}}}$, then $(\xi_V + \rho)|_{\mathfrak{a}_P^{\tilde{\tilde{P}}}}$ belongs to the interior of $-{}^+\mathfrak{a}_P^{\tilde{\tilde{P}}*}$ [13, IV, §6.2], which is a contradiction. So there exists $\alpha_0 \in \Delta_P^{\tilde{\tilde{P}}}$ such that $\langle \xi_V + \rho, \alpha_0^\vee \rangle \geq 0$. Let $\tilde{P} \geq P$ have type $\Delta_P^{\tilde{P}} = \{\alpha_0\}$ and decompose $w = w^{\tilde{P}} w_{\tilde{P}} \in W_P^{\tilde{P}} W_{\tilde{P}}^{\tilde{\tilde{P}}}$. Then it is easy to check that $\tilde{V} = H^{\ell(w_{\tilde{P}})}(\mathfrak{n}_{\tilde{P}}^{\tilde{\tilde{P}}}; \tilde{\tilde{V}})_{w_{\tilde{P}}}$ satisfies $V \preccurlyeq_+ \tilde{V} \preccurlyeq_+ \tilde{\tilde{V}}$ and we can use induction. ☐

22.5**. Lemma.** *If $V \preccurlyeq_+ V'$ then $Q_V^W \vee P' \geq Q_{V'}^W$ and $Q_V'^W \vee P' \geq Q_{V'}'^W$. If $V \preccurlyeq_- V'$ then $Q_V^W \vee P' \leq Q_{V'}^W$ and $Q_V'^W \vee P' \leq Q_{V'}'^W$.*

Proof. By Lemma 22.4 it suffices to consider the case where $\Delta_P^{P'} = \{\alpha_0\}$. Suppose $V \preccurlyeq_+ V'$ and thus $\langle \xi_V + \rho, \alpha_0^\vee \rangle \geq 0$. Let $\alpha' \in \Delta_{P'}$ and let $\alpha \in \Delta_P \setminus \Delta_P^{P'}$ be the unique element for which $\alpha|_{\mathfrak{a}_{P'}} = \alpha'$. Since $\xi_{V'} + \rho_{P'} = (\xi_V + \rho)|_{\mathfrak{a}_{P'}}$ we have

$$\langle \xi_{V'} + \rho, \alpha'^\vee \rangle = \langle \xi_V + \rho, \alpha^\vee \rangle - \langle \beta_{\alpha_0}^{P'}, \alpha^\vee \rangle \langle \xi_V + \rho, \alpha_0^\vee \rangle \ .$$

Since $\langle \beta_{\alpha_0}^{P'}, \alpha^\vee \rangle = c \langle \alpha_0, \alpha^\vee \rangle \leq 0$ (where $c > 0$), the inequality $\langle \xi_{V'} + \rho, \alpha'^\vee \rangle < 0$ implies $\langle \xi_V + \rho, \alpha^\vee \rangle < 0$ and thus $Q_V^W \vee P' \geq Q_{V'}^W$. The other assertions are proven similarly. ☐

22.6. **The case of a closed embedding: weak form.** Let $\mathcal{M}$ be an $\mathcal{L}_W$-module.

If $k \colon Z \hookrightarrow W$ is the inclusion of an admissible subspace possessing a unique maximal stratum and $\mathcal{S} \subseteq \mathfrak{Irr}(\mathcal{L}_W)$ is any subset, define

$$\begin{aligned} k^*\mathcal{S} &= \{\, V \in \mathfrak{Irr}(\mathcal{L}_Z) \mid V \preccurlyeq_+ \tilde{V} \text{ for some } \tilde{V} \in \mathcal{S} \,\} \\ k^!\mathcal{S} &= \{\, V \in \mathfrak{Irr}(\mathcal{L}_Z) \mid V \preccurlyeq_- \tilde{V} \text{ for some } \tilde{V} \in \mathcal{S} \,\} \ . \end{aligned} \tag{22.6.1}$$

Proposition. *Let $\hat{\imath}_R \colon \widehat{X}_R \cap W \hookrightarrow W$ be the inclusion of a closed stratum and let $\mathcal{M}$ be an $\mathcal{L}_W$-module. Then*

$$\operatorname{SS}_w(\hat{\imath}_R^*\mathcal{M}) \subseteq \hat{\imath}_R^* \operatorname{SS}_w(\mathcal{M}) \qquad \textit{and} \qquad \operatorname{SS}_w(\hat{\imath}_R^!\mathcal{M}) \subseteq \hat{\imath}_R^! \operatorname{SS}_w(\mathcal{M}) \ .$$

For $V \in \operatorname{SS}_w(\hat{\imath}_R^*\mathcal{M})$ we have estimates

$$\begin{aligned} c(V; \hat{\imath}_R^*\mathcal{M}) &\geq \min_{V \preccurlyeq_+ \tilde{V}} (c(\tilde{V}; \mathcal{M}) + [\tilde{V} : V]) \\ d(V; \hat{\imath}_R^*\mathcal{M}) &\leq \max_{V \preccurlyeq_+ \tilde{V}} (d(\tilde{V}; \mathcal{M}) + [\tilde{V} : V]) \ . \end{aligned} \tag{22.6.2}$$

For $V \in \operatorname{SS}_w(\hat{\imath}_R^!\mathcal{M})$ we have estimates

$$\begin{aligned} c(V; \hat{\imath}_R^!\mathcal{M}) &\geq \min_{V \preccurlyeq_- \tilde{V}} (c(\tilde{V}; \mathcal{M}) + [\tilde{V} : V] + \dim \mathfrak{a}_P^{\tilde{P}}) \\ d(V; \hat{\imath}_R^!\mathcal{M}) &\leq \max_{V \preccurlyeq_- \tilde{V}} (d(\tilde{V}; \mathcal{M}) + [\tilde{V} : V] + \dim \mathfrak{a}_P^{\tilde{P}}) \ . \end{aligned} \tag{22.6.3}$$

Proof. We need to show for all $R \in \mathcal{P}(W)$ that

$$\text{if } V \in \operatorname{SS}_w(\hat{\imath}_R^*\mathcal{M}) \text{ then there exists } \tilde{V} \in \operatorname{SS}_w(\mathcal{M}) \text{ with } V \preccurlyeq_+ \tilde{V} \tag{22.6.4}$$

and

$$c(\tilde{V}; \mathcal{M}) + [\tilde{V} : V] \leq c(V; \hat{\imath}_R^*\mathcal{M}) \leq d(V; \hat{\imath}_R^*\mathcal{M}) \leq d(\tilde{V}; \mathcal{M}) + [\tilde{V} : V] \ , \tag{22.6.5}$$

and the analogous claims for $\operatorname{SS}_w(\hat{\imath}_R^!\mathcal{M})$.

First some notation. Let $P \in \mathcal{P}(\widehat{X}_R \cap W)$ and let V be an irreducible L_P-module. Let $Q \in [Q_V^R, Q_V'^R]$. (Here we write Q_V^R instead of $Q_V^{\widehat{X}_R \cap W}$ and similarly for $Q_V'^R$.) Set $U = (P, R) \cap S$ and let $T = (Q, R) \cap S = Q \vee U$:



By Proposition 3.4(i)(iii) we have $i_P^* \hat{\imath}_Q^! \hat{\imath}_R^* \mathcal{M} = i_P^* \hat{\imath}_T^! \mathcal{M}$ and $i_P^* \hat{\imath}_Q^! \hat{\imath}_R^! \mathcal{M} = i_P^* \hat{\imath}_Q^! \mathcal{M}$. Thus the condition that $V \in \mathrm{SS}_w(\hat{\imath}_R^* \mathcal{M})$ becomes

$$H^j(i_P^* \hat{\imath}_T^! \mathcal{M})_V \neq 0 \tag{22.6.6}$$

for some j and some Q as above, and the condition that $V \in \mathrm{SS}_w(\hat{\imath}_R^! \mathcal{M})$ becomes

$$H^j(i_P^* \hat{\imath}_Q^! \mathcal{M})_V \neq 0\ . \tag{22.6.7}$$

The following case of (22.6.4) is easy and will be used below:

(22.6.8) if $V \in \mathrm{SS}_w(\hat{\imath}_R^* \mathcal{M})$ and $Q_V'^W = Q_V'^R \vee U$, then $V \in \mathrm{SS}_w(\mathcal{M})$.

For $Q_V^W \in [Q_V^R, Q_V^R \vee U]$ and thus $T \in [Q_V^W, Q_V'^W]$. Hence (22.6.6) implies that $V \in \mathrm{SS}_w(\mathcal{M})$. Similarly

(22.6.9) if $V \in \mathrm{SS}_w(\hat{\imath}_R^! \mathcal{M})$ and $Q_V^W = Q_V^R$, then $V \in \mathrm{SS}_w(\mathcal{M})$.

For simplicity we now consider the general case in detail only for $V \in \mathrm{SS}_w(\hat{\imath}_R^* \mathcal{M})$; the changes necessary for $V \in \mathrm{SS}_w(\hat{\imath}_R^! \mathcal{M})$ will be indicated at the end. We will use induction on $\#\Delta_P^S$. Since the case where $\#\Delta_P^S = 0$ is trivial, we may assume that (22.6.4) and (22.6.5) are true for all $R \in \mathcal{P}(W)$ and all irreducible $L_{P'}$-modules V' satisfying $\#\Delta_{P'}^S < \#\Delta_P^S$.

Let $P' \geq P$ have type $\Delta_P^{P'} = \{\, \alpha \in \Delta_P^S \setminus \Delta_P^R \mid \langle \xi_V + \rho, \alpha^\vee \rangle > 0 \,\}$ and let $T' \leq T$ and $U' \leq U$ have types $\Delta_P^{T'} = \Delta_P^T \setminus \Delta_P^{P'}$ and $\Delta_P^{U'} = \Delta_P^U \setminus \Delta_P^{P'}$ respectively:



We can assume that $P' > P$ since otherwise $V \in \mathrm{SS}_w(\mathcal{M})$ by (22.6.8). Consider the short exact sequence

$$0 \to i_P^* \hat\imath_{T'}^! \mathcal{M} \to i_P^* \hat\imath_T^! \mathcal{M} \to i_P^* i_{P'*}(i_{P'}^* \hat\imath_T^! \mathcal{M}) \to 0$$

and corresponding long exact sequence (which is the analogue of the long exact sequence of a triple)

$$\cdots \to H^j(i_P^* \hat\imath_{T'}^! \mathcal{M})_V \to H^j(i_P^* \hat\imath_T^! \mathcal{M})_V \to H^j(i_P^* i_{P'*}(i_{P'}^* \hat\imath_T^! \mathcal{M}))_V \to \cdots \,. \tag{22.6.10}$$

Since by (22.6.6) the middle term is nonzero, either the first or last term must be nonzero as well. If the first term of (22.6.10) is nonzero, then we have similarly to (22.6.8) that $V \in \mathrm{SS}_w(\mathcal{M})$ (since $Q_V^W \leq Q_V^R \vee U' \leq Q \vee U' = T' \leq Q_V'^R \vee U' = Q_V'^W$ in the current situation). If instead the last term of (22.6.10) is nonzero, it follows from (3.6.6) that $H^\ell(\mathfrak{n}_P^{P'}; H^{j-\ell}(i_{P'}^* \hat\imath_T^! \mathcal{M}))_V \neq 0$. Thus

$$H^{j-\ell}(i_{P'}^* \hat\imath_T^! \mathcal{M})_{V'} \neq 0 \tag{22.6.11}$$

for some irreducible $L_{P'}$-module V' satisfying $V \preccurlyeq_+ V'$ and $\ell = [V' : V]$.

Choose $R' \in [P', S] \cap \mathcal{P}(W)$ such that $Q' \equiv R' \cap T \leq Q_{V'}'^{R'}$. (For example, one can set $Q' = Q_{V'}'^W \cap T$ and take $R' = (Q', T) \cap S'$. Another choice of R' will be made in §23.) Let $S' = R' \vee T$:



By Proposition 3.4(i)(iii), $i_{P'}^* \hat\imath_T^! \mathcal{M} = i_{P'}^* \hat\imath_{Q'}^! \hat\imath_{R'}^* \hat\imath_{S'}^! \mathcal{M}$ and so (22.6.11) becomes

$$H^{j-\ell}(i_{P'}^* \hat\imath_{Q'}^! (\hat\imath_{R'}^* \hat\imath_{S'}^! \mathcal{M}))_{V'} \neq 0 \,. \tag{22.6.12}$$

Furthermore we claim that

$$Q^{R'}_{V'} \leq Q' \leq Q'^{R'}_{V'} \ . \tag{22.6.13}$$

The second inequality of (22.6.13) is our hypothesis on Q'; the first inequality of (22.6.13) follows since

$$Q' = R' \cap T = R' \cap (Q \vee U) \geq R' \cap (Q^R_V \vee U) \geq Q^{R'}_V \vee P' \geq Q^{R'}_{V'} \ ,$$

where at the last stage we use Lemma 22.5.

Equations (22.6.12) and (22.6.13) show that $V' \in \mathrm{SS}_w(\hat{\imath}^*_{R'}\hat{\imath}^!_{S'}\mathcal{M})$. By the inductive hypothesis we can apply (22.6.4) to V' and conclude there exists an irreducible $L_{\tilde{P}}$-module $\tilde{V} \in \mathrm{SS}_w(\hat{\imath}^!_{S'}\mathcal{M})$ with $V' \preccurlyeq_+ \tilde{V}$. By transitivity of $\preccurlyeq_+$ (Lemma 22.4) we have $V \preccurlyeq_+ \tilde{V}$. However $S' \geq T \vee \tilde{P} \geq Q^W_V \vee \tilde{P} \geq Q^W_{\tilde{V}}$ by Lemma 22.5, and thus $\tilde{V} \in \mathrm{SS}_w(\mathcal{M})$ by (22.6.9). This proves (22.6.4). Equation (22.6.5) also follows since $[\tilde{V} : V] = [\tilde{V} : V'] + [V' : V]$.

The proof for $\mathrm{SS}_w(\hat{\imath}^!_R\mathcal{M})$ is similar except that we let $P' \geq P$ have type $\Delta^{P'}_P = \{\, \alpha \in \Delta^S_P \setminus \Delta^R_P \mid \langle \xi_V + \rho, \alpha^\vee \rangle < 0 \,\}$ and use the long exact sequence

(22.6.14)

$$\cdots \to H^j(i^*_P\hat{\imath}^!_Q\mathcal{M})_V \to H^j(i^*_P\hat{\imath}^!_{Q\vee P'}\mathcal{M})_V \to H^j(i^*_P i_{P'*}(i^*_{P'}\hat{\imath}^!_{Q\vee P'}\mathcal{M}))_V \to \cdots \ .$$

It is now the first term that is nonzero and hence either the middle or last term (in degree $j-1$) must be nonzero as well. Furthermore we choose R' for which $Q' \equiv R' \cap (Q \vee P') \geq Q^{R'}_{V'}$ and let $S' = (Q', Q \vee P') \cap S$:



(For example we can take $R' = (Q \vee P', Q^W_{V'} \vee Q \vee P') \cap S$ for which $Q' = Q \vee P'$ and $S' = S$.) Equation (22.6.12) must be replaced by

$$H^{j-1-\ell}(i^*_{P'}\hat{\imath}^!_{Q'}(\hat{\imath}^!_{R'}\hat{\imath}^*_{S'}\mathcal{M}))_{V'} \neq 0 \ . \quad \square$$

22.7. *Remark.* The proposition is false in general if SS_w is replaced by SS. We will see in §23 a situation in which this replacement is valid.

22.8. **The case of an inclusion of a fiber.** Say we are given a connected normal $\mathbb{Q}$-subgroup $L_{R,\ell} \subseteq L_R$ for a given $R \in \mathcal{P}$ and set $L_{R,h} = L_R/L_{R,\ell}$; we have a almost direct product decomposition $L_R = \widetilde{L_{R,h}}L_{R,\ell}$ where $\widetilde{L_{R,h}}$ is a lift of $L_{R,h}$. Let $X_R(L_{R,\ell}) \subseteq \widehat{X}_R$ be the partial compactification of X_R in the "$L_{R,\ell}$-directions" as in §3.5. Let $\hat{\imath}_{R,\ell} \colon \widehat{X}_{R,\ell} \hookrightarrow X_R(L_{R,\ell})$ be the inclusion of a generic fiber of the flat bundle $\pi \colon X_R(L_{R,\ell}) \to X_{R,h}$.

Note that any $V \in \mathfrak{Irr}(L_R)$ may be written as $\tilde{V}_h \otimes V_\ell$, where $\tilde{V}_h \in \mathfrak{Irr}(\widetilde{L_{R,h}})$ and $V_\ell \in \mathfrak{Irr}(L_{R,\ell})$; the restriction $\mathrm{Res}^{L_P}_{L_{P,\ell}} V$ is a sum of copies of V_ℓ. For a subset

$\mathcal{S} \subseteq \mathfrak{Irr}(\mathcal{L}_{X_R(L_{R,\ell})})$, define

$$\operatorname{Res}^*_\ell \mathcal{S} = \operatorname{Res}^!_\ell \mathcal{S} = \left\{ V_\ell \in \mathfrak{Irr}(\mathcal{L}_{\widehat{X}_{R,\ell}}) \;\middle|\; \begin{array}{l} V_\ell \text{ is a component of } \operatorname{Res}^{L_P}_{L_{P,\ell}} V \text{ for some} \\ L_P\text{-module } V \in \mathcal{S} \end{array} \right\} . \tag{22.8.1}$$

Proposition. *Let* $\mathcal{M}$ *be an* $\mathcal{L}_{X_R(L_{R,\ell})}$*-module and assume that* $X_{R,h}$ *is equal-rank. Then* $\operatorname{SS}(\hat{\imath}^*_{R,\ell}\mathcal{M}) = \operatorname{Res}^*_\ell \operatorname{SS}(\mathcal{M})$ *and for* $V_\ell \in \operatorname{SS}(\hat{\imath}^*_{R,\ell}\mathcal{M})$ *we have the estimate*

$$[c(V_\ell; \hat{\imath}^*_{R,\ell}\mathcal{M}), d(V_\ell; \hat{\imath}^*_{R,\ell}\mathcal{M})] = \bigcup_{\operatorname{Res}^*_\ell V = V_\ell} [c(V;\mathcal{M}), d(V;\mathcal{M})] . \tag{22.8.2}$$

The same result holds for $\hat{\imath}^!_{R,\ell}\mathcal{M}$ *except for the shift in degree of* $-\dim D_{R,h}$.

Proof. For $P_\ell \leq Q_\ell \in \mathcal{P}(\widehat{X}_{R,\ell})$ we compute via (3.5.1) that

$$i^*_{P_\ell}\hat{\imath}^!_{Q_\ell}(\hat{\imath}^*_{R,\ell}\mathcal{M}) = i^*_{P_\ell}\hat{\imath}^*_{P,\ell}(i^*_P\hat{\imath}^!_Q\mathcal{M}) , \tag{22.8.3}$$

where $P \leq Q \in \mathcal{P}(X_R(L_{R,\ell}))$ are defined as in §3.5. Thus an irreducible module $V_\ell \in \mathfrak{Irr}(L_{P_\ell})$ appears in the cohomology of (22.8.3) if and only if $H(i^*_P\hat{\imath}^!_Q\mathcal{M})_V \neq 0$ for some $V = \tilde{V}_h \otimes V_\ell \in \mathfrak{Irr}(L_P)$. Furthermore in this case $Q_\ell \in [Q^{R,\ell}_{V_\ell}, Q'^{R,\ell}_{V_\ell}]$ if and only if $Q \in [Q_V, Q'_V]$. ☐

23. Functoriality under Inverse Images (continued)

Let F_R be a stratum of a Satake compactification $X^*_\sigma = \Gamma\backslash D^*_\sigma$. In this section we present a generalization of Proposition 22.6 (replacing SS_w by SS) in the case that $\hat{\imath}_R$ is the inclusion of $\pi^{-1}(F_R) = X_R(L_{R,\ell})$, under the assumption that D^*_σ has equal-rank rational boundary components. We begin with two lemmas, for which this assumption is not necessary.

We will use the notation of §§3.5 and 21.

23.1. **Lemma.** *Assume* $V \preccurlyeq_\pm V'$ *where* V *and* V' *are irreducible* L_P*- and* $L_{P'}$*-modules respectively. Set* $R = P^\dagger$ *and* $R' = P'^\dagger$ *and assume* $P' \cap R = P$*. Then*

$$R' \cap (Q^R_V \vee P') = Q^{R'}_{V'} \qquad \text{and} \qquad R' \cap (Q'^R_V \vee P') = Q'^{R'}_{V'} \tag{23.1.1}$$

Proof. For $\alpha \in \Delta_P$ let $\widetilde{\alpha}$ denote the unique element of Δ whose restriction to $\mathfrak{a}_P$ is α. In general $\langle \alpha, \alpha_0^\vee \rangle \neq 0$ for α, $\alpha_0 \in \Delta_P$ if and only if $\Psi \cup \{\widetilde{\alpha}, \widetilde{\alpha}_0\}$ is connected for some $\Psi \subseteq \Delta^P$. This is easy to verify if $\#\Delta^P = 1$ and then one uses induction.

We claim that

$$R \vee P' \geq R'. \tag{23.1.2}$$

To see this, note that if $\alpha \in \Delta_P$ and $\alpha \notin \Delta^R_P \cup \Delta^{P'}_P$, then $\{\tilde{\alpha}\} \cup \kappa(\Delta^P)$ is σ-connected and $\alpha|_{\mathfrak{a}_{P'}} = \alpha' \neq 0$. The first assertion implies that $\{\tilde{\alpha}\} \cup \kappa(\Delta^{P'})$ is σ-connected. Thus $\alpha' = \tilde{\alpha}|_{\mathfrak{a}_{P'}} \notin \Delta^{R'}_{P'}$ and therefore $\alpha \notin \Delta^{R'}_P$. This proves the claim.

Now let $\alpha' \in \Delta^{R'}_{P'}$ and let α be the unique element of $\Delta_P \setminus \Delta^{P'}_P$ for which $\alpha|_{\mathfrak{a}_{P'}} = \alpha'$. Equation (23.1.2) implies in fact that $\alpha \in \Delta^R_P \setminus \Delta^{P'}_P$. We claim that

$$\alpha|_{\mathfrak{a}^{P'}_P} = 0 . \tag{23.1.3}$$

To prove the claim let $\alpha_0 \in \Delta^{P'}_P$. As above if $\langle \alpha, \alpha_0^\vee \rangle \neq 0$ then $\Psi \cup \{\widetilde{\alpha}, \widetilde{\alpha}_0\}$ is connected for some $\Psi \subseteq \Delta^P$. But the assumption $P' \cap R = P$ implies that $\widetilde{\alpha}_0 \notin \omega(\kappa(\Delta^P))$ and hence $\widetilde{\alpha}_0 \in \kappa(\Delta^{P'})$. It follows that $\Psi \subseteq \kappa(\Delta^{P'})$ and thus

$\{\widetilde{\alpha}\} \cup \kappa(\Delta^{P'})$ is σ-connected. But then $\alpha' = \tilde{\alpha}|_{\mathfrak{a}_{P'}} \notin \Delta_{P'}^{R'}$, a contradiction. Hence $\langle \alpha, \alpha_0^\vee \rangle = 0$ for all $\alpha_0 \in \Delta_P^{P'}$, which proves (23.1.3).

Now equation (23.1.3) implies that

$$\langle \xi_{V'} + \rho, \alpha'^\vee \rangle = \langle \xi_V + \rho, \alpha^\vee \rangle \tag{23.1.4}$$

in view of $\xi_{V'} + \rho_{P'} = (\xi_V + \rho)|_{\mathfrak{a}_{P'}}$. It follows that $\alpha' \in \Delta_{P'}^{Q_{V'}^{R'}}$ if and only if $\alpha \in \Delta_P^{Q_V^R}$, that is, $\alpha' \in \Delta_{P'}^{Q_V^R \vee P'}$. Since $\Delta_{P'}^{(Q_V^R \vee P') \cap R'} = \Delta_{P'}^{(Q_V^R \vee P')} \cap \Delta_{P'}^{R'}$, this proves the first equality of (23.1.1); the other follows similarly. □

23.2. **Lemma.** *Assume $V \preccurlyeq_\pm V'$ where V and V' are irreducible L_P- and $L_{P'}$-modules respectively. Set $R = P^\dagger$ and $R' = P'^\dagger$ and assume $P' \cap R = P$. If $V|_{M_P}$ is conjugate self-contragredient and $F_{R'}$ is equal-rank, then $V'|_{M_{P'}}$ is conjugate self-contragredient.*

Proof. Write $V' = \widetilde{V'_h} \otimes V'_\ell$, a tensor product of irreducible modules according to the almost direct product decomposition $L_{P'} = \widetilde{L_{P',h}} L_{P',\ell}$. The module $\widetilde{V'_h}$ is automatically conjugate self-contragredient since $\widetilde{L_{P',h}} \cong \widetilde{L_{R',h}}$ has no $\mathbb{Q}$-split center and $D_{R',h}$ is equal-rank. To analyze V'_ℓ, note that $P' \cap R = P$ and $R = P^\dagger$ implies that the parabolic $P/N_{P'}$ of $L_{P'} = \widetilde{L_{P',h}} L_{P',\ell}$ is obtained by removing simple roots of $\widetilde{L_{P',h}}$. It follows that $N_P^{P'} \subseteq \widetilde{L_{P',h}}$ and that L_P factors as an almost direct product

$$L_P = \widetilde{L_{P,h}} L_{P,\ell}^{P',h} L_{P',\ell}.$$

Let $\operatorname{Res}_{L_{P',\ell}}^{L_P}$ denote the resulting functor that restricts a representation of L_P to a representation of $L_{P',\ell}$. The preceding discussion shows that $\operatorname{Res}_{L_{P',\ell}}^{L_P} H^\ell(\mathfrak{n}_P^{P'}; V')$ is isomorphic to a sum of copies of V'_ℓ. On the other hand, V occurs as a component of $H^\ell(\mathfrak{n}_P^{P'}; V')$, so $\operatorname{Res}_{L_{P',\ell}}^{L_P} V$ is also a sum of copies of V'_ℓ. Since $V|_{M_P}$ is conjugate self-contragredient, this implies $V'_\ell|_{M_{P',\ell}}$ is conjugate self-contragredient. □

23.3. **Proposition.** *Let F_R be a stratum of a Satake compactification X_σ^* and assume that all rational boundary components $D_{R',h} \geq D_{R,h}$ are equal-rank. Let $\hat{\imath}_R \colon X_R(L_{R,\ell}) = \pi^{-1}(F_R) \hookrightarrow \widehat{X}$ be the inclusion. If $\mathcal{M}$ is an $\mathcal{L}_{\widehat{X}}$-module, then*

$$\operatorname{SS}(\hat{\imath}_R^* \mathcal{M}) \subseteq \hat{\imath}_R^* \operatorname{SS}(\mathcal{M}) \qquad \text{and} \qquad \operatorname{SS}(\hat{\imath}_R^! \mathcal{M}) \subseteq \hat{\imath}_R^! \operatorname{SS}(\mathcal{M}) \ .$$

Furthermore the estimates (22.6.2) and (22.6.3) hold in the following strengthened form: we can assume that $V \preccurlyeq_\pm \tilde{V}$ may be factored as $V = V_0 \preccurlyeq_\pm V_1 \preccurlyeq_\pm \cdots \preccurlyeq_\pm V_N = \tilde{V}$ such that

$$P_{i+1} \cap P_i^\dagger = P_i \qquad \text{for } 0 \leq i < N, \tag{23.3.1}$$

$$V_i|_{M_{P_i}} \text{ is conjugate self-contragredient} \qquad \text{for } 0 \leq i \leq N, \text{ and} \tag{23.3.2}$$

$$(\xi_{V_i} + \rho)|_{\mathfrak{a}_{P_i}^{P_{i+1}}} \in \pm \mathfrak{a}_{P_i}^{P_{i+1}*+} \qquad \text{for } 0 \leq i < N. \tag{23.3.3}$$

Proof. We first note that by Lemma 21.4 the σ-saturated parabolic $\mathbb{Q}$-subgroups R have $\mathbb{Q}$-type $\omega(\Upsilon)$ for some σ-connected $\Upsilon \subseteq \Delta$. Equivalently, R is σ-saturated if and only if

$$\gamma^\vee \not\perp \kappa(\Delta^R) \cup \{{}_{\mathbb{Q}}\mu\} \qquad \text{for all } \gamma \in \Delta \setminus \Delta^R \ , \tag{23.3.4}$$

where ${}_{\mathbb{Q}}\mu$ is the highest $\mathbb{Q}$-weight of σ.

Once again we only consider the case of an irreducible L_P-module $V \in \mathrm{SS}(\hat\imath_R^*\mathcal{M})$; the case $V \in \mathrm{SS}(\hat\imath_R^!\mathcal{M})$ involves only minor changes. Since $P \in \mathcal{P}(X_R(L_{R,\ell}))$, we have $R = P^\dagger$, a σ-saturated parabolic $\mathbb{Q}$-subgroup. Let $U(F_R) = \coprod_{F_{R'} \geq F_R} F_{R'}$ be the open star neighborhood of the stratum $F_R \subseteq X_\sigma^*$ and set $W = \pi^{-1}(U(F_R))$. In order to apply the proof of Proposition 22.6 to $V \in \mathrm{SS}(\hat\imath_R^*\mathcal{M})$, we must replace $\mathcal{M}$ by its restriction to W; this is permissible by Proposition 22.2. Now recall that at one point in that proof we obtain an irreducible $L_{P'}$-module $V' \in \mathrm{SS}_w(\hat\imath_{R'}^*\hat\imath_{S'}^!\mathcal{M})$ satisfying $V \preccurlyeq_+ V'$ and $P' \cap R = P$ and we were free to choose R' provided that $R' \cap (Q \vee U) \leq Q_{V'}^{\prime R'}$. We now choose $R' = P'^\dagger$. Since $Q \in [Q_V^R, Q_V^{\prime R}]$ and $R' \cap U = P'$ by (23.1.2), Lemma 23.1 implies these conditions are satisfied and Lemma 23.2 implies that $V'|_{M_{P'}}$ is conjugate self-contragredient given that $V|_{M_P}$ is, so $V' \in \mathrm{SS}(\hat\imath_{R'}^*\hat\imath_{S'}^!\mathcal{M})$. Also $(\xi_V + \rho)|_{\mathfrak{a}_P^{P'}} \in \mathfrak{a}_P^{P'*+}$ since by definition of P' all $\alpha \in \Delta_P^{P'}$ satisfy $\langle \xi_V + \rho, \alpha^\vee \rangle > 0$.

Finally we need to verify that the passing by induction from V to V' preserves our new assumptions. Observe that $R' < S'$ and thus S' is also σ-saturated by the criterion (23.3.4). Let $U^{S'}(F_{R'}) = U(F_{R'}) \cap F_{S'}^*$ be the open star neighborhood of the stratum $F_{R'}$ in the induced Satake compactification $F_{S'}^* = \mathrm{cl}(F_{S'})$. Set $W' = \pi^{-1}(U^{S'}(F_{R'}))$. Consider the diagram of inclusions where the vertical arrows are open embeddings:

$$\begin{tikzcd}
\widehat{X}_{R'} \cap W \arrow[r, hook, "\hat\imath_{R'}"] & \widehat{X}_{S'} \cap W \\
X_{R'}(L_{R',\ell}) \arrow[u, hook, "k_{R'}"] \arrow[r, hook, "\hat\imath_{R'}"'] & W' \arrow[u, hook, "k_{S'}"'] .
\end{tikzcd}$$

By Proposition 22.2, $V' \in \mathrm{SS}(k_{R'}^*\hat\imath_{R'}^*\hat\imath_{S'}^!\mathcal{M}) = \mathrm{SS}(\hat\imath_{R'}^*k_{S'}^*\hat\imath_{S'}^!\mathcal{M})$, and we can use induction. □

24. The Basic Lemma

We now address the issue of estimating the term $[\tilde V : V] = \ell(w)$ in (22.6.2) and (22.6.3) within the context of Proposition 23.3. Our tool is a basic lemma which, under the assumption that $V|_{M_P}$ is conjugate self-contragredient, establishes the fundamental relationship between the geometry of ξ_V *vis a vis* an $S_P^{\tilde P}$-root α and the contribution to the length of w attributable to α. This basic lemma is important in other contexts; a simple form already occurred as Lemma 17.6.

For simplicity we will assume that $\tilde V = E$ is an irreducible G-module. We use the notation of §17.6.

24.1. Let $\mathfrak{h}$ be a fundamental Cartan subalgebra of $\mathfrak{l}_P$ and let $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})$ be a θ-stable positive system for $\mathfrak{l}_P$. Let V be an irreducible L_P-module with highest weight $\mu \in \mathfrak{h}_\mathbb{C}^*$ and assume $V|_{M_P}$ is conjugate self-contragredient. We have defined a reductive $\mathbb{R}$-subgroup $L_P(\mu) \subseteq L_P$ in §§10.1 and 10.2 with roots

$$\{\, \gamma \in \Phi(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C}) \mid \langle \mu, \gamma^\vee \rangle = 0 \,\} . \tag{24.1.1}$$

Via a lift of L_P to P, we obtain an adjoint representation of $\mathfrak{l}_P$ (and hence $\mathfrak{l}_P(\mu)$) on $\mathfrak{n}_{P\mathbb{C}}$. There is a unique decomposition

$$\mathfrak{n}_{P\mathbb{C}} = \bigoplus_{\substack{\mathfrak{l}_P(\mu)\text{-irreducible} \\ \text{submodules} \\ F\subseteq \mathfrak{n}_{P\mathbb{C}}}} F,$$

which coarsens the decomposition into root spaces. Set $\tau'_P = -\tau_P$ and observe that if F is an irreducible component as above with weights $\Phi(F, \mathfrak{h}_\mathbb{C})$, then $\tau'_P(\Phi(F, \mathfrak{h}_\mathbb{C}))$ is the set of weights of another irreducible component, which we denote $\tau'_P(F)$; this follows from (8.2.2). In this way we obtain an action of τ'_P on the $\mathfrak{l}_P(\mu)$-irreducible components of $\mathfrak{n}_{P\mathbb{C}}$, generalizing the action of τ'_P on the roots of $\mathfrak{n}_{P\mathbb{C}}$. Let[15]

$$\mathfrak{n}_P(\mu) = \bigoplus_{\tau'_P(F)=F} F;$$

since the action of $\mathfrak{l}_P(\mu)$ on $\mathfrak{n}_{P\mathbb{C}}$ preserves $\mathfrak{n}_{\alpha\mathbb{C}}$ for any $\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)$, we may likewise define $\mathfrak{n}_\alpha(\mu)$. Although $\mathfrak{n}_P(\mu)$ and $\mathfrak{n}_\alpha(\mu)$ depend on the choice of a lift of L_P to G, their dimensions are independent of this choice. The dimension of $\mathfrak{n}_P(\mu)$ can vary depending on the choice of $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})$ however; let $\mathfrak{n}_P(V)$ denote any one of the $\mathfrak{n}_P(\mu)$ with maximal dimension.

24.2. **The Basic Lemma.** *Let P be a parabolic $\mathbb{Q}$-subgroup and let $w \in W_P$. Let $V = H^{\ell(w)}(\mathfrak{n}_P; E)_w$ have highest weight μ with respect to a θ-stable positive system. Assume $V|_{M_P}$ is conjugate self-contragredient. For any $\alpha \in \Phi(\mathfrak{n}_P, \mathfrak{a}_P)$ we have:*

(i) $\langle \xi_V + \rho, \alpha^\vee \rangle \le 0 \implies \ell_\alpha(w) \ge \frac{1}{2}(\dim \mathfrak{n}_\alpha + \dim \mathfrak{n}_\alpha(\mu))$.

(ii) $\langle \xi_V + \rho, \alpha^\vee \rangle = 0 \implies \ell_\alpha(w) = \frac{1}{2}\dim \mathfrak{n}_\alpha$ *and* $\mathfrak{n}_\alpha(\mu) = 0$.

(iii) $\langle \xi_V + \rho, \alpha^\vee \rangle \ge 0 \implies \ell_\alpha(w) \le \frac{1}{2}(\dim \mathfrak{n}_\alpha - \dim \mathfrak{n}_\alpha(\mu))$.

Furthermore if $F \subseteq \mathfrak{n}_{\alpha\mathbb{C}}$ is an irreducible $\mathfrak{l}_P(\mu)$-submodule, then Φ_w contains respectively (i) *at least one,* (ii) *exactly one,* (iii) *at most one of $\Phi(F, \mathfrak{h}_\mathbb{C})$ and $\Phi(\tau'_P(F), \mathfrak{h}_\mathbb{C})$.*

Proof. Let $W^{L_P(\mu)}$ denote the Weyl group of $L_P(\mu)$ and let $\lambda \in \mathfrak{h}_\mathbb{C}$ be the highest weight of E relative to $\Phi^+ = \Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C}) \cup \Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_\mathbb{C})$. We begin by establishing

$$s \in W^{L_P(\mu)} \quad \Rightarrow \quad s\Phi_w = \Phi_w \text{ and } sw\lambda = w\lambda \tag{24.2.1}$$

following [15]. Express the highest weight $\mu = w(\lambda+\rho) - \rho$ of V as $w\lambda - \sum_{\gamma\in\Phi_w}\gamma$. Since s may be written as a product of reflections in simple coroots orthogonal to μ we see that $\mu = s\mu$ and hence

$$w\lambda - \sum_{\gamma\in\Phi_w} \gamma = \mu = s\mu = sw\lambda - \sum_{\gamma\in s\Phi_w}\gamma.$$

Note that $s\Phi_w \subseteq \Phi^+$; by canceling the terms of the right hand sum that lie in Φ_w with the corresponding terms of the left hand sum one may compute that

$$\lambda = w^{-1}sw\lambda - \sum_{\gamma\in(w^{-1}\Psi)\cap\Phi^+} \gamma, \tag{24.2.2}$$

where $\Psi = s\Phi_w \cup -(\Phi^+ \setminus s\Phi_w)$. Since λ is dominant, we may replace $w^{-1}sw\lambda$ in (24.2.2) by $\lambda - \sum_i m_i\alpha_i$ for α_i simple and $m_i \ge 0$. In addition since the roots of

---

[15] The quantities $\mathfrak{l}_P(\mu)$ and $\mathfrak{n}_P(\mu)$ are defined quite differently despite the similarity of notation. We trust that with this warning there should be no confusion.

$(w^{-1}\Psi) \cap \Phi^+$ being subtracted in (24.2.2) are positive, it follows that all $m_i = 0$ and that $(w^{-1}\Psi) \cap \Phi^+ = \emptyset$; these two facts are equivalent to the desired (24.2.1).

Now consider an $\mathfrak{l}_P(\mu)$-irreducible submodule $F \subseteq \mathfrak{n}_{\alpha\mathbb{C}}$. Since by (24.2.1) Φ_w is stable under $W^{L_P(\mu)}$, either all extremal weights of $\Phi(F, \mathfrak{h}_{\mathbb{C}})$ belong to Φ_w or none do. Since furthermore membership in Φ_w is given by a linear inequality on positive roots and $\Phi(F, \mathfrak{h}_{\mathbb{C}})$ is the convex hull of the extremal weights, this implies that either all of $\Phi(F, \mathfrak{h}_{\mathbb{C}})$ is contained in Φ_w or none of it is. An application of Lemma 17.6 to any $\gamma \in \Phi(F, \mathfrak{h}_{\mathbb{C}})$ concludes the proof. □

24.3. **Corollary.** *Let P be a parabolic $\mathbb{Q}$-subgroup and let V be an irreducible constituent of $H^i(\mathfrak{n}_P; E)$ for which $V|_{M_P}$ is conjugate self-contragredient. Then*

$$(\xi_V + \rho)|_{\mathfrak{a}_P} \in -\mathfrak{a}_P^{*+} \implies i \geq \frac{1}{2}(\dim \mathfrak{n}_P + \dim \mathfrak{n}_P(V))$$

and

$$(\xi_V + \rho)|_{\mathfrak{a}_P} \in \mathfrak{a}_P^{*+} \implies i \leq \frac{1}{2}(\dim \mathfrak{n}_P - \dim \mathfrak{n}_P(V)).$$

24.4. *Remark.* All results in this section and their proofs generalize immediately to the situation in which E is replaced by a "virtual" irreducible G-module, whose highest weight λ is allowed to be any dominant element of $\mathfrak{h}_{\mathbb{C}}^*$, not necessarily integral. They also generalize to parabolic k-subgroups where k is any subfield of $\mathbb{R}$.

25. The Basic Lemma in the Presence of Equal-Rank Real Boundary Components

We need to replace the term $\dim \mathfrak{n}_P(V)$ appearing in Corollary 24.3 by a more geometric expression. This can be done in the presence of a Satake compactification with equal-rank *real* boundary components.

25.1. Let V be an irreducible L_P-module and assume that $V|_{M_P}$ is conjugate self-contragredient. Let $D_P(V)$ be defined as in §10.2. Let $D \subseteq {}_{\mathbb{R}}D_\sigma^*$ be a Satake compactification. The decomposition (21.4.1) induces a decomposition of symmetric spaces

$$D_P(V) = D_{P,h}(V) \times D_{P,\ell}(V)\ . \tag{25.1.1}$$

Here $D_{P,h}(V)$ is defined using $\operatorname{Res}^{L_P}_{\widetilde{L_{P,h}}} V$ and similarly for $D_{P,\ell}(V)$; see the last comment in §10.2.

25.2. **Lemma.** *Let P be a parabolic $\mathbb{Q}$-subgroup and let $D \subseteq {}_{\mathbb{R}}D_\sigma^*$ be a Satake compactification for which we assume all real boundary components $D_{R',h} \geq D_{P^\dagger,h}$ are equal-rank. Let V be an irreducible constituent of $H(\mathfrak{n}_P; E)$ for which $V|_{M_P}$ is conjugate self-contragredient. Then*

$$\dim \mathfrak{n}_P(V) \geq \dim \mathfrak{a}_P^G + \dim D_{P,\ell}(V)\ .$$

Remark. A similar result was proved by Borel in [7, §5.5] under the weaker assumptions that D and $D_{P^\dagger,h}$ are equal-rank and a certain condition (B) holds. The condition (B) needed to be verified for each desired case and Borel expressed the hope that it could be replaced by a more conceptual argument. The proof offered below, while not transparent, is at least free from case by case analysis.

Note also that condition (B) does not hold[16] in one case of interest where the lemma above *does* apply: $G = \mathrm{SO}(p,q)$ with $p+q$ odd with the Satake compactification in which σ is connected only to the short simple $\mathbb{R}$-root. In this case there are imaginary noncompact roots in $\mathfrak{l}_P$ (relative to a fundamental Cartan subalgebra) which are strongly orthogonal to all real roots in $\mathfrak{n}_P$; the hypothesis that the intermediate real boundary components are equal-rank and its use below in Sublemma 25.3 was needed precisely to handle this situation.

Proof of Lemma 25.2. We first establish notation. Let θ be a Cartan involution of $L_P(\mathbb{R})$ with Cartan decomposition $\mathfrak{l}_P = \mathfrak{k}_P + \mathfrak{p}_P + \mathfrak{a}_P$. Let $\mathfrak{h} = \mathfrak{b}_{P,\mathfrak{k}} + \mathfrak{b}_{P,\mathfrak{p}} + \mathfrak{a}_P$ be a fundamental θ-stable Cartan subalgebra of $\mathfrak{l}_P$ and choose a θ-stable positive system $\Phi^+(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ for $\mathfrak{l}_{P\mathbb{C}}$; thus $\tau_P = \theta$. Let μ be the highest weight of V and let $L_P(\mu)$ be defined as in §10.1 with Cartan decomposition $\mathfrak{k}_P(\mu) + \mathfrak{p}_P(\mu) + \mathfrak{a}_P$. Let $\Phi^+(\mathfrak{p}_P(\mu)_{\mathbb{C}}, \mathfrak{b}_{P,\mathfrak{k}\mathbb{C}})$ denote the positive nonzero weights of $\mathfrak{b}_{P,\mathfrak{k}\mathbb{C}}$ in $\mathfrak{p}_P(\mu)_{\mathbb{C}}$, counted with multiplicity. We extend the preceding notation to $L_{P,h}$ and $L_{P,\ell}$ by adding the appropriate subscript. Finally note that any reductive subalgebra $\mathfrak{r} \subseteq \mathfrak{l}_{P\mathbb{C}}$ acts on $\mathfrak{n}_{P\mathbb{C}}$ by the adjoint action of a lift of $\mathfrak{l}_{P\mathbb{C}}$ and define $\mathfrak{n}_P(\mathfrak{r})$ to be the sum of the irreducible $\mathfrak{r}$-submodules whose set of weights are τ_P'-stable; thus $\mathfrak{n}_P(\mathfrak{l}_P(\mu)) = \mathfrak{n}_P(\mu)$ as in §24.1. It will suffice to prove $\dim \mathfrak{n}_P(\mu) \geq \dim \mathfrak{a}_P^G + \dim D_{P,\ell}(\mu)$.

Clearly

$$\dim \mathfrak{a}_P^G + \dim D_{P,\ell}(\mu) = 2 \cdot \#\Phi^+(\mathfrak{p}_{P,\ell}(\mu)_{\mathbb{C}}, \mathfrak{b}_{P,\mathfrak{k}\mathbb{C}}) + \dim \mathfrak{b}_{P,\mathfrak{p}} + \dim \mathfrak{a}_P^G \ . \tag{25.2.1}$$

We begin by bounding the last two terms. Consider $\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}})$ as defined above; the set $\Phi(\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}})$ consists of the positive roots γ satisfying $\tau_P'\gamma = \gamma$, that is, the real roots. We claim that

$$\dim \mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}) \geq \dim \mathfrak{b}_{P,\mathfrak{p}} + \dim \mathfrak{a}_P^G \ . \tag{25.2.2}$$

In fact there exists a set of strongly orthogonal roots $\{\delta_1, \dots, \delta_r\}$ satisfying $\tau_P'\delta_i = \delta_i$ and which span $\mathfrak{b}_{P,\mathfrak{p}}^* + \mathfrak{a}_P^{G*}$. To see this, proceed as in [7] to write the centralizer in ${}^0\mathfrak{g}$ of $\mathfrak{b}_{P,\mathfrak{k}}$ as $\mathfrak{b}_{P,\mathfrak{k}} + \mathfrak{z}_1$. Then $\mathfrak{z}_1$ has both a split Cartan subalgebra $\mathfrak{b}_{P,\mathfrak{p}} + \mathfrak{a}_P^G$ and, since by hypothesis ${}^0G = L_{G,h}$ is equal-rank, a compact Cartan subalgebra. The existence of $\{\delta_1, \dots, \delta_r\}$ then follows from [30], [52, Prop. 11], [53].

We now attempt to bound the rest of (25.2.1). For $\gamma_{\mathfrak{k}} \in \Phi^+(\mathfrak{p}_{P,\ell}(\mu)_{\mathbb{C}}, \mathfrak{b}_{P,\ell,\mathfrak{k}\mathbb{C}})$ choose $\gamma \in \Phi^+(\mathfrak{l}_{P,\ell}(\mu)_{\mathbb{C}}, \mathfrak{b}_{P,\ell\mathbb{C}})$ such that $\gamma|_{\mathfrak{b}_{P,\ell,\mathfrak{k}\mathbb{C}}} = \gamma_{\mathfrak{k}}$. The root $\tau_P\gamma$ is also in $\Phi^+(\mathfrak{l}_{P,\ell}(\mu)_{\mathbb{C}}, \mathfrak{b}_{P,\ell\mathbb{C}})$ since $\tau_P(\mu|_{\mathfrak{b}_P}) = \mu|_{\mathfrak{b}_P}$. Let $\mathfrak{r}_\gamma$ denote the reductive subalgebra $\mathfrak{h}_{\mathbb{C}} \subseteq \mathfrak{r}_\gamma \subseteq \mathfrak{l}_P(\mu)_{\mathbb{C}}$ with root system $\Phi \cap \operatorname{Span}\{\gamma, \tau_P\gamma\}$. Clearly

$$\Phi(\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}}) \subseteq \Phi(\mathfrak{n}_P(\mathfrak{r}_\gamma), \mathfrak{h}_{\mathbb{C}}) \subseteq \Phi(\mathfrak{n}_P(\mathfrak{l}_P(\mu)_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}}) = \Phi(\mathfrak{n}_P(\mu), \mathfrak{h}_{\mathbb{C}}).$$

We will see below in Sublemma 25.3(i)(iii) that there exists a pair of roots $\eta \neq \tau_P'\eta \in \Phi(\mathfrak{n}_P(\mathfrak{r}_\gamma), \mathfrak{h}_{\mathbb{C}}) \setminus \Phi(\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}}) \subseteq \Phi(\mathfrak{n}_P(\mu), \mathfrak{h}_{\mathbb{C}})$. This would imply

$$\dim \mathfrak{n}_P(\mu) - \dim \mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}) \geq 2 \cdot \#\Phi^+(\mathfrak{p}_{P,\ell}(\mu)_{\mathbb{C}}, \mathfrak{b}_{P,\mathfrak{k}\mathbb{C}}) \tag{25.2.3}$$

(and hence prove the lemma in view of (25.2.1) and (25.2.2)) except that we have to deal with the possibility that the sets $\Phi(\mathfrak{n}_P(\mathfrak{r}_\gamma), \mathfrak{h}_{\mathbb{C}}) \setminus \Phi(\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}})$ for the different $\gamma_{\mathfrak{k}}$ may not be disjoint.

Since $|\gamma| = |\tau_P\gamma|$ and no multiple of $\gamma - \tau_P\gamma$ can be a root (recall that $\mathfrak{l}_{P\mathbb{C}}$ has no roots negated by τ_P), one easily checks that $\{\gamma, \tau_P\gamma\}$ is a base for the root system of $\mathfrak{r}_\gamma$ and the type is A_1 (if $\gamma = \tau_P\gamma$), $A_1 \times A_1$, or A_2. Now consider $\gamma_{\mathfrak{k}} \neq \gamma_{\mathfrak{k}}' \in$

---

[16] In [7, §6.6] it is mistakenly asserted that condition (B) holds in this case.

$\Phi^+(\mathfrak{p}_{P,\ell}(\mu)_{\mathbb{C}}, \mathfrak{b}_{P,\ell,\mathfrak{k}\mathbb{C}})$ and choose corresponding roots $\gamma \neq \gamma' \in \Phi^+(\mathfrak{l}_{P,\ell}(\mu)_{\mathbb{C}}, \mathfrak{b}_{P,\ell\mathbb{C}})$. Suppose that $\Phi(\mathfrak{n}_P(\mathfrak{r}_\gamma), \mathfrak{h}_{\mathbb{C}}) \setminus \Phi(\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}})$ and $\Phi(\mathfrak{n}_P(\mathfrak{r}_{\gamma'}), \mathfrak{h}_{\mathbb{C}}) \setminus \Phi(\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}})$ have nonempty intersection. Then there exists a root $\eta \in \Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ for which $\tau'_P\eta - \eta$ is a nonzero $\mathbb{Z}$-linear combination of both $\{\gamma, \tau_P\gamma\}$ and $\{\gamma', \tau_P\gamma'\}$. Since $\tau'_P\eta - \eta$ is τ_P-invariant, this means that $\gamma + \tau_P\gamma$ and $\gamma' + \tau_P\gamma'$ are scalar multiples of each other. But $(1/2)(\gamma + \tau_P\gamma)$ belongs to either the reduced root system $\Phi(\mathfrak{l}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ (if $\gamma = \tau_P\gamma$) or the reduced root system $\Phi(\mathfrak{k}_{P\mathbb{C}}, \mathfrak{b}_{P,\mathfrak{k}\mathbb{C}})$ (if $\gamma \neq \tau_P\gamma$), and similarly for $(1/2)(\gamma' + \tau_P\gamma')$. Thus $\gamma + \tau_P\gamma$ and $\gamma' + \tau_P\gamma'$ can be scalar multiples of each other only if γ (say) satisfies $\gamma \neq \tau_P\gamma$ and γ' satisfies $\gamma' = \tau_P\gamma'$. It follows from our comments earlier then that $\gamma' = \gamma + \tau_P\gamma$ and $\mathfrak{r}_\gamma$ has type A_2.

So $\Phi(\mathfrak{n}_P(\mathfrak{r}_\gamma), \mathfrak{h}_{\mathbb{C}}) \setminus \Phi(\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}})$ can intersect with at most one other such set and it will suffice to show that in this case $\Phi(\mathfrak{n}_P(\mathfrak{r}_\gamma), \mathfrak{h}_{\mathbb{C}}) \setminus \Phi(\mathfrak{n}_P(\mathfrak{h}_{\mathbb{C}}), \mathfrak{h}_{\mathbb{C}})$ contains an additional pair of roots $\eta' \neq \tau'_P\eta'$. But if such a pair of roots did not exist, the only nontrivial $\mathfrak{r}_\gamma$-submodule of $\mathfrak{n}_P(\mathfrak{r}_\gamma)$ would have weights (after interchanging η and $\tau'_P\eta$ if necessary) $\{\eta, \eta + \gamma = \tau'_P\eta - \tau_P\gamma, \tau'_P\eta\}$ and hence $\{\eta, \gamma, \tau_P\gamma\}$ would be the base of a root subsystem of type A_3. Sublemma 25.3(ii) following shows this situation cannot occur. ☐

25.3. Sublemma. *As in the proof of Lemma* 25.2 *let* $\gamma_{\mathfrak{k}} \in \Phi^+(\mathfrak{p}_{P,\ell\mathbb{C}}, \mathfrak{b}_{P,\ell,\mathfrak{k}\mathbb{C}})$ *and choose* $\gamma \in \Phi^+(\mathfrak{l}_{P,\ell\mathbb{C}}, \mathfrak{b}_{P,\ell\mathbb{C}})$ *such that* $\gamma|_{\mathfrak{b}_{P,\ell,\mathfrak{k}\mathbb{C}}} = \gamma_{\mathfrak{k}}$.

- (i) *If* $\tau_P\gamma \neq \gamma$*, then there exists a root* $\delta = \tau'_P\delta \in \Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ *such that* (*after interchanging* γ *and* $\tau_P\gamma$ *if necessary*) $\eta = \delta - \gamma$ *and* $\tau'_P\eta = \delta + \tau_P\gamma$ *are also roots.*
- (ii) *If in* (i) *the roots* $\{\eta, \gamma, \tau_P\gamma\}$ *are a base of a root subsystem of type* A_3*, then there also exists a root* $\widetilde{\delta} \neq \delta$ *with* $\widetilde{\delta} = \tau'_P\widetilde{\delta}$ *and one of* $\widetilde{\delta} \pm \gamma$ *a root.*
- (iii) *If* $\tau_P\gamma = \gamma$*, then there exists a root* $\eta \in \Phi(\mathfrak{n}_{P\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ *such that* $\tau'_P\eta = \eta + k\gamma$ *for some* $k \in \mathbb{Z} \setminus \{0\}$.

Proof. Let $\{\delta_1, \dots, \delta_r\}$ be a set of strongly orthogonal roots spanning $\mathfrak{b}^*_{P,\mathfrak{p}} + \mathfrak{a}^{G\,*}_P$ and satisfying $\tau'_P\delta_i = \delta_i$ as in the proof of Lemma 25.2. If $\tau_P\gamma \neq \gamma$ then $\gamma|_{\mathfrak{b}_{P,\mathfrak{p}} + \mathfrak{a}^G_P} \neq 0$ and so there exists $\delta = \delta_i$ such that $\langle \gamma, \delta^\vee \rangle \neq 0$. By interchanging γ and $\tau_P\gamma$ if necessary, we may assume $\langle \gamma, \delta^\vee \rangle > 0$. Then $\eta = \delta - \gamma$ is a root which proves (i). Part (ii) will follow if we show there exists $\widetilde{\delta} = \delta_j \neq \delta_i$ with $\langle \gamma, \widetilde{\delta}^\vee \rangle \neq 0$ as well. But if such a $\widetilde{\delta}$ did not exist, $\gamma|_{\mathfrak{b}_{P,\mathfrak{p}} + \mathfrak{a}^G_P} = c\delta$ for some scalar c and it is easy to check that the conditions on c implied by the equalities $|\delta - \gamma| = |\gamma| = |\delta|$ and $2(\gamma, \tau_P\gamma)/|\gamma|^2 = -1$ for a Weyl group invariant inner product on a root system of type A_3 are inconsistent.

If $\tau_P\gamma = \gamma$ we claim that there is a nontrivial γ-root string through some root in $\mathbb{R}\gamma + \mathfrak{b}^*_{P,\mathfrak{p}} + \mathfrak{a}^{G\,*}_P$. On such a root string the operator τ'_P acts as reflection in γ and hence the root string is τ'_P-stable. Thus part (iii) follows from the claim if we set η to be any root in the root string other than at the center.

The claim is obvious if $\delta_i + \gamma$ or $\delta_i - \gamma$ is a root for some i we are done, so we assume that γ is strongly orthogonal to $\{\delta_1, \dots, \delta_r\}$. Recall [29, Chapter VII, §7] that a parabolic $\mathbb{R}$-subgroup R is called *cuspidal*[17] if $L_R/Z(L_R)$ is equal-rank, where $Z(L_R)$ denotes the center. Let $R \subseteq P$ be a cuspidal parabolic $\mathbb{R}$-subgroup with ${}_{\mathbb{R}}\mathfrak{b}_R = \mathfrak{b}_{P,\mathfrak{k}}$ and split component ${}_{\mathbb{R}}\mathfrak{a}_R = \mathfrak{b}_{P,\mathfrak{p}} + \mathfrak{a}_P$; to construct R apply [29,

---

[17] This usage is different from that in [33] or [9]; R being cuspidal in those references corresponds here to R being defined over $\mathbb{Q}$.

Prop. 7.87]. Since $\tau_R = \tau_P = \theta$ on $\mathfrak{h}$, γ remains an imaginary noncompact root for $\mathfrak{l}_{R\mathbb{C}}$. Thus there is a Cayley transformation [29, Chapter VI, §7] which transforms γ into a real root. That is, there exists a inner automorphism $\mathbf{c}_\gamma$ of $\mathfrak{l}_{R\mathbb{C}}$ such that $\mathfrak{h}' \equiv \mathfrak{l}_R \cap \mathbf{c}_\gamma(\mathfrak{h}_\mathbb{C})$ is a θ-stable Cartan subalgebra and $\mathbf{c}_\gamma \circ (s_\gamma \circ \theta) = \theta \circ \mathbf{c}_\gamma$ on $\mathfrak{h}$. Let $R' \subset R$ be a cuspidal parabolic $\mathbb{R}$-subgroup with ${}_\mathbb{R}\mathfrak{b}_{R'} = \mathbf{c}_\gamma(\operatorname{Ker}\gamma \cap {}_\mathbb{R}\mathfrak{b}_R)$ and ${}_\mathbb{R}\mathfrak{a}_{R'} = \mathbb{R}\mathbf{c}_\gamma(\gamma^\vee) + {}_\mathbb{R}\mathfrak{a}_R$. The roots $\{\delta_1, \dots, \delta_r, \mathbf{c}_\gamma(\gamma)\}$ form a strongly orthogonal basis of ${}_\mathbb{R}\mathfrak{a}_{R'}^{G*}$ and to prove our claim we need to demonstrate that there is a nontrivial $\mathbf{c}_\gamma(\gamma)$-root string through some root in ${}_\mathbb{R}\mathfrak{a}_{R'}^{G*}$.

From our hypotheses on the Satake compactification we now construct another independent set $\{\delta'_1, \dots, \delta'_{r+1}\}$ of roots spanning ${}_\mathbb{R}\mathfrak{a}_{R'}^{G*}$. Define a sequence of parabolic $\mathbb{R}$-subgroups

$$R' = R'_0 \subset R'_1 \subset \cdots \subset R'_{r+1} = G$$

inductively by choosing R'_i to have type ${}_\mathbb{R}\Delta_{R'}^{R'_i} = \{\alpha_1, \dots, \alpha_i\}$ where $\alpha_i \in {}_\mathbb{R}\Delta_{R'}^{R', R'^{\dagger}_{i-1}}$. Thus the corresponding real Satake boundary component strictly increases at each stage,

$$D_{R'^\dagger,h} = D_{R'^\dagger_0,h} \lneq D_{R'^\dagger_1,h} \lneq \cdots \lneq D_{R'^\dagger_{r+1},h} = D\ .$$

On the other hand, since $L_{P,h}$ is equal-rank and γ was a root of $\mathfrak{l}_{P,\ell}$, we have $L_{R',h} = L_{P,h}$ and hence $D_{P^\dagger,h} = D_{R'^\dagger,h}$. We now claim that each parabolic subgroup R'_i is cuspidal. This follows since $L_{R'_i,\ell}/Z(L_{R'_i})$ is an almost direct factor of $L_{R'_{i-1},\ell}/Z(L_{R'_{i-1}})$, which is equal-rank by induction, and $L_{R'_i,h}$ is equal-rank since the boundary component $D_{R'^\dagger_i,h}$ is equal-rank by hypothesis. Thus for each $i = 1, \dots, r+1$ there exists a root δ'_i which spans ${}_\mathbb{R}\mathfrak{a}_{R'_{i-1}}^{R'_i\,*}$ [30], [52], [53].

Since $\mathbf{c}_\gamma(\gamma) \in {}_\mathbb{R}\mathfrak{a}_{R'}^{G*}$ is nonzero there exists $1 \le i \le r+1$ such that $\langle \mathbf{c}_\gamma(\gamma), \delta_i'^\vee \rangle \neq 0$. The desired nontrivial $\mathbf{c}_\gamma(\gamma)$-root string would now exist provided $\mathbf{c}_\gamma(\gamma) \neq \pm\delta'_i$. To show $\mathbf{c}_\gamma(\gamma)$ cannot equal $\pm\delta'_i$, write ${}_\mathbb{R}\Delta_{R'}^{R} = \{\alpha_k\}$ where $1 \le k \le r+1$. Thus $\mathbf{c}_\gamma(\gamma)$ is a multiple of α_k. On the other hand, δ'_i is proportional to the projection of α_i to ${}_\mathbb{R}\mathfrak{a}_{R'_{i-1}}^{R'_i}$, that is, $\delta'_i \sim \alpha_i + \sum_{j<i} c_j \alpha_j$ where all $c_j = 0$ if and only if $\alpha_i \in {}_\mathbb{R}\mathfrak{a}_{R'_{i-1}}^{R'_i}$. So if $\mathbf{c}_\gamma(\gamma)$ were to equal $\pm\delta'_i$ then $k = i$ and $\alpha_i \in {}_\mathbb{R}\mathfrak{a}_{R'_{i-1}}^{R'_i}$. This implies that the simple $\mathbb{R}$-roots underlying α_i as opposed to $\{\alpha_1, \dots, \alpha_{i-1}\}$ belong to different components of $L_{R'_i,h}$. This implies that the condition $\alpha_i \in {}_\mathbb{R}\Delta_{R'}^{R', R'^{\dagger}_{i-1}}$ actually means $\alpha_i \in {}_\mathbb{R}\Delta_{R'}^{R', R'^\dagger}$. But since γ is a root of $\mathfrak{l}_{P,\ell}$, $\mathbf{c}_\gamma(\gamma)$ is a root of $\mathfrak{l}_{R,\ell}$ and hence a linear combination of roots in ${}_\mathbb{R}\Delta_{R'}^{R^\dagger} = {}_\mathbb{R}\Delta_{R'}^{R'^\dagger}$ and so can't be proportional to α_i. □

25.4. **Corollary.** *Let P be a parabolic $\mathbb{Q}$-subgroup and let $D \subseteq {}_\mathbb{R}D^*_\sigma$ be a Satake compactification for which we assume all real boundary components $D_{R',h} \geq D_{P^\dagger,h}$ are equal-rank. Let V be an irreducible constituent of $H^i(\mathfrak{n}_P; E)$ for which $V|_{M_P}$ is conjugate self-contragredient. Then*

$$(\xi_V + \rho)|_{\mathfrak{a}_P} \in -\mathfrak{a}_P^{*+} \implies i \geq \frac{1}{2}(\dim \mathfrak{n}_P + \dim \mathfrak{a}_P^G + \dim D_{P,\ell}(V))$$

and

$$(\xi_V + \rho)|_{\mathfrak{a}_P} \in \mathfrak{a}_P^{*+} \implies i \leq \frac{1}{2}(\dim \mathfrak{n}_P - \dim \mathfrak{a}_P^G - \dim D_{P,\ell}(V)).$$

Proof. Combine Lemma 25.2 and Corollary 24.3. □

Remark. For the Baily-Borel-Satake compactification in the Hermitian case, the second inequality (for ξ_V in a larger cone) is the content of [47, Prop. 11.1]. Shortly after [47] appeared, Saper and Stern noted that the result generalized to Satake compactifications of equal-rank symmetric spaces where all real boundary components were equal-rank—a proof replaced Lemmas 11.6 and 11.7 from [47] with a case by case analysis based on the list appearing in [58, (A2)]. The proof here via Corollary 24.3 and Lemma 25.2 is independent of classification theory.

26. Restriction to Fibers of $\pi\colon \widehat{X} \to X^*_\sigma$

26.1**. Theorem.** *Let F_R be a stratum of a Satake compactification X^*_σ and assume that all the real boundary components $D_{R',h} \geq D_{R,h}$ of the associated Satake compactification ${}_{\mathbb{R}}D^*_\sigma$ are equal-rank. Let $x \in F_R$ and consider the inclusions $\hat\imath_{R,\ell}\colon \pi^{-1}(x) = \widehat{X}_{R,\ell} \hookrightarrow \widehat{X}$ and $\hat\imath_R\colon \pi^{-1}(F_R) = X_R(L_{R,\ell}) \hookrightarrow \widehat{X}$. Let $\mathcal{M}$ be an $\mathcal{L}_{\widehat{X}}$-module. Then*

$$\mathrm{SS}(\hat\imath^*_{R,\ell}\mathcal{M}) \subseteq \operatorname{Res}^*_\ell \hat\imath^*_R \,\mathrm{SS}(\mathcal{M}) \qquad \text{and} \qquad \mathrm{SS}(\hat\imath^!_{R,\ell}\mathcal{M}) \subseteq \operatorname{Res}^!_\ell \hat\imath^!_R \,\mathrm{SS}(\mathcal{M}) \ . \tag{26.1.1}$$

We have the estimates

$$\begin{aligned} d(\hat\imath^*_{R,\ell}\mathcal{M}) \leq \sup_{\substack{\tilde V \in \mathrm{SS}(\mathcal{M}) \\ F_{\tilde R} \geq F_R}} \big(d(\tilde V;\mathcal{M}) &+ \frac{1}{2}(\dim D_{\tilde P,\ell} + \dim D_{\tilde P,\ell}(\tilde V)) \\ &+ \frac{1}{2}\operatorname{codim}_{F^*_{\tilde R}} F_R - \dim \mathfrak{a}^{\tilde P}_{\tilde P \cap R} \big) \end{aligned} \tag{26.1.2}$$

and

$$\begin{aligned} c(\hat\imath^!_{R,\ell}\mathcal{M}) \geq \inf_{\substack{\tilde V \in \mathrm{SS}(\mathcal{M}) \\ F_{\tilde R} \geq F_R}} \big(c(\tilde V;\mathcal{M}) &- \frac{1}{2}(\dim D_{\tilde P,\ell} - \dim D_{\tilde P,\ell}(\tilde V)) \\ &+ \frac{1}{2}\operatorname{codim}_{F^*_{\tilde R}} F_R + \dim \mathfrak{a}^{\tilde P}_{\tilde P \cap R} \big). \end{aligned} \tag{26.1.3}$$

In these estimates, $\tilde V$ is an irreducible $L_{\tilde P}$-module and we set $\tilde R = \tilde P^\dagger$.

Proof. Equation (26.1.1) is simply Propositions 23.3 and 22.8. These propositions further imply that if $V_\ell \in \mathrm{SS}(\hat\imath^*_{R,\ell}\mathcal{M})$, then there exists $V \in \mathrm{SS}(\hat\imath^*_R\mathcal{M})$ and $\tilde V \in \mathrm{SS}(\mathcal{M})$ such that $\operatorname{Res}^*_\ell V = V_\ell$ and there exists

$$V = V_0 \preccurlyeq_+ \cdots \preccurlyeq_+ V_N = \tilde V$$

satisfying (23.3.1)–(23.3.3). Furthermore, as in the proof of Lemma 23.2, we have an almost direct product factorization

$$L_{P_i} = \widetilde{L_{P_i,h}} L_{P_i,\ell} = \widetilde{L_{P_i,h}} L^{P_{i+1},h}_{P_i,\ell} L_{P_{i+1},\ell} \ .$$

Let $D^{P_{i+1},h}_{P_i,\ell}$ denote the symmetric space associated to $L^{P_{i+1},h}_{P_i,\ell}$. Thus we can apply Corollary 25.4 to the parabolic $(P_i/N_{P_{i+1}}) \cap L_{P_{i+1},h} \subseteq L_{P_{i+1},h}$ and the induced Satake compactification of $D_{P_{i+1},h}$ in order to conclude that

$$[V_{i+1} : V_i] \leq \frac{1}{2}(\dim \mathfrak{n}^{P_{i+1}}_{P_i} - \dim \mathfrak{a}^{P_{i+1}}_{P_i} - \dim D^{P_{i+1},h}_{P_i,\ell}(V_i)) \tag{26.1.4}$$

Now $\operatorname{Res}^{L_{P_i}}_{L_{P_{i+1},\ell}}(V_i)$ is a direct sum of copies of $V_{P_{i+1},\ell}$, so $D_{P_i,\ell}(V_i) = D^{P_{i+1},h}_{P_i,\ell}(V_i) \times D_{P_{i+1},\ell}(V_{i+1})$. Consequently if we sum (26.1.4) for $i = 0, \dots, N-1$ we obtain

$$
\begin{aligned}
[\tilde V : V] &\le \frac{1}{2}(\dim \mathfrak{n}_P^{\tilde P} - \dim \mathfrak{a}_P^{\tilde P} - \dim D^{\tilde P,h}_{P,\ell}(V)) \\
&= -\frac{1}{2}(\dim D^{\tilde P,h}_{P,\ell} + \dim D^{\tilde P,h}_{P,\ell}(V)) + \frac{1}{2}\operatorname{codim}_{F^*_{\tilde R}} F_R - \dim \mathfrak{a}_P^{\tilde P}
\end{aligned}
\tag{26.1.5}
$$

since

$$
\operatorname{codim}_{F^*_{\tilde R}} F_R = \dim \mathfrak{n}_P^{\tilde P} + \dim \mathfrak{a}_P^{\tilde P} + \dim D^{\tilde P,h}_{P,\ell}.
$$

We now calculate that (26.1.2) holds by using (10.3.2), (22.6.2), (22.8.2), and (26.1.5). The proof of (26.1.3) is similar. □

26.2. **Corollary.** *In the above situation, assume further that* $\mathrm{SS}_{\mathrm{ess}}(\mathcal{M}) = \{E\}$ *for* $E \in \mathfrak{Irr}(G)$ *with* $c(E;\mathcal{M}) = d(E;\mathcal{M}) = 0$. *Then*

$$
d(\hat\imath^*_{R,\ell}\mathcal{M}) \le \frac{1}{2}\operatorname{codim} F_R - \dim \mathfrak{a}_R^G \qquad \text{and} \qquad c(\hat\imath^!_{R,\ell}\mathcal{M}) \ge \frac{1}{2}\operatorname{codim} F_R + \dim \mathfrak{a}_R^G.
$$

Proof. Any $V \in \mathrm{SS}(\mathcal{M})$ satisfies $V \preccurlyeq_0 E$ by Corollary 9.3 and therefore has the form $H^{\ell(w)}(\mathfrak{n}_P; E)_w$ where w is fundamental by Lemma 9.1. However since D is equal-rank the only fundamental parabolic $\mathbb{R}$-subgroup is G. Hence $\mathrm{SS}(\mathcal{M}) = \{E\}$. Now apply (26.1.2) and (26.1.3). □

27. APPLICATION: THE CONJECTURE OF RAPOPORT AND GORESKY-MACPHERSON

27.1. **Theorem.** *Let* X^* *be a Satake compactification of* X *for which all real boundary components* $D_{R,h}$ *are equal-rank, and let* $\pi\colon \widehat X \to X^*$ *be the projection from the reductive Borel-Serre compactification. Let* p *be a middle perversity. Then there is a natural quasi-isomorphism* $\pi_*\mathcal{I}_p\mathcal{C}(\widehat X;\mathbb{E}) \cong \mathcal{I}_p\mathcal{C}(X^*;\mathbb{E})$.

Remark. When D is Hermitian and X^* is the Baily-Borel-Satake compactification, the equal-rank hypothesis is automatic. In this case the theorem was conjectured independently by Rapoport [39] and Goresky and MacPherson [24]; a proof in this case for $\mathbb{Q}$-rank $G = 1$ was given by Saper and Stern [40, Appendix].

Proof. For $x \in F_R$, a proper stratum of X^*, let $i_x\colon \{x\} \hookrightarrow X^*$ and $\hat\imath_{R,\ell}\colon \pi^{-1}(x) = \widehat X_{R,\ell} \hookrightarrow \widehat X$ be the inclusion maps. Let $q(k) = k - 2 - p(k)$ be the dual middle perversity. By the local characterization of intersection cohomology [23], [10, V, 4.2] on X^* we need to verify that

(i) $\pi_*\mathcal{I}_p\mathcal{C}(\widehat X;\mathbb{E})$ is $\mathcal{X}$-clc,
(ii) $\pi_*\mathcal{I}_p\mathcal{C}(\widehat X;\mathbb{E})|_X$ is quasi-isomorphic to $\mathbb{E}$,
(iii) $H^i(i_x^*\pi_*\mathcal{I}_p\mathcal{C}(\widehat X;\mathbb{E})) = 0$ for $x \in F_R$, $i > p(\operatorname{codim} F_R)$, and
(iv) $H^i(i_x^!\pi_*\mathcal{I}_p\mathcal{C}(\widehat X;\mathbb{E})) = 0$ for $x \in F_R$, $i < \dim X - q(\operatorname{codim} F_R)$.

Condition (i) follows from [10, V, 10.16] and condition (ii) is obvious. For the others, calculate

$$
\begin{aligned}
H(i_x^*\pi_*\mathcal{I}_p\mathcal{C}(\widehat X;\mathbb{E})) &\cong H(\widehat X_{R,\ell}; \hat\imath^*_{R,\ell}\mathcal{I}_p\mathcal{C}(\widehat X;\mathbb{E})) && \text{by [10, V, 10.7],} \\
&\cong H(\widehat X_{R,\ell}; \hat\imath^*_{R,\ell}\mathcal{I}_p\mathcal{C}(E)) && \text{by Thm. 4.1 and Prop. 5.4}
\end{aligned}
$$

and similarly $H(i_x^* \pi_* \mathcal{I}_p\mathcal{C}(\widehat{X};\mathbb{E})) \cong H(\widehat{X}_{R,\ell}; \hat{\imath}^!_{R,\ell}\mathcal{I}_p\mathcal{C}(E))$. Now since all strata of X^* have even dimension by the equal-rank hypothesis, $p(\operatorname{codim} F_R) = \frac{1}{2}\operatorname{codim} F_R - 1$ and $\dim X - q(\operatorname{codim} F_R) = \frac{1}{2}\operatorname{codim} F_R + 1$. Thus we need to demonstrate that

$$\begin{aligned} H^i(\widehat{X}_{R,\ell}; \hat{\imath}^*_{R,\ell}\mathcal{I}_p\mathcal{C}(E)) &= 0 \qquad \text{for } i \geq \frac{1}{2}\operatorname{codim} F_R, \\ H^i(\widehat{X}_{R,\ell}; \hat{\imath}^!_{R,\ell}\mathcal{I}_p\mathcal{C}(E)) &= 0 \qquad \text{for } i \leq \frac{1}{2}\operatorname{codim} F_R. \end{aligned} \tag{27.1.1}$$

Under the equal-rank hypotheses, the possible types of irreducible $\mathbb{Q}$-root systems that can occur in G are B_n, BC_n, C_n, and G_2 [58, (A.2)]. Thus we can apply Corollary 17.2 to see that the essential micro-support of $\mathcal{I}_p\mathcal{C}(E)$ is simply $\{E\}$; Corollary 26.2 then implies that $d(\hat{\imath}^*_{R,\ell}\mathcal{M}) < \frac{1}{2}\operatorname{codim} F_R$ and $c(\hat{\imath}^!_{R,\ell}\mathcal{M}) > \frac{1}{2}\operatorname{codim} F_R$. The proof (27.1.1) is concluded by applying our vanishing theorem for the cohomology of $\mathcal{L}$-modules, Theorem 10.4. □

27.2. If we replace the use of Corollary 17.2 by Theorem 16.3 in the preceding proof we obtain the following theorem.

Theorem. *Let X^* be a Satake compactification of X for which all real boundary components $D_{R,h}$ are equal-rank, and let $\pi\colon \widehat{X} \to X^*$ be the projection from the reductive Borel-Serre compactification. Let η be a middle weight profile and let p be a middle perversity. Then there is a natural quasi-isomorphism $\pi_*\mathcal{W}^\eta\mathcal{C}(\widehat{X};\mathbb{E}) \cong \mathcal{I}_p\mathcal{C}(X^*;\mathbb{E})$.*

Remark. When D is Hermitian and X^* is the Baily-Borel-Satake compactification, this theorem was the main result of [22].

References

1. J. Arthur, *A trace formula for reductive groups. I. Terms associated to classes in $G(\mathbf{Q})$*, Duke Math. J. **45** (1978), 911–953.
2. W. Baily and A. Borel, *Compactification of arithmetic quotients of bounded symmetric domains*, Ann. of Math. **84** (1966), 442–528.
3. W. Ballmann, *Lectures on spaces of nonpositive curvature*, Birkhäuser, Basel, 1995.
4. A. Beilinson, J. Bernstein, and P. Deligne, *Faisceaux pervers*, Analyse et topologie sur les espaces singuliers (I), CIRM, 6–10 juillet 1981, Astérisque, vol. 100, 1982.
5. A. Borel, *Ensembles fondamentaux pour les groupes arithmétiques*, Colloque sur la Théorie des Groupes Algébriques, Tenu à Bruxelles du 5 au 7 juin 1962, Centre Belge de Recherches Mathématiques, 1962, pp. 23–40.
6. ———, *Stable real cohomology of arithmetic groups*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 235–272.
7. ———, *A vanishing theorem in relative Lie algebra cohomology*, Algebraic Groups: Utrecht 1986 (Berlin) (A. M. Cohen, W. H. Hesselink, W. L. J. van der Kallen, and J. R. Strooker, eds.), Lecture Notes in Mathematics, vol. 1271, Springer-Verlag, 1987, pp. 1–16.
8. ———, *Linear algebraic groups*, Springer-Verlag, Berlin, 1991.
9. A. Borel and W. Casselman, *L^2-cohomology of locally symmetric manifolds of finite volume*, Duke Math. J. **50** (1983), 625–647.
10. A. Borel et al., *Intersection cohomology*, Birkhäuser, Boston, 1984.
11. A. Borel and J.-P. Serre, *Corners and arithmetic groups*, Comment. Math. Helv. **48** (1973), 436–491.
12. A. Borel and J. Tits, *Groupes réductifs*, I.H.E.S. Publ. Math. **27** (1965), 55–151.
13. A. Borel and N. Wallach, *Continuous cohomology, discrete subgroups, and representation of reductive groups*, Ann. of Math. Stud., vol. 94, Princeton University Press, Princeton, 1980.
14. N. Bourbaki, *Groupes et algèbres de Lie, chapitres 4, 5 et 6*, Éléments de mathématique, Masson, Paris, 1981.

15. P. Cartier, *Remarks on "Lie algebra cohomology and the generalized Borel-Weil theorem", by B. Kostant*, Ann. of Math. **74** (1961), 388–390.
16. W. Casselman, *L^2-cohomology for groups of real rank one*, Representation Theory of Reductive Groups (P. Trombi, ed.), Birkhäuser, Boston, 1983, pp. 69–84.
17. ———, *Geometric rationality of Satake compactifications*, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., **9**, Cambridge Univ. Press, Cambridge, 1997, pp. 81–103.
18. J. Cheeger, *On the Hodge theory of Riemannian pseudomanifolds*, Geometry of the Laplace Operator (Providence), Proc. Sympos. Pure Math., vol. 36, Amer. Math. Soc., 1980, pp. 91–146.
19. G. de Rham, *Variétés différentiables*, Hermann, Paris, 1973.
20. J. Franke, *Harmonic analysis in weighted L_2-spaces*, Ann. Sci. École Norm. Sup. (4) **31** (1998), 181–279.
21. M. Gaffney, *A special Stokes' theorem for complete Riemannian manifolds*, Ann. of Math. **60** (1954), 140–145.
22. M. Goresky, G. Harder, and R. MacPherson, *Weighted cohomology*, Invent. Math. **116** (1994), 139–213.
23. M. Goresky and R. MacPherson, *Intersection homology II*, Invent. Math. **72** (1983), 77–129.
24. ———, *Weighted cohomology of Satake compactifications*, Centre de recherches mathématiques, preprint #1593, 1988.
25. ———, *The topological trace formula*, J. Reine Angew. Math. **560** (2003), 77–150.
26. L. Hörmander, *L^2 estimates and existence theorems for the $\overline{\partial}$ operator*, Acta Math. **113** (1965), 89–152.
27. M. Kashiwara and P. Schapira, *Sheaves on manifolds*, Springer-Verlag, Berlin, 1990.
28. G. Kempf, *Linear systems on homogeneous spaces*, Ann. of Math. **103** (1976), 557–591.
29. A. Knapp, *Lie groups, beyond an introduction*, Birkhäuser, Boston, 1996.
30. B. Kostant, *On the conjugacy of real Cartan subalgebras. I.*, Proc. Nat. Acad. Sci. U. S. A. **41** (1955), 967–970.
31. ———, *Lie algebra cohomology and the generalized Borel-Weil theorem*, Ann. of Math. **74** (1961), 329–387.
32. V. Lakshmibai, C. Musili, and C. Seshadri, *Geometry of G/P. III. Standard monomial theory for a quasi-minuscule P*, Proc. Indian Acad. Sci. Sect. A Math. Sci. **88A** (1979), 93–177.
33. R. P. Langlands, *On the functional equations satisfied by Eisenstein series*, Lecture Notes in Mathematics, vol. 544, Springer-Verlag, Berlin, 1976.
34. Y. Matsushima and S. Murakami, *On vector bundle valued harmonic forms and automorphic forms on symmetric Riemannian manifolds*, Ann. of Math. **78** (1963), 365–416.
35. C. Moeglin and J.-L. Waldspurger, *Spectral decomposition and Eisenstein series*, Cambridge University Press, Cambridge, 1995.
36. K. Nomizu, *On the cohomology of compact homogeneous spaces of nilpotent Lie groups*, Ann. of Math. **59** (1954), 531–538.
37. M. S. Raghunathan, *Vanishing theorems for cohomology groups associated to discrete subgroups of semisimple Lie groups*, Osaka J. Math. **3** (1966), 243–256.
38. ———, *Corrections to "Vanishing theorems . . . "*, Osaka J. Math. **16** (1979), 295–299.
39. M. Rapoport, 1986, letter to A. Borel.
40. ———, *On the shape of the contribution of a fixed point on the boundary: The case of $\mathbb{Q}$-rank 1 (with an appendix by L. Saper and M. Stern)*, The Zeta functions of Picard modular surfaces (R. P. Langlands and D. Ramakrishnan, eds.), Les Publications CRM, Montréal, 1992, pp. 479–488.
41. L. Saper, *L_2-cohomology and intersection homology of certain algebraic varieties with isolated singularities*, Inv. Math. **82** (1985), 207–255.
42. ———, *L_2-cohomology of Kähler varieties with isolated singularities*, J. Diff. Geom. **36** (1992), 89–161.
43. ———, *Tilings and finite energy retractions of locally symmetric spaces*, Comment. Math. Helv. **72** (1997), 167–202.
44. ———, *Geometric rationality of equal-rank Satake compactifications*, Math. Res. Lett. **11** (2004), no. 5, 653–671.
45. ———, *L^2-cohomology of locally symmetric spaces, I*, `math.RT/0412353`, submitted, 2004.
46. ———, *$\mathcal{L}$-modules and the conjecture of Rapoport and Goresky-MacPherson*, `math.RT/0112250`, to appear in Astérisque, 2005.

47. L. Saper and M. Stern, *L_2-cohomology of arithmetic varieties*, Ann. of Math. **132** (1990), 1–69.
48. I. Satake, *On a generalization of the notion of a manifold*, Proc. Nat. Acad. Sci. U. S. A. **42** (1956), 359–363.
49. ———, *On representations and compactifications of symmetric Riemannian spaces*, Ann. of Math. **71** (1960), 77–110.
50. ———, *On compactifications of the quotient spaces for arithmetically defined discontinuous groups*, Ann. of Math. **72** (1960), 555–580.
51. J. Schwermer, *Eisenstein series and cohomology of arithmetic groups: the generic case*, Invent. Math. **116** (1994), 481–511.
52. M. Sugiura, *Conjugate classes of Cartan subalgebras in real semisimple Lie algebras*, J. Math. Soc. Japan **11** (1959), 374–434.
53. ———, *Correction to my paper: "Conjugate classes of Cartan subalgebras in real semisimple Lie algebras"*, J. Math. Soc. Japan **23** (1971), 379–383.
54. W. Thurston, *The geometry and topology of three-manifolds*, M.S.R.I., `http://www.msri.org/publications/books/gt3m/`, 1980.
55. W. T. van Est, *A generalization of the Cartan-Leray sequence, I, II*, Indag. Math. **20** (1958), 399–413.
56. S. Zucker, *L_2 cohomology of warped products and arithmetic groups*, Invent. Math. **70** (1982), 169–218.
57. ———, *Satake compactifications*, Comment. Math. Helv. **58** (1983), 312–343.
58. ———, *L_2-cohomology and intersection homology of locally symmetric varieties, II*, Compositio Math. **59** (1986), 339–398.

Department of Mathematics, Duke University, Box 90320, Durham, NC 27708, U.S.A.
E-mail address: `saper@math.duke.edu`
URL: `http://www.math.duke.edu/faculty/saper`